

OPTIMAL HOMOGENIZATION RATES IN STOCHASTIC HOMOGENIZATION OF NONLINEAR UNIFORMLY ELLIPTIC EQUATIONS AND SYSTEMS

JULIAN FISCHER AND STEFAN NEUKAMM

ABSTRACT. We derive optimal-order homogenization rates for random nonlinear elliptic PDEs with monotone nonlinearity in the uniformly elliptic case. More precisely, for a random monotone operator on \mathbb{R}^d with stationary law (i. e. spatially homogeneous statistics) and fast decay of correlations on scales larger than the microscale $\varepsilon > 0$, we establish homogenization error estimates of the order ε in case $d \geq 3$, respectively of the order $\varepsilon |\log \varepsilon|^{1/2}$ in case $d = 2$. Previous results in nonlinear stochastic homogenization have been limited to a small algebraic rate of convergence ε^δ . We also establish error estimates for the approximation of the homogenized operator by the method of representative volumes of the order $(L/\varepsilon)^{-d/2}$ for a representative volume of size L . Our results also hold in the case of systems for which a (small-scale) $C^{1,\alpha}$ regularity theory is available.

CONTENTS

1. Introduction	2
1.1. Summary of results	4
1.2. Examples	6
1.3. Notation	7
2. Main results	8
2.1. Assumptions and setting	10
2.2. Optimal-order homogenization error estimates	12
2.3. Optimal-order error estimates for the approximation of the homogenized operator by periodic RVEs	14
2.4. The decorrelation assumption: spectral gap inequality	17
3. Strategy and intermediate results	18
3.1. Key objects: Localized correctors, the two-scale expansion, and linearized correctors	18
3.2. The strategy of proof for corrector estimates	25
3.3. Quantitative two-scale approximation	34
3.4. The approximation of the homogenized operator by periodic representative volumes	36
4. Proof of the main results	39
4.1. Ingredients from regularity theory	39
4.2. The convergence rate of the solutions	41

Key words and phrases. random material, stochastic homogenization, convergence rate, representative volume, nonlinear elliptic equation, monotone operator.

SN acknowledges partial support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – project number 405009441.

4.3. Estimates on the random fluctuations of the RVE approximation for the effective material law	48
4.4. Estimates for the error introduced by localization	49
4.5. Coupling error for RVEs	55
4.6. Estimate on the systematic error of the RVE method	58
5. Corrector estimates for the nonlinear PDE	61
5.1. Estimates on linear functionals of the corrector and the flux corrector	61
5.2. Estimates on the corrector for the monotone system	68
6. Corrector estimates for the linearized system	73
6.1. Estimates on linear functionals of the corrector and the flux corrector for the linearized PDE	73
6.2. Estimates on the corrector	80
7. Structural properties of the effective equation	81
Appendix A. Auxiliary results from regularity theory	83
Appendix B. Qualitative differentiability of correctors	86
Appendix C. Meyers' estimate for elliptic equations with massive term	87
References	90

1. INTRODUCTION

In the present work, we establish quantitative homogenization results with optimal rates for nonlinear elliptic PDEs of the form

$$(1) \quad -\nabla \cdot (A_\varepsilon(x, \nabla u_\varepsilon)) = f \quad \text{in } \mathbb{R}^d,$$

where A_ε is a random monotone operator whose correlations decay quickly on scales larger than a microscopic scale ε . For scalar problems and also certain systems, we obtain the optimal convergence rate $O(\varepsilon)$ of the solutions u_ε towards the solution u_{hom} of the homogenized problem

$$(2) \quad -\nabla \cdot (A_{\text{hom}}(\nabla u_{\text{hom}})) = f \quad \text{in } \mathbb{R}^d$$

in three or more spatial dimensions $d \geq 3$. In two spatial dimensions $d = 2$, we obtain the optimal convergence rate $O(\varepsilon |\log \varepsilon|^{1/2})$ upon including a lower-order term in the PDEs (1) and (2).

Our results may be seen as the optimal quantitative counterpart in the case of 2-growth to the qualitative stochastic homogenization theory for monotone systems developed by Dal Maso and Modica [17, 18], respectively as the nonlinear counterpart of the optimal-order stochastic homogenization theory for linear elliptic equations developed by Gloria and Otto [32, 33] and Gloria, Otto, and the second author [29, 31]. Just like for [17, 18], a key motivation for our work is the homogenization of nonlinear materials.

In the context of random materials, the first – and to date also the only – homogenization rates for elliptic PDEs with monotone nonlinearity were obtained by Armstrong and Smart [9], Armstrong and Mourrat [7], and Armstrong, Ferguson, and Kuusi [2] in the form of a small algebraic convergence rate ε^δ for some $\delta > 0$. The optimal convergence rates derived in the present work improve substantially upon their rate. However, in contrast to the works of Armstrong et al. we make no attempt to reach optimal stochastic integrability: While we derive an error estimate

of the form

$$(3) \quad \|u_\varepsilon - u_{\text{hom}}\|_{L^p(\mathbb{R}^d)} \leq \begin{cases} \mathcal{C}_f \varepsilon |\log \varepsilon|^{1/2} & \text{for } d = 2, \\ \mathcal{C}_f \varepsilon & \text{for } d \geq 3 \end{cases}$$

with a random constant \mathcal{C}_f whose stretched exponential moments are estimated by $\mathbb{E}[\exp(|\mathcal{C}_f/C(f)|^\nu)] \leq 2$ for some constant $\nu > 0$ and some constant $C(f) > 0$, the homogenization error estimates for linear elliptic PDEs with optimal rate in [4, 34] establish (essentially) Gaussian stochastic moments $\mathbb{E}[\exp(|\mathcal{C}_f/C(f, \mu)|^{2-\mu})] \leq 2$ for any $\mu > 0$. Likewise, the homogenization error estimates for monotone operators with non-optimal rate ε^δ of [2, 7, 9] include optimal stochastic moment bounds.

Before providing a more detailed summary of our results, let us give a brief overview of the previous quantitative results in nonlinear stochastic homogenization. The first – logarithmic – rates of convergence in the stochastic homogenization of a nonlinear second-order elliptic PDE were obtained by Caffarelli and Souganidis [15] in the setting of non-divergence form equations. Subsequently, a rate of convergence ε^δ has been derived both for equations in divergence form and non-divergence form by Armstrong and Smart [9, 8] and Armstrong and Mourrat [7]. In the homogenization of Hamilton-Jacobi equations, a rate of convergence of the order $\varepsilon^{1/8}$ has been obtained by Armstrong, Cardaliaguet, and Souganidis [6]. For forced mean curvature flow, Armstrong and Cardaliaguet [1] have derived a convergence rate of order $\varepsilon^{1/90}$. These rates of convergence are all expected to be non-optimal (compare, for instance, the result for Hamilton-Jacobi equations to the rate of convergence ε known in the periodic homogenization setting [37]).

To the best of our knowledge, the present work constitutes the first optimal-order convergence results for any nonlinear stochastic homogenization problem. However, we are aware of an independent work in preparation by Armstrong, Ferguson, and Kuusi [3], which aims to address the same question. In contrast to our work – which is inspired by the approach to quantitative stochastic homogenization via spectral gap inequalities of [32, 33, 29, 31] – the upcoming work [3] relies on the approach of sub- and superadditive quantities of [9, 4, 2]. Nevertheless, both our present work and the approach of [3, 2] use the concept of correctors for the linearized PDE, see Section 3 for details.

Before turning to a more detailed description of our results, let us briefly comment on the theory of periodic homogenization of nonlinear elliptic equations. A quantitative theory for the periodic homogenization of nonlinear monotone operators has recently been derived by Wang, Xu, and Zhao [42]. A corresponding result for degenerate elliptic equations of p -Laplacian type may be found in [43]. In the periodic homogenization of polyconvex integral functionals, the single-cell formula for the effective material law (which determines the effective material law by a variational problem on a single periodicity cell) may fail [12, 26], a phenomenon associated with possible “buckling” of the microstructure. A related phenomenon of loss of ellipticity may occur in the periodic homogenization of linear elasticity [14, 24, 35, 36]. Note that polyconvex integral functionals occur naturally in the framework of nonlinear elasticity [11]; however, their Euler-Lagrange equations in general lack a monotone structure. However, in periodic homogenization of nonlinear elasticity the single-cell formula is valid for small deformations [39, 40], and rates of convergence may be derived.

1.1. Summary of results. To summarize our results in a continuum mechanical language, we consider the effective macroscopic behavior of a nonlinear and microscopically heterogeneous material. We assume that the behavior of the nonlinear material is described by the solution $u_\varepsilon : \mathbb{R}^d \rightarrow \mathbb{R}^m$ of a second-order nonlinear elliptic system of the form

$$-\nabla \cdot (A_\varepsilon(x, \nabla u_\varepsilon)) = f$$

for some random monotone operator $A_\varepsilon : \mathbb{R}^d \times \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ with correlation length ε and some right-hand side $f \in L^{\frac{d}{d+2}}(\mathbb{R}^d; \mathbb{R}^m)$. We further assume that the random monotone operator A_ε is of the form $A_\varepsilon(x, \xi) := A(\omega_\varepsilon(x), \xi)$, where ω_ε is a random field representing the random heterogeneities in the material; for each realization of the random material (i.e. each realization of the probability distribution), ω_ε selects at each point $x \in \mathbb{R}^d$ a local material law $A(\omega_\varepsilon(x), \cdot) : \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ from a family A of potential material laws. Under some suitable additional conditions, the theory of stochastic homogenization shows that for small correlation lengths $\varepsilon \ll 1$ the above nonlinear elliptic system is well-approximated by a homogenized effective equation. The effective equation again takes the form of a nonlinear elliptic system, however now with a spatially homogeneous effective material law $A_{\text{hom}} : \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$. It is our goal to provide an optimal-order estimate for the difference of the solution u_ε to the solution u_{hom} of the effective equation

$$-\nabla \cdot (A_{\text{hom}}(\nabla u_{\text{hom}})) = f,$$

as well as to give an optimal-order error bound for the approximation of the effective material law A_{hom} by the method of representative volumes.

To be mathematically more precise, we consider a random field taking values in the unit ball of a Hilbert space $\omega_\varepsilon : \Omega \times \mathbb{R}^d \rightarrow \mathbb{H} \cap B_1$ and a family of monotone operators $A : (\mathbb{H} \cap B_1) \times \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ indexed by the unit ball of this Hilbert space. We then define a random monotone operator $A_\varepsilon(x, \cdot) := A(\omega_\varepsilon(x), \cdot)$ by selecting a monotone operator from the family A at each point $x \in \mathbb{R}^d$ according to the value of the random field $\omega_\varepsilon(x)$. Note that the property of ω_ε being Hilbert-space valued is not an essential point and just included for generality: Even the homogenization for a scalar-valued random field (and correspondingly a single-parameter family of monotone operators $A : (\mathbb{R} \cap B_1) \times \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$) would be highly relevant and just as difficult, as it could describe e.g. composite materials.

The conditions on the random field ω_ε and the family of monotone operators A are as follows:

- We assume *spatial statistical homogeneity of the material*: The statistics of the random material should not depend on the position in space. In terms of a mathematical formulation, this assumption corresponds to stationarity of the probability distribution of ω_ε under spatial translations.
- We assume sufficiently fast *decorrelation of the material properties on scales larger than a correlation length ε* . In terms of a mathematical formulation, we make this notion rigorous by assuming that a spectral gap inequality holds. More precisely, we shall assume that ω_ε itself is a Hilbert-space valued random field on \mathbb{R}^d which satisfies a spectral gap inequality and on which the random monotone operator A_ε depends in a pointwise way as $A_\varepsilon(x, \xi) := A(\omega_\varepsilon(x), \xi)$, where the map $A : (\mathbb{H} \cap B_1) \times \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ is Lipschitz in its first variable.

- We assume *uniform coercivity and boundedness* of the monotone operator in the sense that $(A(\omega, \xi_2) - A(\omega(x), \xi_1)) : (\xi_2 - \xi_1) \geq \lambda |\xi_2 - \xi_1|^2$ and $|A(\omega, \xi_2) - A(\omega, \xi_1)| \leq \Lambda |\xi_2 - \xi_1|$ hold for all $\xi_1, \xi_2 \in \mathbb{R}^{m \times d}$ and every $\omega \in \mathbb{H} \cap B_1$ for suitably chosen constants $0 < \lambda < \Lambda < \infty$.
- For some of our results, we shall impose an additional condition, which essentially entails a $C^{1,\alpha}$ regularity theory for the equation (1) on the microscopic scale ε . Namely, we shall assume Lipschitz continuity of the random field ω_ε on the ε -scale with suitable stochastic moment bounds on the local Lipschitz norm and a uniform bound on the second derivative $\partial_\xi^2 A$, along with one of the following three conditions:
 - Our problem consists of a single nonlinear monotone PDE, i. e. $m = 1$.
 - We are in the two-dimensional case $d = 2$.
 - Our system has Uhlenbeck structure, i. e. the nonlinearity has the structure $A(\omega, \xi) = \rho(\omega, |\xi|^2)\xi$ for some scalar function ρ , and the same is true for the homogenized operator.

Under these assumptions, we establish the following quantitative stochastic homogenization results with optimal rates for the nonlinear elliptic PDE (1).

- The solution u_ε to the nonlinear PDE with fluctuating random material law (1) can be approximated by the solution u_{hom} to a homogenized effective PDE of the form (2). In case $d = 2$ or $d = 1$, we include a lower-order term in the PDEs, see (6) and (7). The homogenized effective material law is given by a monotone operator $A_{\text{hom}} : \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ which is independent of the spatial variable $x \in \mathbb{R}^d$ and satisfies analogous uniform ellipticity and boundedness properties. The error $u - u_{\text{hom}}$ is estimated by

$$\|u - u_{\text{hom}}\|_{L^p(\mathbb{R}^d)} \leq \begin{cases} \mathcal{C}(f) \varepsilon & \text{for } d \geq 3, \\ \mathcal{C}(f) \varepsilon |\log \varepsilon|^{1/2} & \text{for } d = 2, \\ \mathcal{C}(f) \varepsilon^{1/2} & \text{for } d = 1, \end{cases}$$

where $p = \frac{2d}{d-2}$ in case $d \geq 2$ and $p = 2$ in case $d \in \{1, 2\}$. Here, $\mathcal{C}(f)$ denotes a random constant with bounded stretched exponential moments in the sense

$$\mathbb{E} \left[\exp \left(\left(\frac{\mathcal{C}(f)}{C(f)} \right)^{\bar{v}} \right) \right] \leq 2$$

for some universal constant $\bar{v} > 0$ (which is in particular independent of the right-hand side f) and for some constant $C(f) = C(\|f\|_{L^1}, \|f\|_{L^{d+1}})$. Without the additional small-scale regularity assumption, we still achieve half of the rate of convergence $\varepsilon^{1/2}$ for $d \geq 3$ respectively $\varepsilon^{1/2} |\log \varepsilon|^{1/4}$ for $d = 2$, a result that we also establish for the Dirichlet problem in bounded domains.

- The homogenized effective operator A_{hom} may be approximated by the method of representative volumes, and this approximation is subject to the following a priori error estimate: If a box of size $L \geq \varepsilon$ is chosen as the representative volume, the error estimate

$$|A_{\text{hom}}^{\text{RVE}}(\xi) - A_{\text{hom}}(\xi)| \leq \mathcal{C}(L, \xi) (1 + |\xi|)^C |\xi| \left(\frac{L}{\varepsilon} \right)^{-d/2}$$

holds true for every $\xi \in \mathbb{R}^{m \times d}$, where $A_{\text{hom}}^{\text{RVE}}$ denotes the approximation of A_{hom} by the method of representative volumes and where again $\mathcal{C}(L, \xi)$ denotes a random constant with bounded stretched exponential moments (independently of L and ξ). The systematic error is of higher order

$$|\mathbb{E}[A_{\text{hom}}^{\text{RVE}}(\xi)] - A_{\text{hom}}(\xi)| \leq C(1 + |\xi|)^C |\xi| \left(\frac{L}{\varepsilon}\right)^{-d} \left|\log \frac{L}{\varepsilon}\right|^{d+2},$$

at least in case $d \leq 4$ (which includes the physically relevant cases $d = 2$ and $d = 3$). Without the additional small-scale regularity assumption, we achieve almost the same overall estimate $|A_{\text{hom}}^{\text{RVE}}(\xi) - A_{\text{hom}}(\xi)| \leq \mathcal{C}(L, \xi) |\xi| (L/\varepsilon)^{-d/2} (\log \frac{L}{\varepsilon})^C$, but not the improved bound for the systematic error.

Note that the rates of convergence $\|u_\varepsilon - u_{\text{hom}}\|_{L^{2d/(d-2)}} \leq C\varepsilon$ in case $d \geq 3$ respectively $\|u_\varepsilon - u_{\text{hom}}\|_{L^2} \leq C\varepsilon |\log \varepsilon|^{1/2}$ in case $d = 2$ coincide with the optimal rate of convergence in the homogenization of linear elliptic PDEs, see e.g. [4, 29, 31, 34]. Similarly, the rate of convergence for the representative volume approximation $|A_{\text{hom}}^{\text{RVE}}(\xi) - A_{\text{hom}}(\xi)| \leq \mathcal{C}(L/\varepsilon)^{-d/2}$ coincides with the corresponding optimal rate for linear elliptic PDEs, as does (essentially) the higher-order convergence rate for the systematic error. As linear elliptic PDEs may be regarded as a particular case of our nonlinear PDE (1), our rates of convergence are optimal.

1.2. Examples. To illustrate our results, let us mention two examples of random nonlinear elliptic PDEs and systems to which our theorems apply, as well as an important class of random fields ω_ε which satisfy our assumptions.

We first give an example for the random field ω_ε . Let $\theta : \mathbb{R}^k \rightarrow \mathbb{R}^k \cap B_1$ be any Lipschitz map taking values only in the unit ball. Let $Y_\varepsilon : \mathbb{R}^d \rightarrow \mathbb{R}^k$ be any stationary Gaussian random field whose correlations decay sufficiently quickly in the sense

$$|\text{Cov}[Y_\varepsilon(x), Y_\varepsilon(y)]| \lesssim \frac{1}{1 + \left(\frac{|x-y|}{\varepsilon}\right)^{d+\delta}}$$

for some $\delta > 0$. Set $\mathbb{H} := \mathbb{R}^k$. Then the random field $\omega_\varepsilon : \mathbb{R}^d \rightarrow \mathbb{H}$ defined by

$$\omega_\varepsilon(x) := \theta(Y_\varepsilon(x))$$

satisfies a spectral gap inequality with correlation length ε in the sense of Definition 11; for a proof see e. g. [20]. As stationarity is immediate, any such ω_ε satisfies our key assumptions on the random field (P1) and (P2) stated in Section 2.1 below. Note in particular that the spectral gap assumption allows for the presence of (sufficiently quickly decaying, namely integrable) long-range correlations. Typical realizations for two such random fields are depicted in Figure 1.

To state the first example of a random monotone operator satisfying our assumptions, consider any two deterministic spatially homogeneous monotone operators $A_1 : \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ and $A_2 : \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ subject to the ellipticity and Lipschitz continuity assumptions (A1) and (A2). Furthermore, consider any random field $\omega_\varepsilon : \mathbb{R}^d \rightarrow [0, 1]$. Then the operator

$$A_\varepsilon(x, \xi) := \omega_\varepsilon(x)A_1(\xi) + (1 - \omega_\varepsilon(x))A_2(\xi)$$




FIGURE 1. Typical realizations of random fields obtained by applying a nonlinear map pointwise to a stationary Gaussian random field with only short-range correlations (left), respectively to a stationary Gaussian random field with barely integrable correlations (right).

satisfies our assumptions (A1)–(A3). Note that this operator corresponds to the PDE

$$-\nabla \cdot \left(\omega_\varepsilon(x) A_1(\nabla u) + (1 - \omega_\varepsilon(x)) A_2(\nabla u) \right) = f.$$

The additional small-scale regularity assumption (R) is satisfied whenever the operators A_1 and A_2 have uniformly bounded second derivatives, the random field ω_ε is regular enough, and one of the three following conditions holds: The equation is scalar ($m = 1$), the spatial dimension is at most two ($d \leq 2$), or both A_1 and A_2 as well as the homogenized operator A_{hom} are of Uhlenbeck structure.

As a second simple example of a monotone operator, for any random field $\omega_\varepsilon : \mathbb{R}^d \rightarrow [0, 1]$ the operator

$$A_\varepsilon(x, \xi) := \frac{1 + |\xi|^2}{1 + (1 + \omega_\varepsilon(x)) |\xi|^2} \xi$$

satisfies our assumptions, possibly with the exception of the additional regularity condition (R). Note that this operator corresponds to the PDE

$$-\nabla \cdot \left(\frac{1 + |\nabla u|^2}{1 + \omega_\varepsilon(x) |\nabla u|^2} \nabla u \right) = f.$$

The additional small-scale regularity assumption – stated in (R) below – is satisfied in the scalar case $m = 1$ as well as in the low-dimensional case $d \leq 2$, provided that the random field ω_ε is sufficiently smooth on the microscopic scale ε .

1.3. Notation. The number of spatial dimensions will be denoted by $d \in \mathbb{N}$. For a measurable function u , we denote by ∇u its (weak) spatial derivative. For a function of two variables $A(\omega, \xi)$, we denote its partial derivatives by $\partial_\omega A$ and $\partial_\xi A$. For a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$, we denote by $\partial_i f$ its partial derivative with respect to the coordinate i . For a matrix-valued function $M : \mathbb{R}^d \rightarrow \mathbb{R}^{m \times d}$, we denote by $\nabla \cdot M$ its divergence with respect to the second index, i. e. $(\nabla \cdot M)_i = \sum_{j=1}^d \partial_j M_{ij}$.

Throughout the paper, we use standard notation for Sobolev spaces. In particular, we denote by $H^1(\mathbb{R}^d)$ the space of all measurable functions $u : \mathbb{R}^d \rightarrow \mathbb{R}$ whose weak spatial derivative ∇u exists and which satisfy $\|u\|_{H^1} := (\int_{\mathbb{R}^d} |u|^2 + |\nabla u|^2 dx)^{1/2} < \infty$. Similarly, we denote by $H^1(\mathbb{R}^d; \mathbb{R}^m)$ the space of \mathbb{R}^m -valued

vector fields with the analogous properties and the analogous norm. For $d \geq 3$, we denote by $\dot{H}^1(\mathbb{R}^d; \mathbb{R}^m)$ the space of all measurable functions u with $\|u\|_{\dot{H}^1} := (\int_{\mathbb{R}^d} |\nabla u|^2 dx)^{1/2} + \|u\|_{L^{2d/(d-2)}} < \infty$. By $H_{\text{loc}}^1(\mathbb{R}^d)$ we denote the space of all measurable functions $u : \mathbb{R}^d \rightarrow \mathbb{R}$ for which all restrictions $u|_{B_r}$ to finite balls ($0 < r < \infty$) belong to $H^1(B_r)$. For a box $[0, L]^d$, we denote by $H_{\text{per}}^1(\mathbb{R}^d)$ the closure in the $H^1([0, L]^d)$ norm of the smooth L -periodic functions. By $H_{\text{uloc}}^1(\mathbb{R}^d)$, we denote the space of measurable functions u whose weak derivative ∇u exists and which satisfy the bound $\|u\|_{H_{\text{uloc}}^1} := \sup_{x \in \mathbb{R}^d} (\int_{B_1(x)} |\nabla u|^2 + |u|^2 d\tilde{x})^{1/2} < \infty$.

In order not to overburden notation, we shall frequently suppress the dependence on the spatial variable x in many expressions, for instance we will write $A(\omega_\varepsilon, \xi)$ or $A(\omega_\varepsilon, \nabla u)$ instead of $A(\omega_\varepsilon(x), \xi)$ respectively $A(\omega_\varepsilon(x), \nabla u(x))$. By an expression like $\partial_\xi A(\omega, \xi)\Xi$, we denote the derivative of A with respect to the second variable at the point (ω, ξ) evaluated in direction Ξ . Similarly, by $\partial_\omega \partial_\xi A(\omega, \xi)\delta\omega\Xi$ we denote the second mixed derivative of A with respect to its two variables at the point (ω, ξ) evaluated in directions $\delta\omega$ and Ξ . We use notation like δF or $\delta\omega_\varepsilon$ to indicate infinitesimal changes (i. e. differentials) of various quantities and functions.

For two numbers $a, b \in \mathbb{R}$, we denote by $a \wedge b$ the minimum of a and b . We write $a \sim b$ to indicate that two constants $a, b \in (0, \infty)$ are of a similar order of magnitude. For a matrix $M \in \mathbb{R}^{m \times d}$, we denote by $|M| := \sum_{i,j} |M_{ij}|^2$ its Frobenius norm. By $\mathbb{R}_{\text{skew}}^{d \times d}$ we denote the set of skew-symmetric matrices of dimension $d \times d$.

By $B_r(x)$ we denote the ball of radius r centered at x . By B_r we will denote the ball of radius r around the origin. For two sets $A, B \subset \mathbb{R}^d$ and a point $x \in \mathbb{R}^d$, we denote by $A + B := \{a + b : A \in A, b \in B\}$ their Minkowski sum, respectively by $x + A$ the translation of A by x . By \mathbb{H} we will denote a Hilbert space; we will denote its unit ball by $\mathbb{H} \cap B_1$.

By C and c we will denote – typically large respectively typically small – non-negative constants, whose precise value may change from occurrence to occurrence but which only depend on a certain set of parameters. For a set M , we denote by $\#M$ the number of its elements.

We write $\omega_\varepsilon \sim \mathbb{P}$ to indicate that a random field ω_ε is distributed according to the probability distribution \mathbb{P} . For two random variables or random fields X and Y , we write $X \sim Y$ to indicate that their laws coincide.

Whenever we use the terms “coefficient field” or “monotone operator”, we shall implicitly assume measurability.

2. MAIN RESULTS

Before stating our main results and the precise setting, let us introduce the key objects in the homogenization of nonlinear elliptic PDEs and systems with monotone nonlinearity. To fix a physical setting, we will here give an outline of the meaning of the objects in the context of electric fields and associated currents. However, a major motivation for the present work – and in particular for the choice to include the case of nonlinear elliptic systems – stems from the homogenization of nonlinear elastic materials. While in this context the monotone structure is lost [11], it may be retained for small deformations [16, 25, 39, 40, 44]. A corresponding result in the context of stochastic homogenization will be established in [21]. The homogenization of monotone operators may also be viewed as a simple but necessary first step towards a possible quantitative homogenization theory for nonlinear elastic materials for larger deformations.

In the context of electric fields and currents, we are concerned with a scalar equation (i. e. $m = 1$); the functions u and u_{hom} in (1) and (2) correspond (up to a sign) to the electric potential in the heterogeneous material respectively in the homogenized picture. Their gradients ∇u respectively ∇u_{hom} are the associated electric fields. The monotone functions $A(\omega_\varepsilon(x), \xi)$ respectively $A_{\text{hom}}(\xi)$ are the material law and describe the electric current created by a given electric field ξ . Note that in contrast to existing optimal results in stochastic homogenization, the material law may be nonlinear in ξ . Finally, the PDEs (1) and (2) correspond to prescribing the sources and sinks of the electric current.

The central object in the quantitative homogenization of elliptic PDEs is the *homogenization corrector* ϕ_ξ , which in our context is an \mathbb{R}^m -valued random field on \mathbb{R}^d . It provides a bridge between the microscopic (heterogeneous) and the macroscopic (homogenized) picture: For a given constant macroscopic field gradient $\xi \in \mathbb{R}^{m \times d}$, the corrector ϕ_ξ provides the correction yielding the associated microscopic field gradient $\xi + \nabla \phi_\xi$. The corrector ϕ_ξ is defined as the unique (up to a constant) sublinearly growing distributional solution to the PDE

$$(4) \quad -\nabla \cdot (A(\omega_\varepsilon, \xi + \nabla \phi_\xi)) = 0,$$

see Definition 1 for the precise definition. Note that *a priori*, similarly to the linear elliptic case, the existence and uniqueness of such a solution is unclear.

The effective (homogenized) material law A_{hom} (see Definition 1 for the precise definition) may be determined in terms of the homogenization corrector: In principle, at each point $x \in \mathbb{R}^d$ the microscopic material law $A(\omega_\varepsilon(x), \cdot) : \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ associates a current $A(\omega_\varepsilon(x), \xi_\varepsilon)$ to a given electric field ξ_ε ; likewise, the effective macroscopic material law $A_{\text{hom}}(\cdot) : \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ associates a current $A_{\text{hom}}(\xi)$ to a given macroscopic electric field ξ . As the macroscopic current corresponds to an “averaged” microscopic current, the macroscopic material law should be obtained by averaging the microscopic flux. More precisely, the homogenization corrector ϕ_ξ associates a microscopic electric field $\xi + \nabla \phi_\xi$ to a given macroscopic electric field ξ ; therefore the macroscopic current $A_{\text{hom}}(\xi)$ should be given by the “average” of the microscopic current $A(\omega_\varepsilon, \xi + \nabla \phi_\xi)$. In our setting, due to stationarity and ergodicity “averaging” corresponds to taking the expected value at an arbitrary point $x \in \mathbb{R}^d$ (and we will suppress the point $x \in \mathbb{R}^d$ in the notation). In other words, we have

$$A_{\text{hom}}(\xi) = \mathbb{E}[A(\omega_\varepsilon, \xi + \nabla \phi_\xi)] \stackrel{a.s.}{=} \lim_{r \rightarrow \infty} \int_{B_r} A(\omega_\varepsilon(x), \xi + \nabla \phi_\xi(x)) dx.$$

Our main results on the quantitative approximation of the solution u_ε to the nonlinear elliptic PDE with randomly fluctuating material law

$$-\nabla \cdot (A(\omega_\varepsilon, \nabla u_\varepsilon)) = f$$

by the solution u_{hom} to the homogenized equation

$$-\nabla \cdot (A_{\text{hom}}(\nabla u_{\text{hom}})) = f$$

are stated in Theorem 2 and Theorem 3 in the case of the full space \mathbb{R}^d . The case of a bounded domain – however with a lower rate of convergence – is considered in Theorem 4.

Our second main result – the error estimates for the approximation of the effective material law A_{hom} by the method of representative volumes – is stated in Theorem 9 and Corollary 10.

2.1. Assumptions and setting. We denote by $d \in \mathbb{N}$ the spatial dimension and by $m \in \mathbb{N}$ the system size; in particular, the case $m = 1$ corresponds to a scalar PDE. Let λ and Λ , $0 < \lambda \leq \Lambda < \infty$, denote ellipticity and boundedness constants. Let H and $H \cap B_1$ denote a Hilbert space and the open unit ball in H , respectively. We denote by

$$A : H \times \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$$

a family of operators, indexed by a parameter in H . We require A to satisfy the following conditions:

- (A1) Each operator $A(\omega, \cdot)$ in the family is monotone in the second variable in the sense

$$(A(\omega, \xi_2) - A(\omega, \xi_1)) \cdot (\xi_2 - \xi_1) \geq \lambda |\xi_2 - \xi_1|^2$$

for every parameter $\omega \in H$ and all $\xi_1, \xi_2 \in \mathbb{R}^{m \times d}$.

- (A2) Each operator $A(\omega, \cdot)$ is continuously differentiable in the second variable and Lipschitz in the sense

$$|A(\omega, \xi_2) - A(\omega, \xi_1)| \leq \Lambda |\xi_2 - \xi_1|$$

for every parameter $\omega \in H$ and all $\xi_1, \xi_2 \in \mathbb{R}^{m \times d}$. Furthermore, we have $A(\omega, 0) = 0$ for every parameter $\omega \in H$.

- (A3) The operator $A(\omega, \xi)$ and its derivative $\partial_\xi A(\omega, \xi)$ are differentiable in the parameter ω with bounded derivative in the sense

$$|\partial_\omega A(\omega, \xi)| \leq \Lambda |\xi|, \quad |\partial_\omega \partial_\xi A(\omega, \xi)| \leq \Lambda,$$

for every $\omega \in H$ and all $\xi \in \mathbb{R}^{m \times d}$. Here, ∂_ω and ∂_ξ denote the Fréchet derivative with respect to the first variable and the partial derivative with respect to the second variable, respectively. Furthermore, $|\cdot|$ denotes the operator norm on $H \rightarrow \mathbb{R}^{m \times d}$ and $H \times \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$, respectively.

Throughout our paper, we will reserve the term *parameter field* for a measurable function $\tilde{\omega} : \mathbb{R}^d \rightarrow H \cap B_1$. With help of the operator family A , we may associate to each parameter field $\tilde{\omega}$ a space-dependent monotone material law $\mathbb{R}^d \ni x \mapsto A(\tilde{\omega}(x), \cdot)$. We denote the space of all parameter fields by Ω and equip Ω with the $L^1_{loc}(\mathbb{R}^d; H)$ topology. We then equip the space of parameter fields Ω with a probability measure \mathbb{P} and write $\omega_\varepsilon : \mathbb{R}^d \rightarrow H \cap B_1$ to denote a random parameter field sampled with \mathbb{P} .

It will be our second key assumption that the probability measure \mathbb{P} describes a stationary random field with correlation length ε (which is also the reason why we include the index “ ε ” in our notation). To be precise, we impose the following conditions on \mathbb{P} :

- (P1) \mathbb{P} is *stationary* in the sense that the probability distribution of $\omega_\varepsilon(\cdot + y)$ coincides with the probability distribution of $\omega_\varepsilon(\cdot)$ for all $y \in \mathbb{R}^d$. From a physical viewpoint this corresponds to the assumption of statistical spatial homogeneity of the random material: While each sample of the random material is spatially heterogeneous, the underlying probability distribution is spatially homogeneous.
- (P2) \mathbb{P} features *fast decorrelation on scales $\geq \varepsilon$* in the sense of the spectral gap assumption of Definition 11 below. Here, and throughout the paper $0 < \varepsilon \leq 1$ is fixed and denotes the *correlation length* of the material.

Note that this corresponds to a quantitative assumption of ergodicity by assuming a decorrelation in the coefficient field ω_ε on scales $\geq \varepsilon$.

Under the previous conditions homogenization occurs (in fact (P2) can be weakend to qualitative assumption of ergodicity). In particular, we may introduce the corrector of stochastic homogenization and define a homogenized monotone operator (i. e. a homogenized material law) A_{hom} as follows.

Definition 1 (Corrector and homogenized operator). *Let the assumptions (A1)–(A3) and (P1)–(P2) be in place. Then for all $\xi \in \mathbb{R}^{m \times d}$ there exists a unique random field $\phi_\xi : \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}^m$, called the corrector associated with ξ , with the following properties:*

- (a) *For \mathbb{P} -almost every realization of the random field ω_ε the corrector $\phi_\xi(\omega_\varepsilon, \cdot)$ has the regularity $\phi_\xi(\omega_\varepsilon, \cdot) \in H_{\text{loc}}^1(\mathbb{R}^d; \mathbb{R}^m)$, satisfies $\int_{B_1} \phi_\xi(\omega_\varepsilon, \cdot) dx = 0$, and solves the corrector equation (4) in the sense of distributions.*
- (b) *The gradient of the corrector $\nabla \phi_\xi$ is stationary in the sense that*

$$\nabla \phi_\xi(\omega_\varepsilon, \cdot + y) = \nabla \phi_\xi(\omega_\varepsilon(\cdot + y), \cdot) \quad \text{a.e. in } \mathbb{R}^d$$

holds for \mathbb{P} -a.e. ω_ε and all $y \in \mathbb{R}^d$.

- (c) *The gradient of the corrector $\nabla \phi_\xi$ has finite second moments and vanishing expectation, that is*

$$\mathbb{E}[\nabla \phi_\xi] = 0, \quad \mathbb{E}[|\nabla \phi_\xi|^2] < \infty.$$

- (d) *The corrector \mathbb{P} -almost surely grows sublinearly at infinity in the sense*

$$\lim_{R \rightarrow \infty} \frac{1}{R^2} \int_{B_R} |\phi_\xi(\omega_\varepsilon, x)|^2 dx = 0.$$

Moreover, for each $\xi \in \mathbb{R}^{m \times d}$ we may define

$$(5) \quad A_{\text{hom}}(\xi) := \mathbb{E}[A(\omega_\varepsilon, \xi + \nabla \phi_\xi)],$$

where the right-hand side in this definition is independent of the spatial coordinate x . The map $A_{\text{hom}} : \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ is called the effective operator or the effective material law.

We shall see that the homogenized material law $A_{\text{hom}} : \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ defined by (5) inherits the monotone structure from the heterogeneous material law $A(\omega_\varepsilon, \cdot)$, see Theorem 6 below.

For some of our results we will assume that the following microscopic regularity condition is satisfied. Note that the condition essentially implies a small-scale $C^{1,\alpha}$ regularity theory (i. e. a $C^{1,\alpha}$ theory on the ε scale) for the heterogeneous equation and a global $C^{1,\alpha}$ regularity theory for the homogenized (effective) equation.

- (R) Suppose that at least one of the following three conditions is satisfied:
 - The equation is scalar (i. e. $m = 1$).
 - The number of spatial dimension is at most two (i. e. $d \leq 2$).
 - The system is of Uhlenbeck structure in the sense that there exists a function $\rho : \mathbb{H} \cap B_1 \times \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$ with $A(\omega, \xi) = \rho(\omega, |\xi|^2)\xi$ for all $\omega \in \mathbb{H} \cap B_1$ and all $\xi \in \mathbb{R}^{m \times d}$; furthermore, the effective operator given by (5) is also of Uhlenbeck structure.

Suppose in addition that the second derivative of A with respect to the second variable exists and satisfies the bound $|\partial_\xi \partial_\xi A(\omega, \xi)| \leq \Lambda$ for all $\omega \in \mathbb{H} \cap B_1$ and all $\xi \in \mathbb{R}^{m \times d}$.

Suppose furthermore that the random field ω_ε is Lipschitz regular on small scales in the following sense: There exists a random field \mathcal{C} with uniformly bounded stretched exponential moments $\mathbb{E}[\exp(|\mathcal{C}(x)|^\nu)] \leq 2$ for all $x \in \mathbb{R}^d$ for some $\nu > 0$ such that

$$\sup_{y \in B_\varepsilon(x)} |\nabla \omega_\varepsilon(y)| \leq \mathcal{C}(x) \varepsilon^{-1}$$

holds for all $x \in \mathbb{R}^d$.

2.2. Optimal-order homogenization error estimates. Our first main result is an optimal-order estimate on the homogenization error in the stochastic homogenization of nonlinear uniformly elliptic PDEs (and systems) with monotone nonlinearity. Note that our rate of convergence coincides with the optimal rate of convergence for linear elliptic PDEs and systems [4, 29, 31, 32, 34], which form a subclass of the class of elliptic PDEs with monotone nonlinearity.

Theorem 2 (Optimal-order estimates for the homogenization error for $d \geq 3$). *Let $d \geq 3$. Let the assumptions (A1)–(A3) and (P1)–(P2) be in place. Suppose furthermore that the small-scale regularity condition (R) holds. Let the effective (homogenized) monotone operator $A_{\text{hom}} : \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ be given by the defining formula (5) and let $p > d$. Then for any $f \in L^1(\mathbb{R}^d; \mathbb{R}^m) \cap L^p(\mathbb{R}^d; \mathbb{R}^m)$ the unique weak solution $u_\varepsilon \in \dot{H}^1(\mathbb{R}^d; \mathbb{R}^m)$ to the nonlinear elliptic PDE with heterogeneous coefficients*

$$-\nabla \cdot (A(\omega_\varepsilon, \nabla u_\varepsilon)) = f$$

may be approximated by the solution $u_{\text{hom}} \in \dot{H}^1(\mathbb{R}^d; \mathbb{R}^m)$ to the homogenized effective equation

$$-\nabla \cdot (A_{\text{hom}}(\nabla u_{\text{hom}})) = f$$

up to an error of the order

$$\|u_\varepsilon - u_{\text{hom}}\|_{L^{\frac{2d}{d-2}}(\mathbb{R}^d)} \leq \mathcal{C}(\omega_\varepsilon, f) \varepsilon.$$

Here, $\mathcal{C}(\omega_\varepsilon, f)$ is a random constant whose values may depend on ω_ε and f , but whose (stretched exponential) stochastic moments are uniformly estimated by

$$\mathbb{E} \left[\exp \left(\frac{\mathcal{C}(\omega_\varepsilon, f)^{\bar{\nu}}}{C} \right) \right] \leq 2$$

for some $\bar{\nu} > 0$ depending only on $d, m, \lambda, \Lambda, \rho, p$, and on ν from assumption (R), as well as some $C > 0$ depending only on upper bounds for $\|f\|_{L^1}$ and $\|f\|_{L^p}$ as well as on p .

Since by Theorem 6 the effective material law A_{hom} inherits the monotone structure from the heterogeneous material law $A(\omega, \cdot)$, the solvability of the homogenized equation is guaranteed for any $f \in L^{\frac{2d}{d-2}}(\mathbb{R}^d; \mathbb{R}^m)$ provided that $d \geq 3$.

In the case of low dimension $d \leq 2$ the rate of convergence becomes limited by the central limit theorem scaling. In particular, as for linear elliptic equations, the case $d = 2$ is critical, leading to a logarithmic correction. Furthermore, even for the Poisson equation – which may be regarded as a very particular case of our PDEs

(1) or (2) – the gradient of solutions of the whole-space problem may fail to be square-integrable. For this reason we include a lower order term in our PDEs.

Theorem 3 (Optimal-order estimates for the homogenization error for $d = 2$ and $d = 1$). *Let $d = 2$ or $d = 1$ and let otherwise the assumptions of Theorem 2 be in place. Suppose furthermore $\varepsilon \leq 1$ and $(1 + |x|)^\kappa f \in L^2(\mathbb{R}^d)$ for some $\kappa > 0$. Then the unique weak solution $u_\varepsilon \in H^1(\mathbb{R}^d; \mathbb{R}^m)$ to the nonlinear elliptic PDE with heterogeneous coefficients*

$$(6) \quad -\nabla \cdot (A(\omega_\varepsilon, \nabla u_\varepsilon)) + u_\varepsilon = f$$

may be approximated by the solution $u_{\text{hom}} \in H^1(\mathbb{R}^d; \mathbb{R}^m)$ to the homogenized effective equation

$$(7) \quad -\nabla \cdot (A_{\text{hom}}(\nabla u_{\text{hom}})) + u_{\text{hom}} = f$$

up to an error of the order

$$\|u_\varepsilon - u_{\text{hom}}\|_{L^2(\mathbb{R}^d)} \leq \begin{cases} \mathcal{C}\varepsilon^{1/2} & \text{for } d = 1, \\ \mathcal{C}\varepsilon |\log \varepsilon|^{1/2} & \text{for } d = 2. \end{cases}$$

Here, \mathcal{C} is a random constant which satisfies stretched exponential stochastic moment bounds in the sense $\mathbb{E}[\exp(\mathcal{C}^\nu/C)] \leq 2$ for some $\nu > 0$ as in Theorem 2 and some $C > 0$ depending only on upper bounds for $\|f\|_{L^1}$, $\|(1 + |x|)^\kappa f\|_{L^2}$, and $\|f\|_{L^p}$ as well as on p and κ .

In the absence of the small-scale regularity condition (R), we still obtain half of the optimal rate of convergence. However, we also directly obtain this result in bounded domains. Note that in order to recover the optimal rates of convergence from Theorem 2 and Theorem 3 also for the Dirichlet problem on bounded domains, one would need to construct boundary correctors, as done for linear elliptic PDEs for instance in [10] in the setting of periodic homogenization or in [5, 23] in the setting of stochastic homogenization.

Theorem 4 (Estimates for the homogenization error on bounded domains). *Let $d \geq 2$, let $\mathcal{O} \subset \mathbb{R}^d$ be a bounded domain of class $C^{1,1}$, and let the assumptions (A1)–(A3) and (P1)–(P2) be in place. Define A_{hom} as in formula (5). Let $u_{\text{Dir}} \in H^2(\mathcal{O}; \mathbb{R}^m)$ and $f \in L^2(\mathbb{R}^d; \mathbb{R}^m)$. Then the unique weak solution $u_\varepsilon \in u_{\text{Dir}} + H_0^1(\mathcal{O}; \mathbb{R}^m)$ to the nonlinear elliptic PDE with heterogeneous coefficients*

$$-\nabla \cdot (A(\omega_\varepsilon, \nabla u_\varepsilon)) = f \quad \text{in } \mathcal{O}$$

may be approximated by the solution u_{hom} to the homogenized effective equation

$$-\nabla \cdot (A_{\text{hom}}(\nabla u_{\text{hom}})) = f \quad \text{in } \mathcal{O}$$

(with the same Dirichlet boundary conditions $u_{\text{hom}} \in u_{\text{Dir}} + H_0^1(\mathcal{O}; \mathbb{R}^m)$) up to an error of the order

$$\begin{aligned} & \|u_\varepsilon - u_{\text{hom}}\|_{L^2(\mathcal{O})} \\ & \leq \begin{cases} \mathcal{C}(\omega_\varepsilon, f) (\|f\|_{L^2(\mathcal{O})} + \|u_{\text{Dir}}\|_{H^2(\mathcal{O})}) \varepsilon^{1/2} |\log \varepsilon|^{1/4} & \text{for } d = 2, \\ \mathcal{C}(\omega_\varepsilon, f) (\|f\|_{L^2(\mathcal{O})} + \|u_{\text{Dir}}\|_{H^2(\mathcal{O})}) \varepsilon^{1/2} & \text{for } d \geq 3. \end{cases} \end{aligned}$$

Here, $\mathcal{C}(\omega_\varepsilon, f)$ is a random constant whose values may depend on ω_ε , \mathcal{O} , u_{Dir} , and f , but whose (stretched exponential) stochastic moments are uniformly estimated by

$$\mathbb{E} \left[\exp \left(\frac{\mathcal{C}(\omega_\varepsilon, f)^{\bar{\nu}}}{C(\mathcal{O})} \right) \right] \leq 2$$

for some $\bar{\nu} > 0$ depending only on d, m, λ, Λ , and ρ , and some $C(\mathcal{O}) > 0$ depending only on \mathcal{O} .

Remark 5. An error estimate analogous to Theorem 4 also holds on the full space \mathbb{R}^d , provided that in case $d \geq 3$ we measure the error $u_\varepsilon - u_{\text{hom}}$ in the $L^{2d/(d-2)}$ norm as in Theorem 2 and assume $f \in L^1 \cap L^{\frac{2d}{d-2}}$, and provided that in case $d = 2$ we include a massive term in the equation as in Theorem 3.

We next establish several structural properties of the homogenized operator A_{hom} , including in particular the statement that the homogenized operator A_{hom} inherits the monotone structure from A_ε . Note that these properties are actually true even in the context of qualitative stochastic homogenization, and their proof is fairly elementary. The monotonicity of the effective operator A_{hom} has (essentially) already been established by Dal Maso and Modica [18, 17], at least in the setting of convex integral functionals. We also give sufficient criteria for frame-indifference and isotropy of the homogenized operator A_{hom} .

Theorem 6 (Structure properties of the homogenized equation). *Let the assumptions (A1)–(A3) and (P1)–(P2) be in place. Define the effective (homogenized) material law $A_{\text{hom}} : \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ by (5). Then A_{hom} has the following properties:*

- a) *The map A_{hom} is monotone and Lipschitz continuous in the sense that*

$$(A_{\text{hom}}(\xi_2) - A_{\text{hom}}(\xi_1)) \cdot (\xi_2 - \xi_1) \geq \lambda |\xi_2 - \xi_1|^2$$

and

$$|A_{\text{hom}}(\xi_2) - A_{\text{hom}}(\xi_1)| \leq C(d, m) \frac{\Lambda^2}{\lambda} |\xi_2 - \xi_1|$$

hold for all $\xi_1, \xi_2 \in \mathbb{R}^{m \times d}$. Furthermore, A_{hom} is continuously differentiable and satisfies $A_{\text{hom}}(0) = 0$.

- b) *If the operator A is frame-indifferent in the sense $A(\omega, O\xi) = O^{-1}A(\omega, \xi)$ for all $O \in \text{SO}(m)$, all $\xi \in \mathbb{R}^{m \times d}$, and all $\omega \in \mathbb{H}$, the operator A_{hom} inherits the frame-indifference in the sense $A_{\text{hom}}(O\xi) = O^{-1}A_{\text{hom}}(\xi)$ for all $O \in \text{SO}(m)$ and all $\xi \in \mathbb{R}^{m \times d}$.*
- c) *If the law of the operator $A_\varepsilon(x, \xi) := A(\omega_\varepsilon(x), \xi)$ is isotropic in the sense that the law of the rotated operator $A_\varepsilon^O(x, \xi) := A_\varepsilon(x, \xi O)O$ coincides with the law of A_ε for all $O \in \text{SO}(d)$, the operator A_{hom} inherits the isotropy in the sense $A_{\text{hom}}(\xi O) = A_{\text{hom}}(\xi)O^{-1}$ for all $O \in \text{SO}(d)$ and all $\xi \in \mathbb{R}^{m \times d}$.*

2.3. Optimal-order error estimates for the approximation of the homogenized operator by periodic RVEs. To perform numerical simulations based on the homogenized PDE (2), the effective material law A_{hom} must be determined. The theoretical expression (5) for the effective material law A_{hom} is essentially an average over all possible realizations of the random material; it is therefore a quantity that is not directly computable. To numerically determine the effective (homogenized) material law A_{hom} in practice, the method of representative volumes

is typically employed: A finite sample of the random medium is chosen, say, a cube $[0, L]^d$ of side length $L \gg \varepsilon$, and the cell formula from homogenization theory is evaluated on this sample.

In most cases, a representative volume of mesoscopic size $\varepsilon \ll L \ll 1$ is sufficient to approximate the effective material law A_{hom} well. For this reason, at least in the case of a clear separation of scales $\varepsilon \ll 1$ numerical simulations based on the homogenized equation (2) and the RVE method are several to many orders of magnitude faster than numerical simulations based on the original PDE (1).

In our next theorem, we establish a priori error estimates for the approximation of the effective material law A_{hom} by the method of representative volumes. Our a priori error estimates are of optimal order in the physically relevant cases $d \leq 4$, at least if the issue of boundary layers on the RVE is addressed appropriately. More precisely, we show that the homogenized coefficients A_{hom} may be approximated by the representative volume element method on an RVE of size L up to an error of the order of the natural fluctuations $(\frac{L}{\varepsilon})^{-d/2}$. This rate of convergence coincides with the optimal rate in the case of linear elliptic PDEs and systems, see [32, 33].

There exist various options to define the RVE approximation, see e. g. [22, Section 1.4] for a discussion in the linear elliptic setting. In the present work, we shall consider the case of periodic RVEs: One considers an L -periodic approximation \mathbb{P}_L of the probability distribution \mathbb{P} of the random field ω_ε , that is an L -periodic variant $\omega_{\varepsilon,L}$ of the random field ω_ε (see below for a precise definition), and imposes periodic boundary conditions on the RVE boundary $\partial[0, L]^d$.

Before rigorously defining the notion of periodization, we first note that to any periodic parameter field $\tilde{\omega}_L$ one can unambiguously (and deterministically) associate a homogenized coefficient via the classical periodic homogenization formula.

Definition 7 (Periodic RVE approximation). *Let $A : \mathbb{H} \times \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ be a family of monotone operators satisfying (A1)–(A3). To any L -periodic parameter field $\tilde{\omega}_L$ and any $\xi \in \mathbb{R}^{m \times d}$ we associate the RVE approximation $A^{\text{RVE},L}(\tilde{\omega}_L, \xi)$ for the effective coefficient $A_{\text{hom}}(\xi)$ given by*

$$A^{\text{RVE},L}(\tilde{\omega}_L, \xi) := \int_{[0,L]^d} A(\tilde{\omega}_L, \xi + \nabla \phi_\xi) dx,$$

where the (periodic) corrector $\phi_\xi = \phi_\xi(\tilde{\omega}_L, \cdot)$ is defined as the unique solution in $H_{\text{per}}^1([0, L]^d; \mathbb{R}^m)$ with vanishing mean $\int_{[0,L]^d} \phi_\xi dx = 0$ to the corrector equation

$$-\nabla \cdot (A(\tilde{\omega}_L(x), \xi + \nabla \phi_\xi)) = 0 \quad \text{in } \mathbb{R}^d.$$

We next define our notion of L -periodic approximation of the coefficient field ω_ε . The main condition will be that the statistics of ω_ε and $\omega_{\varepsilon,L}$ must coincide on balls of the form $B_{\frac{L}{4}}(x_0)$ around any $x_0 \in \mathbb{R}^d$.

Definition 8 (L -periodic approximation of \mathbb{P}). *Let \mathbb{P} satisfy the assumptions (P1) and (P2). Let $L \geq \varepsilon$. We say that \mathbb{P}_L is an L -periodic approximation to \mathbb{P} , if \mathbb{P}_L is stationary in the sense of (P1), concentrates on L -periodic parameter fields, satisfies the periodic spectral gap of Definition 11b, and the following property holds: For \mathbb{P} -a.e. random field ω_ε and \mathbb{P}_L -a.e. random field $\omega_{\varepsilon,L}$ we have*

$$\omega_\varepsilon \sim \mathbb{P} \text{ and } \omega_{\varepsilon,L} \sim \mathbb{P}_L \quad \Rightarrow \quad \omega_\varepsilon|_{B_{\frac{L}{4}}} \sim \omega_{\varepsilon,L}|_{B_{\frac{L}{4}}},$$

i.e., if ω_ε is distributed with \mathbb{P} , and $\omega_{\varepsilon,L}$ is distributed with \mathbb{P}_L , then the restricted random fields $\omega_\varepsilon|_{B_{\frac{L}{4}}}$ and $\omega_{\varepsilon,L}|_{B_{\frac{L}{4}}}$ have the same distribution.

For such an L -periodic approximation \mathbb{P}_L , we abbreviate the representative volume approximation as introduced in Definition 7 by $A^{\text{RVE},L}(\xi) := A^{\text{RVE},L}(\omega_{\varepsilon,L}, \xi)$.

Note, however, that the existence of such an L -periodic approximation \mathbb{P}_L of \mathbb{P} has to be proven on a case-by-case basis, depending on the probability distribution. To give one example, for a random field ω_ε of the form $\omega_\varepsilon(x) := \xi(Y_\varepsilon(x))$, where Y_ε is a stationary Gaussian random field arising as the convolution $Y_\varepsilon(x) := (\beta_\varepsilon * W)(x)$ of white noise W with a kernel β_ε with $\text{supp } \beta_\varepsilon \subset B_\varepsilon$, one may construct an L -periodic approximation simply by replacing the white noise W by L -periodic white noise W_L , i. e. by defining $\omega_{\varepsilon,L}(x) := \xi((\beta_\varepsilon * W_L)(x))$.

For the approximation of the effective material law A_{hom} by such an L -periodic representative volume, we establish the following a priori error estimate. Again, our rates of convergence coincide with the optimal rates of convergence for linear elliptic PDEs and systems [29, 31, 32].

Theorem 9 (Error estimate for periodic RVEs). *Let $A : \mathbb{H} \times \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ satisfy the assumptions (A1) – (A3) and let \mathbb{P} satisfy the assumptions (P1)–(P2). Let $L \geq 2\varepsilon$, let \mathbb{P}_L be an L -periodic approximation of \mathbb{P} in the sense of Definition 8, and denote by $A^{\text{RVE},L}(\xi)$ the corresponding representative volume approximation for the homogenized material law $A_{\text{hom}}(\xi)$.*

(a) (Estimate on random fluctuations). *For all $\xi \in \mathbb{R}^{m \times d}$ we have the estimate on random fluctuations*

$$|A^{\text{RVE},L}(\xi) - \mathbb{E}_L [A^{\text{RVE},L}(\xi)]| \leq C |\xi| \left(\frac{L}{\varepsilon} \right)^{-\frac{d}{2}},$$

where C denotes a random variable that satisfies a stretched exponential moment bound uniformly in L , i. e. there exists $C = C(d, m, \lambda, \Lambda, \rho)$ such that

$$\mathbb{E}_L \left[\exp(C^{1/C}) \right] \leq 2.$$

(b) (Estimate for the systematic error). *Suppose that \mathbb{P} and \mathbb{P}_L additionally satisfy the small-scale regularity condition (R). Then for any $\xi \in \mathbb{R}^{m \times d}$ the systematic error of the representative volume method is of higher order in the sense*

$$|\mathbb{E}_L [A^{\text{RVE},L}(\xi)] - A_{\text{hom}}(\xi)| \leq C(1 + |\xi|)^C |\xi| \left(\frac{L}{\varepsilon} \right)^{-(d \wedge 4)} \left| \log \left(\frac{L}{\varepsilon} \right) \right|^{\alpha_d}.$$

Here, α_d is given by

$$\alpha_d := d \wedge 4 + \begin{cases} 2 & \text{for } d \in \{2, 4\}, \\ 0 & \text{for } d = 3, d = 1, \text{ and } d \geq 5 \end{cases}$$

for some $C = C(d, m, \lambda, \Lambda, \nu, \rho)$. In dimension $d \leq 4$ the estimate is optimal (up to the logarithmic factor). Without the small-scale regularity condition (R), the suboptimal estimate

$$|\mathbb{E}_L [A^{\text{RVE},L}(\xi)] - A_{\text{hom}}(\xi)| \leq C |\xi| \left(\frac{L}{\varepsilon} \right)^{-\frac{d \wedge 4}{2}} \left| \log \left(\frac{L}{\varepsilon} \right) \right|^{\alpha_d}$$

holds, where

$$\alpha_d := \frac{d \wedge 4}{2} + \begin{cases} \frac{1}{2} & \text{for } d \in \{2, 4\}, \\ 1 & \text{for } d = 3, d = 1, \text{ and } d \geq 5. \end{cases}$$

In particular, we derive the following overall a priori error estimate for the method of representative volumes.

Corollary 10 (Total L^2 -error for periodic RVEs). *Let assumptions (A1) – (A3) and (P1) – (P2) be in place. Let $L \geq \varepsilon$. Let \mathbb{P}_L be a L -periodic approximation to \mathbb{P} in the sense of Definition 8.*

(a) *If the small-scale regularity condition (R) is satisfied, then for $2 \leq d \leq 7$ we obtain the optimal estimate*

$$\mathbb{E} [|A^{\text{RVE},L}(\xi) - A_{\text{hom}}(\xi)|^2]^{\frac{1}{2}} \leq C(1 + |\xi|)^C |\xi| \left(\frac{L}{\varepsilon}\right)^{-\frac{d}{2}}.$$

Moreover, for $d > 7$ we obtain the suboptimal estimate

$$\mathbb{E} [|A^{\text{RVE},L}(\xi) - A_{\text{hom}}(\xi)|^2]^{\frac{1}{2}} \leq C(1 + |\xi|)^C |\xi| \left(\frac{L}{\varepsilon}\right)^{-4} \left| \log \left(\frac{L}{\varepsilon}\right) \right|^{C(d)}.$$

(b) *If (R) is not satisfied, then for any $d \geq 2$ we obtain the suboptimal estimate*

$$\mathbb{E} [|A^{\text{RVE},L}(\xi) - A_{\text{hom}}(\xi)|^2]^{\frac{1}{2}} \leq C|\xi| \left(\frac{L}{\varepsilon}\right)^{-\frac{d\wedge 4}{2}} \left| \log \left(\frac{L}{\varepsilon}\right) \right|^{C(d)}.$$

For $d \leq 4$ this estimate is optimal except for the logarithmic factor.

In the above estimates, we have $C = C(d, m, \lambda, \Lambda, \rho)$.

2.4. The decorrelation assumption: spectral gap inequality. We finally state and discuss the quantitative assumption on the decorrelation of the random field ω_ε . The majority of results in the present work require the correlations of the random parameter field ω_ε to decay on scales larger than ε in a quantified way. This quantified decay is enforced by assuming that the probability distribution \mathbb{P} satisfies the following spectral gap inequality. As already mentioned previously, this spectral gap inequality holds for example for random fields of the form $\omega_\varepsilon(x) = \beta(\tilde{\omega}(x))$, where $\beta : \mathbb{R} \rightarrow (-1, 1)$ is a 1-Lipschitz function and where $\tilde{\omega}_\varepsilon$ is a stationary Gaussian random field subject to a covariance estimate of the form

$$|\text{Cov} [\tilde{\omega}_\varepsilon(x), \tilde{\omega}_\varepsilon(y)]| \leq C \left(\frac{\varepsilon}{\varepsilon + |x - y|} \right)^{d+\kappa}$$

for all $x, y \in \mathbb{R}^d$ for some $\kappa > 0$ and some $C < \infty$. The derivation is standard and may be found e. g. in [20].

Definition 11 (Spectral gap inequality encoding fast decorrelation on scales $\geq \varepsilon$).

(a) *We say that the probability distribution \mathbb{P} of random fields ω_ε satisfies a spectral gap inequality with correlation length ε and constant $\rho > 0$ if any random variable $F = F(\omega_\varepsilon)$ satisfies the estimate*

$$(8) \quad \mathbb{E} [|F - \mathbb{E}[F]|^2] \leq \frac{\varepsilon^d}{\rho^2} \mathbb{E} \left[\int_{\mathbb{R}^d} \left(\int_{B_\varepsilon(x)} \left| \frac{\partial F}{\partial \omega_\varepsilon} \right| d\tilde{x} \right)^2 dx \right].$$

Here, $\int_{B_\varepsilon(x)} \left| \frac{\partial F}{\partial \omega_\varepsilon}(\omega_\varepsilon) \right| d\tilde{x}$ stands short for

$$\sup_{\delta\omega_\varepsilon} \limsup_{t \rightarrow 0} \frac{|F(\omega_\varepsilon + t\delta\omega_\varepsilon) - F(\omega_\varepsilon)|}{t},$$

where the sup runs over all random fields $\delta\omega_\varepsilon : \mathbb{R}^d \rightarrow H$ supported in $B_\varepsilon(x)$ with $\|\delta\omega_\varepsilon\|_{L^\infty(\mathbb{R}^d)} \leq 1$.

(b) Let $L \geq \varepsilon$ and let \mathbb{P}_L be a probability distribution of L -periodic random fields $\omega_{L,\varepsilon}$. We say that \mathbb{P}_L satisfies a periodic spectral gap inequality with correlation length ε and constant $\rho > 0$ if any random variable $F = F(\omega_{\varepsilon,L})$ satisfies the estimate

$$(9) \quad \mathbb{E}_L[|F - \mathbb{E}_L[F]|^2] \leq \frac{\varepsilon^d}{\rho^2} \mathbb{E} \left[\int_{[0,L]^d} \left(\int_{B_\varepsilon(x)} \left| \frac{\partial F}{\partial \omega_{\varepsilon,L}} \right| d\tilde{x} \right)^2 dx \right].$$

Here, $\int_{B_\varepsilon(x)} \left| \frac{\partial F}{\partial \omega_{\varepsilon,L}}(\omega_{\varepsilon,L}) \right| d\tilde{x}$ stands short for

$$\sup_{\delta \omega_{\varepsilon,L}} \limsup_{t \rightarrow 0} \frac{|F(\omega_{\varepsilon,L} + t\delta \omega_{\varepsilon,L}) - F(\omega_{\varepsilon,L})|}{t},$$

where the sup runs over all L -periodic random fields $\delta \omega_{\varepsilon,L} : \mathbb{R}^d \rightarrow H$ with $\|\delta \omega_{\varepsilon,L}\|_{L^\infty([0,L]^d)} \leq 1$ and support in $B_\varepsilon(x) + LZ^d$.

For a reader not familiar with the concept of spectral gap inequalities, it may be instructive to insert the spatial average $F(\omega_\varepsilon) := \int_{B_r} \omega_\varepsilon dx$ into the definition (8). For this particular choice, the Fréchet derivative is given by $\frac{\partial F}{\partial \omega_\varepsilon} = \frac{1}{|B_r|} \chi_{B_r}$ and the expression on the right-hand side in (8) is of the order $(\frac{\varepsilon}{r})^d$. Thus, the fluctuations of the spatial average $F(\omega_\varepsilon) = \int_{B_r} \omega_\varepsilon(x) dx$ are (at most) of the order $(\frac{\varepsilon}{r})^{d/2}$, just as expected when averaging $(\frac{r}{\varepsilon})^d$ independent random variables of similar variance. However, the importance of spectral gap inequalities in stochastic homogenization [30, 38, 29] stems from the fact that they also entail fluctuation bounds for appropriate *nonlinear* functionals of the random field ω_ε , as the homogenization corrector ϕ_ξ is a nonlinear function of the random field ω_ε (even in the setting of linear elliptic PDEs). To give a simple example for fluctuation bounds for a nonlinear functional of ω_ε , the reader will easily verify that the spectral gap inequality (8) also implies a fluctuation bound of the order $(\frac{\varepsilon}{r})^{d/2}$ for a quantity like $f(\int_{B_r} g(\omega_\varepsilon(x)) dx)$ with two 1-Lipschitz functions $f, g : \mathbb{R} \rightarrow \mathbb{R}$.

3. STRATEGY AND INTERMEDIATE RESULTS

3.1. Key objects: Localized correctors, the two-scale expansion, and linearized correctors. Before turning to the description of our strategy, we introduce the central objects for our approach. Besides the homogenization corrector ϕ_ξ defined in Definition 1, these key objects include the flux corrector σ_ξ , localized versions ϕ_ξ^T and σ_ξ^T of the corrector and the flux corrector, as well as the homogenization correctors $\phi_{\xi,\Xi}^T$ and flux correctors $\sigma_{\xi,\Xi}^T$ for an associated linearized PDE. At the end of the section we also state optimal stochastic moment bounds for these correctors, which will play a central role in the derivation of our main results.

The *flux corrector* σ_ξ is an important quantity in the quantitative homogenization theory of elliptic PDEs, see [29] for a first reference in the context of quantitative stochastic homogenization. Recall that the corrector ϕ_ξ provides the connection between a given macroscopic constant field gradient ξ and the corresponding microscopic field gradient $\xi + \nabla \phi_\xi$. Similarly, the flux corrector $\sigma_\xi : \mathbb{R}^d \rightarrow \mathbb{R}^m \times \mathbb{R}_{\text{skew}}^{d \times d}$ provides a “vector potential” for the difference between the “microscopic” and the “macroscopic” flux in the sense

$$(10) \quad \nabla \cdot \sigma_\xi = A(\omega_\varepsilon, \xi + \nabla \phi_\xi) - A_{\text{hom}}(\xi).$$

The central importance of the flux corrector σ_ξ is that it allows for an elementary representation of the error of the two-scale expansion, see below.

The flux corrector σ_ξ is a $d-1$ -form, i. e. it satisfies the skew-symmetry condition $\sigma_\xi(x) \in \mathbb{R}^m \times \mathbb{R}_{skew}^{d \times d}$ for each $x \in \mathbb{R}^d$. As σ_ξ is a “vector potential” (a $d-1$ -form), the equation (10) only defines σ_ξ up to gauge invariance. It is standard to fix the gauge by requiring σ_ξ to satisfy

$$(11) \quad -\Delta \sigma_{\xi, jk} = \partial_j (A(\omega_\varepsilon, \xi + \nabla \phi_\xi) \cdot e_k) - \partial_k (A(\omega_\varepsilon, \xi + \nabla \phi_\xi) \cdot e_j).$$

Localized correctors. It is cumbersome to work *directly* with the corrector ϕ_ξ and flux corrector σ_ξ in a rigorous manner; in particular, since we need to consider derivatives of (ϕ_ξ, σ_ξ) with respect to compactly supported perturbations of the parameter field ω_ε . For this reason, we shall frequently work with the *localized correctors* ϕ_ξ^T and σ_ξ^T , which for $T > 0$ and *any* parameter field $\tilde{\omega} : \mathbb{R}^d \rightarrow \mathbb{H} \cap B_1$ can be defined unambiguously on a purely deterministic level.

Lemma 12 (Existence of localized correctors). *Let assumptions (A1)–(A3) be in place. Let $\tilde{\omega} : \mathbb{R}^d \rightarrow \mathbb{H} \cap B_1$ be a parameter field, $T \geq 1$, and $\xi \in \mathbb{R}^{m \times d}$. There exist unique tensor fields $\phi_\xi^T := \phi_\xi^T(\tilde{\omega}, \cdot)$ and $\sigma_\xi^T = \{\sigma_{\xi, jk}^T(\tilde{\omega}, \cdot)\}_{jk}$ with*

$$\phi_\xi^T \in H_{\text{uloc}}^1(\mathbb{R}^d; \mathbb{R}^m), \quad \sigma_\xi^T \in H_{\text{uloc}}^1(\mathbb{R}^d; \mathbb{R}^m \times \mathbb{R}_{skew}^{d \times d}),$$

which solve the PDEs

$$(12a) \quad -\nabla \cdot (A(\tilde{\omega}, \xi + \nabla \phi_\xi^T)) + \frac{1}{T} \phi_\xi^T = 0,$$

$$(12b) \quad q_\xi^T := A(\tilde{\omega}, \xi + \nabla \phi_\xi^T),$$

$$(12c) \quad -\Delta \sigma_{\xi, jk}^T + \frac{1}{T} \sigma_{\xi, jk}^T = \partial_j q_{\xi, k}^T - \partial_k q_{\xi, j}^T,$$

in a distributional sense in \mathbb{R}^d . Furthermore, the map $\tilde{\omega} \mapsto (\phi_\xi^T, \sigma_\xi^T)$ is continuous as a map $L^\infty(\mathbb{H}) \rightarrow H_{\text{uloc}}^1$.

Note that the additional term $\frac{1}{T} \phi_\xi^T$ introduces an exponential localization effect. As a consequence, existence and uniqueness of ϕ_ξ^T follow by standard arguments. For example, existence follows by considering the sequence of solutions to the PDEs $-\nabla \cdot (A(\tilde{\omega}, \chi_{B_r(x_0)} \xi + \nabla w_r)) + \frac{1}{T} w_r = 0$ (which admit a unique solution $w_r \in H^1$ by standard monotone operator theory) and passing to the limit $r \rightarrow \infty$; the exponential localization of Lemma 36 below then yields the convergence and the independence of the limit from the choice of x_0 . The argument for σ_ξ^T is similar. For a detailed proof see [21].

Note also that for a stationary random field ω_ε , by uniqueness the associated localized homogenization corrector ϕ_ξ^T and the localized flux corrector σ_ξ^T inherit the property of stationarity.

The localized correctors approximate the original correctors (ϕ_ξ, σ_ξ) in the sense that

$$(13) \quad (\nabla \phi_\xi^T, \nabla \sigma_\xi^T) \rightarrow (\nabla \phi_\xi, \nabla \sigma_\xi) \quad \text{in } L^2(\Omega \times B_r) \text{ for any } r > 0$$

in the limit $T \rightarrow \infty$; for $d \geq 3$, we even have the convergence of ϕ_ξ^T and σ_ξ^T itself to stationary limits ϕ_ξ and σ_ξ (which however may differ from the ϕ_ξ and σ_ξ from Definition 1 by an additive constant). A proof of these facts is provided in Lemma 26.

Two-scale expansion. The correctors (ϕ_ξ, σ_ξ) provide a link between the gradient of the homogenized solution ∇u_{hom} and the gradient of the original solution ∇u_ε via the *two-scale expansion*: Indeed, the two-scale expansion

$$(14) \quad \hat{u}(x) := u_{\text{hom}}(x) + \phi_{\nabla u_{\text{hom}}(x)}(x)$$

is a (formal) approximate solution to the equation with microscopically varying material law in the sense

$$(15) \quad -\nabla \cdot (A(\omega_\varepsilon, \nabla \hat{u})) = f - \nabla \cdot R$$

with the residual R given by

$$(16) \quad R = (\partial_\xi A(\omega_\varepsilon, \xi + \nabla \phi_\xi) \partial_\xi \phi_\xi - \partial_\xi \sigma_\xi)|_{\xi = \nabla u_{\text{hom}}(x)} : \nabla^2 u_{\text{hom}}.$$

If ε is much smaller than the scale on which ∇u_{hom} changes, the residual R will be small (as we will typically have $\partial_\xi \phi_\xi \sim \varepsilon$ and $\partial_\xi \sigma_\xi \sim \varepsilon$, see Proposition 14 for a more precise statement). In this case, taking the difference of (1) and (15) one obtains

$$-\nabla \cdot (A(\omega_\varepsilon, \nabla u_\varepsilon) - A(\omega_\varepsilon, \nabla \hat{u})) = \nabla \cdot R,$$

which by virtue of the monotonicity of the material law A (see (A1)–(A2)) gives rise to an estimate on $\nabla u_\varepsilon - \nabla \hat{u}$. From this estimate, one may then derive a bound on $u_\varepsilon - u_{\text{hom}}$. For a derivation of the error expression in the two-scale expansion in the form (15), which leads to some subtleties regarding measurability, we refer to the forthcoming paper [21]. In our result, we will replace the term $\partial_\xi \phi_\xi|_{\xi = \nabla u_{\text{hom}}}$ in the two-scale expansion by a piecewise constant approximation for u_{hom} (with interpolation); for this variant of the two-scale expansion, essentially finite differences of the form $\phi_{\xi_1} - \phi_{\xi_2}$ appear in the error expression. Details may be found in Proposition 27 respectively in its proof.

Linearized correctors. In the error expression (16) for the residual of the two-scale expansion the derivatives $\partial_\xi \phi_\xi$ and $\partial_\xi \sigma_\xi$ of the corrector and the flux corrector with respect to the field ξ appear. For this reason, we need optimal-order estimates on these derivatives in order to derive an optimal-order error bound for $u_\varepsilon - u_{\text{hom}}$. More precisely, in our variant of the two-scale expansion we need improved estimates for finite differences $\phi_{\xi_1} - \phi_{\xi_2}$ which reflect both the magnitude of the difference $|\xi_1 - \xi_2|$ and the decorrelation in order to derive optimal-order error estimates. Note, however, that this is basically an equivalent problem to estimating the derivatives $\partial_\xi \phi_\xi$. It is interesting to observe that the derivatives $\partial_\xi \phi_\xi$ and $\partial_\xi \sigma_\xi$ are at the same time the homogenization corrector and the flux corrector for the PDE linearized around the field $\xi + \nabla \phi_\xi$: With the coefficient field

$$a_\xi(x) = a_\xi(\omega_\varepsilon, x) := \partial_\xi A(\omega_\varepsilon(x), \xi + \nabla \phi_\xi(\omega_\varepsilon, x)),$$

we deduce by differentiating (4) and (11) that the equalities $\partial_\xi \phi_\xi \Xi = \phi_{\xi, \Xi}$ and $\partial_\xi \sigma_\xi \Xi = \sigma_{\xi, \Xi}$ hold, where $\phi_{\xi, \Xi}$ and $\sigma_{\xi, \Xi}$ are defined as the solution to the PDEs

$$(17a) \quad -\nabla \cdot (a_\xi(x)(\Xi + \nabla \phi_{\xi, \Xi})) = 0,$$

$$(17b) \quad q_{\xi, \Xi} = a_\xi(\Xi + \nabla \phi_{\xi, \Xi}),$$

$$(17c) \quad -\Delta \sigma_{\xi, \Xi, jk} = \partial_j q_{\xi, \Xi, k} - \partial_k q_{\xi, \Xi, j},$$

i. e. $\phi_{\xi, \Xi}$ and $\sigma_{\xi, \Xi}$ are the homogenization correctors for the linear elliptic PDE with random coefficient field a_ξ .

In our analysis we will first estimate the differences $\phi_{\xi_1}^T - \phi_{\xi_2}^T$ of the localized correctors and obtain the estimate on the differences $\phi_{\xi_1} - \phi_{\xi_2}$ by passing to the limit $T \rightarrow \infty$. For this reason, we introduce the linearized localized correctors $\phi_{\xi, \Xi}^T$ and $\sigma_{\xi, \Xi}^T$ as the unique solutions in H_{uloc}^1 to the PDEs

$$(18a) \quad -\nabla \cdot (a_{\xi}^T(\Xi + \nabla \phi_{\xi, \Xi}^T)) + \frac{1}{T} \phi_{\xi, \Xi}^T = 0,$$

$$(18b) \quad q_{\xi, \Xi}^T = a_{\xi}^T(\Xi + \nabla \phi_{\xi, \Xi}^T),$$

$$(18c) \quad -\Delta \sigma_{\xi, \Xi, jk}^T + \frac{1}{T} \sigma_{\xi, \Xi, jk}^T = \partial_j q_{\xi, \Xi, k}^T - \partial_k q_{\xi, \Xi, j}^T,$$

where we have abbreviated

$$a_{\xi}^T(x) = a_{\xi}^T(\omega_{\varepsilon}, x) := \partial_{\xi} A(\omega_{\varepsilon}(x), \xi + \nabla \phi_{\xi}^T(\omega_{\varepsilon}, x)).$$

By differentiating (12a) and (12c), we identify $\partial_{\xi} \phi_{\xi}^T \Xi = \phi_{\xi, \Xi}^T$ and $\partial_{\xi} \sigma_{\xi}^T \Xi = \sigma_{\xi, \Xi}^T$, a fact made rigorous in Lemma 20.

Note that by our assumption of monotonicity and Lipschitz continuity of A (A1)–(A2), the linearized coefficient fields a_{ξ} and a_{ξ}^T are uniformly elliptic and bounded random coefficient fields with a stationary and ergodic distribution. Therefore, the existence (and the notion) of solution to (17a) – (17c) and (18a) – (18c) would follow by the linear theory of stochastic homogenization (see e. g. [29, Lemma 1]), a fact that we however do not need to use: Again, the existence of solutions to the localized corrector problems (18a) and (18c) is evident for any parameter field $\tilde{\omega}$; see Lemma 20 for a proof.

The linearized corrector problem (17a) has also been central for the homogenization result for the linearized equation by Armstrong, Ferguson, and Kuusi [2], and some of our lemmas are analogues of results from [2]: For instance, our differentiability result for the corrector ϕ_{ξ}^T in Lemma 20 is essentially analogous to [2, Lemma 2.4]. Furthermore, in the proofs of [2] small-scale regularity estimates similar to our estimate (42) have been employed. However, in contrast to [2] we establish optimal-order estimates on the linearized correctors $\phi_{\xi, \Xi}^T$ (see Proposition 14 below), a result that will be of key importance for our main theorems.

Corrector estimates. In view of the form of the formula for the residual (16) of the two-scale expansion it is clear that a key ingredient in the quantification of the homogenization error are estimates on the correctors. In this section we state estimates on the localized correctors and its linearizations.

As the typical size of the correctors ϕ_{ξ}^T and $\phi_{\xi, \Xi}^T$ will be at least of order ε due to small-scale fluctuations, the corrector bounds alone cannot capture the decay of fluctuations in $d \geq 3$ optimally. For $d \geq 3$, it is therefore useful to state estimates also for associated vector potentials for the correctors ϕ_{ξ}^T and $\phi_{\xi, \Xi}^T$. In case $d \geq 3$, we introduce a (vector) potential $\theta_{\xi}^T : \mathbb{R}^d \rightarrow \mathbb{R}^{m \times d}$ for the corrector ϕ_{ξ}^T as the (up to an additive constant unique) sublinearly growing solution to the equation

$$(19a) \quad \Delta \theta_{\xi, i}^T = \partial_i \phi_{\xi}^T$$

which entails

$$(19b) \quad \nabla \cdot \theta_{\xi}^T = \phi_{\xi}^T,$$

as well as the corresponding quantity $\theta_{\xi, \Xi}^T$ defined by

$$(20a) \quad \Delta \theta_{\xi, \Xi, i}^T = \partial_i \phi_{\xi, \Xi}^T$$

which yields

$$(20b) \quad \nabla \cdot \theta_{\xi, \Xi}^T = \phi_{\xi, \Xi}^T.$$

Note that as the equations (19a) and (20a) feature a non-decaying right-hand side but lack a massive term, their solvability is not guaranteed for arbitrary parameter fields $\tilde{\omega}$. For random fields ω_ε subject to (P1)–(P2), in $d \geq 3$ the existence of solutions to (19a) and (20a) is a consequence of standard methods in qualitative stochastic homogenization, see e. g. [29].

Proposition 13 (Estimates on the homogenization corrector for the nonlinear monotone PDE). *Let the assumptions (A1)–(A3) and (P1)–(P2) be in place. Then the localized homogenization correctors ϕ_ξ^T and the localized flux correctors σ_ξ^T – defined as the (unique) solution in H_{uloc}^1 to the PDEs (12a) respectively (12c) – are subject to the estimate*

$$(21) \quad \left(\int_{B_r(x_0)} \left| \phi_\xi^T - \int_{B_r(x_0)} \phi_\xi^T(\tilde{x}) d\tilde{x} \right|^2 dx \right)^{1/2} + \left(\int_{B_r(x_0)} \left| \sigma_\xi^T - \int_{B_r(x_0)} \sigma_\xi^T(\tilde{x}) d\tilde{x} \right|^2 dx \right)^{1/2} \leq \begin{cases} C|\xi|\varepsilon^{1/2}r^{1/2} & \text{for } d = 1, \\ C|\xi|\varepsilon|\log \frac{r}{\varepsilon}|^{1/2} & \text{for } d = 2, \\ C|\xi|\varepsilon & \text{for } d \geq 3 \end{cases}$$

for any $r \geq 2\varepsilon$, any $x_0 \in \mathbb{R}^d$, any $T \geq 2\varepsilon^2$, and any $\xi \in \mathbb{R}^{m \times d}$. Furthermore, for $d \geq 3$ we even have

$$(22a) \quad \left(\int_{B_r(x_0)} |\phi_\xi^T|^2 dx \right)^{1/2} + \left(\int_{B_r(x_0)} |\sigma_\xi^T|^2 dx \right)^{1/2} \leq C|\xi|\varepsilon$$

for any $r \geq 2\varepsilon$, any $x_0 \in \mathbb{R}^d$, any $T \geq 2\varepsilon^2$, and any $\xi \in \mathbb{R}^{m \times d}$, while for $d = 1$ and $d = 2$ we have

$$(22b) \quad \left| \int_{B_r(x_0)} \phi_\xi^T dx \right| + \left| \int_{B_r(x_0)} \sigma_\xi^T dx \right| \leq \begin{cases} C|\xi|\varepsilon^{1/2}\sqrt{T}^{1/2} & \text{for } d = 1, \\ C|\xi|\varepsilon|\log \frac{\sqrt{T}}{\varepsilon}|^{1/2} & \text{for } d = 2. \end{cases}$$

Here, C is a random constant whose values may depend on ω_ε , ξ , the localization parameter T , x_0 , and r , but whose (stretched exponential) stochastic moments are uniformly estimated by

$$\mathbb{E}[\exp(C^{\bar{\nu}})] \leq 2$$

for some $\bar{\nu} > 0$ depending only on d , m , λ , Λ , and ρ (in particular, independently of T).

In the case of three and more spatial dimensions $d \geq 3$, the potential field θ_ξ^T exists and satisfies the estimate

$$(23) \quad \left(\int_{B_r(x_0)} \left| \theta_\xi^T - \int_{B_r(x_0)} \theta_\xi^T(\tilde{x}) d\tilde{x} \right|^2 dx \right)^{1/2} \leq \begin{cases} C|\xi|\varepsilon^{3/2}r^{1/2} & \text{for } d = 3, \\ C|\xi|\varepsilon^2|\log \frac{r}{\varepsilon}|^{1/2} & \text{for } d = 4, \\ C|\xi|\varepsilon^2 & \text{for } d \geq 5 \end{cases}$$

for any $r \geq 2\varepsilon$, any $x_0 \in \mathbb{R}^d$, any $T \geq 2\varepsilon^2$, and any $\xi \in \mathbb{R}^{m \times d}$.

Furthermore, the estimates (21), (22a), and (23) also hold for the L^p norm with $2 \leq p \leq \frac{2d}{(d-2)_+}$ and $p < \infty$ in place of the L^2 norm, up to the following modifications: The constant $\bar{\nu}$ may now also depend on p , and the factor $|\log(r/\varepsilon)|^{1/2}$ in (21) and (23) is replaced by $|\log(r/\varepsilon)|$ in the cases $d = 2$ respectively $d = 4$.

Under the additional small-scale regularity assumption (R), we establish the following estimates on the homogenization corrector $\phi_{\xi, \Xi}^T$ associated with the linearized operator $-\nabla \cdot (\partial_\xi A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \nabla \cdot)$. Recall that this homogenization corrector $\phi_{\xi, \Xi}^T$ is equivalently given as the derivative of the homogenization corrector ϕ_ξ^T with respect to ξ in direction Ξ , see Lemma 20.

Proposition 14 (Estimates on the homogenization corrector for the linearized PDE). *Let the assumptions (A1)–(A3) and (P1)–(P2) be in place. Suppose furthermore that the small-scale regularity condition (R) holds. Then the homogenization corrector for the linearized equation $\phi_{\xi, \Xi}^T := \partial_\xi \phi_\xi^T \Xi$ and the corresponding flux corrector $\sigma_{\xi, \Xi}^T := \partial_\xi \sigma_\xi^T \Xi$ given as the solutions to the PDEs (18a) and (18c) are subject to the estimates*

$$(24) \quad \left(\int_{B_r(x_0)} \left| \phi_{\xi, \Xi}^T - \int_{B_\varepsilon(x_0)} \phi_{\xi, \Xi}^T(\tilde{x}) d\tilde{x} \right|^2 dx \right)^{1/2} \\ + \left(\int_{B_r(x_0)} \left| \sigma_{\xi, \Xi}^T - \int_{B_\varepsilon(x_0)} \sigma_{\xi, \Xi}^T(\tilde{x}) d\tilde{x} \right|^2 dx \right)^{1/2} \\ \leq \begin{cases} \mathcal{C}(1 + |\xi|)^C |\Xi| \varepsilon^{1/2} r^{1/2} & \text{for } d = 1, \\ \mathcal{C}(1 + |\xi|)^C |\Xi| \varepsilon \left| \log \frac{r}{\varepsilon} \right|^{1/2} & \text{for } d = 2, \\ \mathcal{C}(1 + |\xi|)^C |\Xi| \varepsilon & \text{for } d \geq 3, \end{cases}$$

for any $r \geq 2\varepsilon$, any $x_0 \in \mathbb{R}^d$, any $T \geq 2\varepsilon^2$, and any $\xi, \Xi \in \mathbb{R}^{m \times d}$. Furthermore, for $d \geq 3$ we even have

$$(25) \quad \left(\int_{B_r(x_0)} |\phi_{\xi, \Xi}^T|^p dx \right)^{1/p} + \left(\int_{B_r(x_0)} |\sigma_{\xi, \Xi}^T|^p dx \right)^{1/p} \leq \mathcal{C}(1 + |\xi|)^C |\xi| \varepsilon$$

for any $r \geq 2\varepsilon$, any $x_0 \in \mathbb{R}^d$, any $p \in [2, \frac{2d}{(d-2)_+}] \cap [2, \infty)$, any $T \geq 2\varepsilon^2$, and any $\xi, \Xi \in \mathbb{R}^{m \times d}$, while for $d = 1$ and $d = 2$ we have

$$\left| \int_{B_\varepsilon(x_0)} \phi_{\xi, \Xi}^T dx \right| + \left| \int_{B_\varepsilon(x_0)} \sigma_{\xi, \Xi}^T dx \right| \leq \begin{cases} \mathcal{C}(1 + |\xi|)^C |\Xi| \varepsilon^{1/2} \sqrt{T}^{1/2} & \text{for } d = 1, \\ \mathcal{C}(1 + |\xi|)^C |\Xi| \varepsilon \left| \log \frac{\sqrt{T}}{\varepsilon} \right|^{1/2} & \text{for } d = 2. \end{cases}$$

Here, \mathcal{C} is a random constant whose values may depend on ω_ε , ξ , Ξ , r , x_0 , and the localization parameter T , but whose (stretched exponential) stochastic moments are uniformly estimated by

$$\mathbb{E}[\exp(\mathcal{C}^{\bar{\nu}})] \leq 2$$

for some $\bar{\nu} > 0$. The constants $\bar{\nu}$ and \mathcal{C} depend only on d , m , λ , Λ , ν , and ρ (in particular, they are independent of T).

In the case of three and more spatial dimensions $d \geq 3$, the potential field $\theta_{\xi, \Xi}^T$ exists and satisfies the estimate

$$(26) \quad \left(\int_{B_r(x_0)} \left| \theta_{\xi, \Xi}^T - \int_{B_\varepsilon(x_0)} \theta_{\xi, \Xi}^T(\tilde{x}) d\tilde{x} \right|^2 dx \right)^{1/2} \\ \leq \begin{cases} \mathcal{C}(1 + |\xi|)^C |\Xi| \varepsilon^{3/2} r^{1/2} & \text{for } d = 3, \\ \mathcal{C}(1 + |\xi|)^C |\Xi| \varepsilon^2 \left| \log \frac{r}{\varepsilon} \right|^{1/2} & \text{for } d = 4, \\ \mathcal{C}(1 + |\xi|)^C |\Xi| \varepsilon^2 & \text{for } d \geq 5 \end{cases}$$

for any $r \geq 2\varepsilon$, any $x_0 \in \mathbb{R}^d$, any $T \geq 2\varepsilon^2$, and any $\xi, \Xi \in \mathbb{R}^{m \times d}$.

Furthermore, all of these estimates also hold for the correctors associated with the adjoint coefficient field $a_\xi^{T,*}$.

Finally, the estimates (24), (25), and (26) also hold for the L^p norm with $2 \leq p \leq \frac{2d}{(d-2)_+}$ and $p < \infty$ in place of the L^2 norm, up to the following modifications: The constant $\bar{\nu}$ may now also depend on p , and the factor $|\log(r/\varepsilon)|^{1/2}$ in (24) and (26) is replaced by $|\log(r/\varepsilon)|$ in the cases $d = 2$ respectively $d = 4$.

A key consequence of our estimates on the linearized correctors $\phi_{\xi,\Xi}^T$ and $\sigma_{\xi,\Xi}^T$ is the following estimate on differences of the correctors ϕ_ξ and σ_ξ for different values of ξ .

Corollary 15. *Let the assumptions (A1)–(A3) and (P1)–(P2) be in place. Then for $T \rightarrow \infty$ the correctors ϕ_ξ^T and σ_ξ^T (for $d \geq 3$) respectively the renormalized correctors $\phi_\xi^T - \int_{B_\varepsilon(0)} \phi_\xi^T d\tilde{x}$ and $\sigma_\xi^T - \int_{B_\varepsilon(0)} \sigma_\xi^T d\tilde{x}$ (for $d = 1, 2$) converge to limits ϕ_ξ and σ_ξ . The limits ϕ_ξ and σ_ξ satisfy the corrector equations (4) and (10); furthermore, in case $d \geq 3$ they are subject to the estimate*

$$(27) \quad \left(\int_{B_r(x_0)} |\phi_\xi|^p + |\sigma_\xi|^p dx \right)^{1/p} \leq C|\xi|\varepsilon$$

for any $r \geq 2\varepsilon$, any $x_0 \in \mathbb{R}^d$, any $p \in [2, \frac{2d}{d-2}]$, and any $\xi \in \mathbb{R}^{m \times d}$, while in case $d = 1, 2$ they satisfy

$$(28) \quad \left(\int_{B_r(x_0)} |\phi_\xi|^2 + |\sigma_\xi|^2 dx \right)^{1/2} \leq \begin{cases} C|\xi|\varepsilon^{1/2}(r + |x_0|)^{1/2} & \text{for } d = 1, \\ C|\xi|\varepsilon \left| \log \frac{r + |x_0|}{\varepsilon} \right|^{1/2} & \text{for } d = 2 \end{cases}$$

for any $r \geq 2\varepsilon$, any $x_0 \in \mathbb{R}^d$, and any $\xi \in \mathbb{R}^{m \times d}$. Here, C is a random constant with stretched exponential moments in the sense

$$\mathbb{E}[\exp(C^\bar{\nu})] \leq 2$$

for some $\bar{\nu} > 0$ depending only on d, m, λ, Λ , and ρ .

Suppose furthermore that the small-scale regularity condition (R) holds. Then the difference of homogenization correctors $\phi_{\xi_1} - \phi_{\xi_2}$ and $\sigma_{\xi_1} - \sigma_{\xi_2}$ is estimated by

(29)

$$\left(\int_{B_r(x_0)} |\phi_{\xi_1} - \phi_{\xi_2}|^2 + |\sigma_{\xi_1} - \sigma_{\xi_2}|^2 dx \right)^{1/2} \leq \begin{cases} C(1 + |\xi_1|^C + |\xi_2|^C) |\xi_1 - \xi_2| \varepsilon^{1/2} (\varepsilon + |x_0|)^{1/2} & \text{for } d = 1, \\ C(1 + |\xi_1|^C + |\xi_2|^C) |\xi_1 - \xi_2| \varepsilon \left| \log \frac{|x_0| + 2\varepsilon}{\varepsilon} \right|^{1/2} & \text{for } d = 2, \\ C(1 + |\xi_1|^C + |\xi_2|^C) |\xi_1 - \xi_2| \varepsilon & \text{for } d \geq 3 \end{cases}$$

for any $\xi_1, \xi_2 \in \mathbb{R}^{m \times d}$, any $r \geq \varepsilon$, and any $x_0 \in \mathbb{R}^d$, where the random constant C satisfies

$$\mathbb{E}[\exp(C^\bar{\nu})] \leq 2$$

for some $\bar{\nu} > 0$. Here, $\bar{\nu}$ and C depend only on $d, m, \lambda, \Lambda, \rho$, and ν .

3.2. The strategy of proof for corrector estimates. A key difficulty in quantitative stochastic homogenization is the derivation of appropriate estimates on the (localized) homogenization correctors ϕ_ξ^T and the (localized) flux corrector σ_ξ^T : As previously mentioned, corrector bounds form the basis for homogenization error estimates and the analysis of the representative volume approximation. For our purposes, we for example need to show that the homogenization corrector ϕ_ξ^T and the flux corrector σ_ξ^T are at most of order $|\xi|\varepsilon$ (at least in case $d \geq 3$).

In periodic homogenization, that is for ε -periodic operators $-\nabla \cdot A_\varepsilon(x, \nabla \cdot)$, such corrector bounds are an easy consequence of the periodicity: By the defining equation (12a), the corrector ϕ_ξ^T is ε -periodic, has vanishing average on each periodicity cell $[0, \varepsilon]^d$, and is subject to an energy estimate of the form $\int_{[0, \varepsilon]^d} |\nabla \phi_\xi^T|^2 dx \leq C|\xi|^2$. By the Poincaré inequality on the periodicity cell, this implies a bound of order $|\xi|\varepsilon$ on the homogenization corrector ϕ_ξ^T . The derivation of the corresponding estimate for the flux corrector σ_ξ^T is analogous.

In contrast, in quantitative stochastic homogenization the derivation of such estimates on the correctors is much more involved and presents one of the key challenges. Our strategy for the derivation of bounds on ϕ_ξ^T , which is strongly inspired by [32, 33, 29, 31] but streamlined by minimizing the use of elliptic regularity, proceeds as follows:

- As outlined in Proposition 13, it is our goal to prove a corrector estimate of essentially the form

$$\inf_{b \in \mathbb{R}^m} \left(\int_{B_R(x_0)} |\phi_\xi^T - b|^p dx \right)^{1/p} \leq \begin{cases} \mathcal{C}\varepsilon & \text{for } d \geq 3 \\ \mathcal{C}\varepsilon |\log(R/\varepsilon)| & \text{for } d = 2, \\ \mathcal{C}\varepsilon^{1/2} R^{1/2} & \text{for } d = 1 \end{cases}$$

for any $R \geq \varepsilon$, where \mathcal{C} is a random constant with stretched exponential moments in the sense $\mathbb{E}[\exp(\mathcal{C}^{1/C})] \leq 2$.

- In principle, the technical Lemma 25 (see below) reduces the derivation of such estimates on the corrector ϕ_ξ^T to (mostly) the derivation of bounds on stochastic moments of integral functionals of the form

$$(30) \quad F(\omega_\varepsilon) := \int_{\mathbb{R}^d} g \cdot \nabla \phi_\xi^T dx,$$

where g basically takes a weighted average of $\nabla \phi_\xi^T$ on a certain scale r with $\varepsilon \leq r \leq R$.

- The random variables $F(\omega_\varepsilon)$ as defined in (30) have vanishing expectation $\mathbb{E}[F(\omega_\varepsilon)] = 0$ due to the vanishing expectation of the corrector gradient $\mathbb{E}[\nabla \phi_\xi^T] = 0$. The latter property is a consequence of stationarity of the corrector ϕ_ξ^T : Since the probability distribution of the random monotone operator $A(\omega_\varepsilon(x), \cdot)$ is invariant under spatial translations, the expectation of the corrector $\mathbb{E}[\phi_\xi^T(x)]$ is the same for all points $x \in \mathbb{R}^d$. This yields $\mathbb{E}[\nabla \phi_\xi^T(x)] = \nabla \mathbb{E}[\phi_\xi^T(x)] = 0$. As a consequence of the vanishing expectation, it suffices to bound the centered moments of $F(\omega_\varepsilon)$, that is, the stochastic moments of $F(\omega_\varepsilon) - \mathbb{E}[F(\omega_\varepsilon)]$.
- Concentration inequalities – like the spectral gap inequality – are one of the most widespread probabilistic tools for establishing bounds on centered

moments of random variables. In our context, that is for random fields ω_ε with correlations restricted to the length scale ε , they will read for example

$$(31) \quad \mathbb{E} \left[\left| F(\omega_\varepsilon) - \mathbb{E}[F(\omega_\varepsilon)] \right|^2 \right] \leq \frac{\varepsilon^d}{\rho^2} \mathbb{E} \left[\int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} \left| \frac{\partial F}{\partial \omega_\varepsilon} \right| d\tilde{x} \right|^2 dx \right]$$

for all random variables $F = F(\omega_\varepsilon)$. Here, $\frac{\partial F}{\partial \omega_\varepsilon}$ denotes (essentially) the Frechét derivative of F with respect to the field ω_ε and $\rho > 0$ denotes a constant.

Note that in stochastic homogenization it is an *assumption* that a spectral gap inequality like (31) holds for the random field ω_ε . This assumption on the probability distribution of the random field ω_ε *encodes the decorrelation properties* of ω_ε . For instance, the reader may convince oneself that (31) implies an estimate of the order $(\varepsilon/r)^{d/2}$ on the fluctuations of the average $F(\omega_\varepsilon) := \int_{B_r} \omega_\varepsilon dx$ on a scale $r \geq \varepsilon$, as one would expect for e.g. the average of $(r/\varepsilon)^d$ i. i. d. random variables; see also the discussion below Definition 11.

A spectral gap inequality of the form (31) is valid for many classes of random fields, see Figure 1 and the accompanying text. It also implies estimates on higher centered moments, see Lemma 16 below.

- In order to employ the spectral gap inequality (31) to estimate the stochastic moments of random variables $F(\omega_\varepsilon) = \int g \cdot \nabla \phi_\xi^T dx$ as defined in (31), we need to estimate the right-hand side of (31), that is we need to estimate the sensitivity of the functionals $F(\omega_\varepsilon)$ with respect to changes in the coefficient field ω_ε . By standard computations (see the proof of Lemma 17a for details), one may show that the Frechét derivative of F with respect to ω_ε is given by

$$(32) \quad \frac{\partial F}{\partial \omega_\varepsilon} = \partial_\omega A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \otimes \nabla h$$

where h is the solution to the auxiliary linear elliptic PDE

$$(33) \quad -\nabla \cdot (a_\xi^{T,*}(x) \nabla h) + \frac{1}{T} h = \nabla \cdot g$$

(the uniformly elliptic and bounded coefficient field a_ξ^T being given by $a_\xi^T(x) := \partial_\xi A(\omega(x), \xi + \nabla \phi_\xi^T)$).

By the (standard) growth assumptions for $A(\omega_\varepsilon, \cdot)$ in (A3), the representation (32) implies that the sensitivity $\frac{\partial F}{\partial \omega_\varepsilon}$ may be estimated by $C|\xi + \nabla \phi_\xi^T| |\nabla h|$. In conclusion, by the spectral gap inequality (31) respectively its version for higher stochastic moments in Lemma 16 we see that we have

$$(34) \quad \mathbb{E}[|F|^q]^{1/q} \lesssim q\varepsilon^{d/2} \mathbb{E} \left[\left| \int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T| |\nabla h| d\tilde{x} \right|^2 dx \right|^{q/2} \right]^{1/q}.$$

- Let us next discuss how to estimate the right-hand side of (34). Recall that g basically takes a weighted average on a scale r (see (30)), i.e. g is supported on a ball B_r and satisfies an estimate like $|g| \lesssim r^{-d}$. As a consequence, we obtain the deterministic estimate $\int_{\mathbb{R}^d} |\nabla h|^2 dx \lesssim r^{-d}$ by a simple energy estimate for the defining equation (33). If we knew that the small-scale averages of the corrector gradient were bounded in the sense

$\int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T|^2 d\tilde{x} \lesssim |\xi|^2$, by Hölder's inequality we would directly obtain the deterministic bound

$$\int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T| |\nabla h| d\tilde{x} \right|^2 dx \lesssim |\xi|^2 \int_{\mathbb{R}^d} |\nabla h|^2 dx \lesssim |\xi|^2 r^{-d}.$$

By (34) this would imply an estimate of fluctuation order on the functionals (30) of the form $\mathbb{E}[|F|^q]^{1/q} \lesssim q|\xi|(\varepsilon/r)^{d/2}$. This would be precisely the bound we seek to obtain.

However, in the context of stochastic homogenization we cannot expect a uniform bound on the locally averaged corrector gradient $\int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T|^2 d\tilde{x}$, as in general the random field ω_ε may contain geometric configurations which could cause the corrector gradient to be arbitrarily large. As we are dealing with the case of systems, we also cannot expect more than L^p integrability of the gradient ∇h of the auxiliary function for a Meyers exponent p slightly larger than 2. We therefore need (almost) an L^∞ bound on the local averages $\int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T|^2 d\tilde{x}$. It is here useful to introduce an auxiliary quantity, namely the minimal radius $r_{*,T,\xi}(x)$ above which the corrector ϕ_ξ^T satisfies a bound of the form

$$\int_{B_R(x)} |\phi_\xi^T|^2 d\tilde{x} \leq 1 \quad \text{for all } R \geq r_{*,T,\xi}(x)$$

(plus one additional technical condition). This allows us to bound using a trivial estimate and the Caccioppoli inequality

$$\int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T|^2 d\tilde{x} \leq \left(\frac{r_{*,T,\xi}(x)}{\varepsilon} \right)^d \underbrace{\int_{B_{r_{*,T,\xi}(x)}} |\xi + \nabla \phi_\xi^T|^2 d\tilde{x}}_{\leq C}.$$

This estimate is however again not sufficient for our purposes. It is here that *elliptic regularity theory* in form of the hole-filling estimate enters and provides a slight but crucial improvement

$$\int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T|^2 d\tilde{x} \leq C \left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{d-\delta}$$

for some $\delta > 0$ (see Lemma 88 for details).

Together with a version of the spectral gap estimate – see Lemma 16 below – and the vanishing expectation $\mathbb{E}[\nabla \phi_\xi^T] = 0$, this estimation strategy yields a bound of the form

$$(35) \quad \mathbb{E}[|F|^{2q}]^{1/2q} \leq C|\xi|q \left(\frac{\varepsilon}{r} \right)^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q}$$

for any $0 < \tau < 1$ and any q large enough (the latter of which is not a problem).

- One then observes that one may control the moments of $r_{*,T,\xi}$ on the right-hand side of (35) in terms of moments of functionals $F = \int g \cdot \nabla \phi_\xi^T dx$ (see the proof of Proposition 13). It is here that the slight gain δ in the

exponent (from hole-filling) is crucial, as it causes the estimate to buckle, yielding a bound of the form

$$\mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^q \right]^{1/q} \lesssim q^C.$$

The resulting moment bounds on $r_{*,T,\xi}$ then allow to deduce the corrector estimate in Proposition 13.

- The derivation of the estimates for the flux corrector σ_ξ^T and the linearized correctors $\phi_{\xi,\Xi}^T, \sigma_{\xi,\Xi}^T$ follows a similar strategy. However, in the case of the linearized correctors $\phi_{\xi,\Xi}^T, \sigma_{\xi,\Xi}^T$ the derivation of the sensitivity estimates involves an additional integrability issue on small scales, making it necessary to use a $C^{1,\alpha}$ regularity theory on the microscopic scale. It is here that our additional regularity assumption (R) enters. For details, see the derivation of Lemma 21.

Let us now state the lemmas used in the course of the proof of our main results. An immediate consequence of the spectral gap inequality of Definition 11 is the following version for the q -th centered moment; for a proof, we refer to e.g. [19, Proposition 3.1].

Lemma 16. *Suppose \mathbb{P} satisfies (P1) and (P2). Then for any $q \geq 1$ we have*

$$\mathbb{E} \left[\left| F(\omega_\varepsilon) - \mathbb{E}[F(\omega_\varepsilon)] \right|^{2q} \right]^{1/q} \leq Cq^2 \varepsilon^d \mathbb{E} \left[\left| \int_{\mathbb{R}^d} \left(\int_{B_\varepsilon(x)} \left| \frac{\partial F}{\partial \omega_\varepsilon} \right|^2 dx \right)^{q/2} \right|^{1/q} \right].$$

A central step towards the proof of the corrector estimates of Proposition 13 are the following estimates on stochastic moments of linear functionals of the localized corrector and the localized flux corrector.

Lemma 17 (Estimates for linear functionals of the corrector and the flux corrector for the monotone system). *Let the assumptions (A1)–(A3) and (P1)–(P2) be satisfied. Let $\xi \in \mathbb{R}^{m \times d}$, $T \geq 2\varepsilon^2$, $K_{mass} \geq C(d, m, \lambda, \Lambda)$, and let ϕ_ξ^T be the unique nongrowing solution to the corrector equation (12a). Define for any $x_0 \in \mathbb{R}^d$*

(36)

$r_{*,T,\xi}(x_0) := \inf \left\{ r = 2^k \varepsilon : k \in \mathbb{N}_0 \text{ and for all } R = 2^\ell \varepsilon \geq r, \ell \in \mathbb{N}_0, \text{ we have both} \right.$

$$\left. \begin{aligned} & \frac{1}{R^2} \int_{B_R(x_0)} \left| \phi_\xi^T - \int_{B_R(x_0)} \phi_\xi^T \right|^2 dx \leq |\xi|^2 \\ & \text{and } \frac{1}{\sqrt{T}} \left| \int_{B_R(x_0)} \phi_\xi^T dx \right| \leq K_{mass} |\xi| \end{aligned} \right\}.$$

Then the random field $r_{*,T,\xi}$ is stationary and allows for the following estimates.

- a) Let $2 < p < 2 + c$ (where $c = c(d, m, \lambda, \Lambda) > 0$ will be defined in the proof below), let $x_0 \in \mathbb{R}^d$ and $r \geq \varepsilon$, and let F be a functional of the form

$$F(\omega) := \int_{\mathbb{R}^d} g \cdot \nabla \phi_\xi^T dx$$

for some $g \in L^p(\mathbb{R}^d; \mathbb{R}^{m \times d})$ with $\text{supp } g \subset B_r(x_0)$ and

$$\left(\int_{B_r(x_0)} |g|^p dx \right)^{1/p} \leq r^{-d}.$$

Then there exists an exponent $\delta = \delta(d, m, \lambda, \Lambda) > 0$ (from hole-filling) such that the stochastic moments of the functional F_σ may be estimated by

$$\mathbb{E}[|F|^{2q}]^{1/2q} \leq C|\xi|q \left(\frac{\varepsilon}{r}\right)^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q}$$

for any $0 < \tau < 1$ and any $q \geq C$. Here, the constants C may depend on $d, m, \lambda, \Lambda, \rho, K_{\text{mass}}, p$, and τ .

- b) Let $2 < p < 2 + c$, let $x_0 \in \mathbb{R}^d$ and $r \geq \varepsilon$, and let F_σ be a functional of the form

$$F_\sigma(\omega) := \int_{\mathbb{R}^d} g \cdot \nabla \sigma_{\xi,jk}^T dx$$

for some $g \in L^p(\mathbb{R}^d; \mathbb{R}^{m \times d})$ with $\text{supp } g \subset B_r(x_0)$ and

$$\left(\int_{B_r(x_0)} |g|^p dx \right)^{1/p} \leq r^{-d}.$$

Then there exists $\delta = \delta(d, m, \lambda, \Lambda) > 0$ such that the stochastic moments of the functional F_σ may be estimated by

$$\mathbb{E}[|F_\sigma|^{2q}]^{1/2q} \leq C|\xi|q \left(\frac{\varepsilon}{r}\right)^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q}$$

for any $0 < \tau < 1$ and any $q \geq C(d, m, \lambda, \Lambda, p, \tau)$.

- c) Suppose that $d \geq 3$. Let $2 < p < 2 + c$, let $x_0 \in \mathbb{R}^d$ and $r \geq \varepsilon$, and let F_θ be a functional of the form

$$F_\theta(\omega) := \int_{\mathbb{R}^d} g \cdot \nabla \theta_{\xi,i}^T dx$$

for some $g \in L^p(\mathbb{R}^d; \mathbb{R}^{m \times d})$ with $\text{supp } g \subset B_r(x_0)$ and

$$\left(\int_{B_r(x_0)} |g|^p dx \right)^{1/p} \leq r^{-d}.$$

Then there exists $\delta > 0$ such that the stochastic moments of the functional F_θ may be estimated by

$$\mathbb{E}[|F_\theta|^{2q}]^{1/2q} \leq Cr|\xi|q \left(\frac{\varepsilon}{r}\right)^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q}$$

for any $0 < \tau < 1$ and any $q \geq C(d, m, \lambda, \Lambda, p, \tau)$.

In particular, all of our estimates are independent of $T \geq 2\varepsilon^2$.

We also obtain the following estimates on averages of the correctors.

Lemma 18. *Given the assumptions of Lemma 17, for any $0 < \tau < 1$, any $q \geq C$, any $x_0 \in \mathbb{R}^d$, any $r \geq \varepsilon$, and any $R \geq r$ we have the estimate*

$$(37) \quad \begin{aligned} & \mathbb{E} \left[\left| \int_{B_r(x_0)} \phi_\xi^T dx - \int_{B_R(x_0)} \phi_\xi^T dx \right|^{2q} \right]^{1/2q} \\ & + \mathbb{E} \left[\left| \int_{B_r(x_0)} \sigma_\xi^T dx - \int_{B_R(x_0)} \sigma_\xi^T dx \right|^{2q} \right]^{1/2q} \\ & \leq Cq|\xi| \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q} \left(\sum_{l=0}^{\log_2 \frac{R}{r}} (2^l r)^2 \left(\frac{\varepsilon}{2^l r} \right)^d \right)^{1/2}. \end{aligned}$$

Furthermore, for any $R \geq \sqrt{T}$ we have the bound

$$(38) \quad \begin{aligned} & \mathbb{E} \left[\left| \int_{B_R(x_0)} \phi_\xi^T dx \right|^q \right]^{1/q} + \mathbb{E} \left[\left| \int_{B_R(x_0)} \sigma_\xi^T dx \right|^q \right]^{1/q} \\ & \leq Cq|\xi| \sqrt{T} \left(\frac{\varepsilon}{R} \right)^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q}. \end{aligned}$$

We then establish the following moment bounds on the minimal radius $r_{*,T,\xi}$, which will enable us to deduce the corrector bounds in Proposition 13.

Lemma 19 (Moment bound on the minimal radius). *Given the assumptions of Lemma 17, there exists $K_{mass} = K_{mass}(d, m, \lambda, \Lambda)$ such that the minimal radius $r_{*,T,\xi}$ has stretched exponential moments in the sense*

$$\mathbb{E} \left[\exp \left(\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{1/C} \right) \right] \leq 2$$

for some constant $C = C(d, m, \lambda, \Lambda, \rho)$.

It will be central for our optimal-order homogenization error estimates to obtain optimal-order estimates on *differences of correctors* for different macroscopic field gradients ξ , like $\phi_{\xi_1} - \phi_{\xi_2}$ for $\xi_1, \xi_2 \in \mathbb{R}^{m \times d}$. To estimate such differences, we will rely on estimates for the derivative $\phi_{\xi, \Xi}^T$ of the corrector ϕ_ξ^T with respect to ξ in direction Ξ . Formally differentiating the corrector equation (12a) with respect to ξ , we obtain for $\phi_{\xi, \Xi}^T := \partial_\xi \phi_\xi^T \Xi$ the PDE

$$-\nabla \cdot (\partial_\xi A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T)(\Xi + \nabla \phi_{\xi, \Xi}^T)) + \frac{1}{T} \phi_{\xi, \Xi}^T = 0$$

(see also (18a) above). It is interesting that this PDE again takes the form of a corrector equation, namely the corrector equation for the linear elliptic equation with random coefficient field $\partial_\xi A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T)$.

We next show that this differentiation can be justified rigorously.

Lemma 20 (Differentiability of the corrector with respect to ξ). *Let $T > 0$ and assume (A1)–(A2). For any $\xi \in \mathbb{R}^{m \times d}$ and any parameter field $\tilde{\omega}$, let ϕ_ξ^T denote the unique solution in $H_{\text{uloc}}^1(\mathbb{R}^d; \mathbb{R}^m)$ to the localized corrector equation (12a). Denote by $\phi_{\xi, \Xi}^T$ the unique solution in $H_{\text{uloc}}^1(\mathbb{R}^d; \mathbb{R}^m)$ to the localized linearized corrector equation (18a). Then the map $\xi \mapsto \phi_\xi^T$ – as a map $\mathbb{R}^{m \times d} \rightarrow H_{\text{uloc}}^1(\mathbb{R}^d; \mathbb{R}^m)$ – is differentiable with respect to ξ in the Fréchet sense with*

$$\partial_\xi \phi_\xi^T \Xi = \phi_{\xi, \Xi}^T$$

for any $\Xi \in \mathbb{R}^{m \times d}$.

Similarly, denoting by σ_ξ^T and $\sigma_{\xi, \Xi}^T$ the unique solutions in H_{uloc}^1 to the PDEs (12c) and (18c), the map $\xi \mapsto \sigma_\xi^T$ – as a map $\mathbb{R}^{m \times d} \rightarrow H_{\text{uloc}}^1(\mathbb{R}^d; \mathbb{R}^m \times \mathbb{R}_{\text{skew}}^{d \times d})$ – is differentiable with respect to ξ in the Fréchet sense with

$$\partial_\xi \sigma_\xi^T \Xi = \sigma_{\xi, \Xi}^T$$

for any $\Xi \in \mathbb{R}^{m \times d}$.

In order to establish appropriate estimates on the linearized corrector $\phi_{\xi, \Xi}^T$, we again start by deriving estimates for linear functionals of the corrector gradient.

Lemma 21 (Estimates for linear functionals of the corrector and the flux corrector for the linearized equation). *Let the assumptions (A1)–(A3) and (P1)–(P2) be satisfied. Suppose in addition that the regularity condition (R) holds. Let $\xi, \Xi \in \mathbb{R}^{m \times d}$, $K_{\text{mass}} \geq C(d, m, \lambda, \Lambda)$, $T \geq 2\varepsilon^2$, and let $\phi_{\xi, \Xi}^T$ be the unique solution in H_{uloc}^1 to the corrector equation for the linearized problem (18a). Define for any $x_0 \in \mathbb{R}^d$*

(39)

$r_{*, T, \xi, \Xi}(x_0) := \inf \left\{ r = 2^k \varepsilon : k \in \mathbb{N}_0 \text{ and for all } R = 2^\ell \varepsilon \geq r, \ell \in \mathbb{N}_0, \text{ we have both} \right.$

$$\begin{aligned} & \frac{1}{R^2} \int_{B_R(x_0)} \left| \phi_{\xi, \Xi}^T - \int_{B_R(x_0)} \phi_{\xi, \Xi}^T \right|^2 dx \leq |\Xi|^2 \\ & \text{and } \frac{1}{\sqrt{T}} \left| \int_{B_R(x_0)} \phi_{\xi, \Xi}^T dx \right| \leq K_{\text{mass}} |\Xi| \left. \right\}. \end{aligned}$$

Then the random field $r_{*, T, \xi, \Xi}$ is stationary and allows for the following estimates.

- a) Let $2 < p < 2 + c$ (where $c = c(d, m, \lambda, \Lambda) > 0$ will be defined in the proof below), let $x_0 \in \mathbb{R}^d$ and $r \geq \varepsilon$, and let F be a functional of the form

$$F(\omega) := \int_{\mathbb{R}^d} g \cdot \nabla \phi_{\xi, \Xi}^T dx$$

for some $g \in L^p(\mathbb{R}^d; \mathbb{R}^{m \times d})$ with $\text{supp } g \subset B_r(x_0)$ and

$$\left(\int_{B_r(x_0)} |g|^p dx \right)^{1/p} \leq r^{-d}.$$

Then there exists an exponent $\delta = \delta(d, m, \lambda, \Lambda) > 0$ (from hole-filling) such that the stochastic moments of the functional F may be estimated by

$$\mathbb{E}[|F|^{2q}]^{1/2q} \leq C(1 + |\xi|^C) |\Xi| q^C \left(\frac{\varepsilon}{r} \right)^{d/2} \mathbb{E} \left[\left(\frac{r_{*, T, \xi, \Xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q}$$

for any $0 < \tau \leq c = c(d, m, \lambda, \Lambda)$ and any $q \geq C$. Here, the constant C may depend on $d, m, \lambda, \Lambda, \rho, \nu, K_{\text{mass}}, p$, and τ .

- b) Let $2 < p < 2 + c$, let $x_0 \in \mathbb{R}^d$ and $r \geq \varepsilon$, and let F_σ be a functional of the form

$$F_\sigma(\omega) := \int_{\mathbb{R}^d} g \cdot \nabla \sigma_{\xi, \Xi, jk}^T dx$$

for some $g \in L^p(\mathbb{R}^d; \mathbb{R}^{m \times d})$ with $\text{supp } g \subset B_r(x_0)$ and

$$\left(\int_{B_r(x_0)} |g|^p dx \right)^{1/p} \leq r^{-d}.$$

Then the stochastic moments of the functional F_σ may be estimated by

$$\mathbb{E}[|F_\sigma|^{2q}]^{1/2q} \leq C(1 + |\xi|^C) |\Xi| q^C \left(\frac{\varepsilon}{r} \right)^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi,\Xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q}$$

for any $0 < \tau \leq c(d, m, \lambda, \Lambda)$ and any $q \geq C$.

- c) Suppose that $d \geq 3$. Let $2 < p < 2 + c$, let $x_0 \in \mathbb{R}^d$ and $r \geq \varepsilon$, and let F_θ be a functional of the form

$$F_\theta(\omega) := \int_{\mathbb{R}^d} g \cdot \nabla \theta_{\xi, \Xi, i}^T dx$$

for some $g \in L^p(\mathbb{R}^d; \mathbb{R}^{m \times d})$ with $\text{supp } g \subset B_r(x_0)$ and

$$\left(\int_{B_r(x_0)} |g|^p dx \right)^{1/p} \leq r^{-d}.$$

Then the stochastic moments of the functional F_θ may be estimated by

$$\mathbb{E}[|F_\theta|^{2q}]^{1/2q} \leq Cr(1 + |\xi|^C) |\Xi| q^C \left(\frac{\varepsilon}{r} \right)^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi,\Xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q}$$

for any $0 < \tau \leq c(d, m, \lambda, \Lambda)$ and any $q \geq C$.

We also derive estimates on the averages of the linearized correctors that are analogous to the ones of Lemma 18.

Lemma 22. *Given the assumptions of Lemma 21, for any $0 < \tau \leq c(d, m, \lambda, \Lambda)$, any $x_0 \in \mathbb{R}^d$, any $r \geq \varepsilon$, and any $R \geq r$ we have the estimate*

(40)

$$\begin{aligned} & \mathbb{E} \left[\left| \int_{B_r(x_0)} \phi_{\xi, \Xi}^T dx - \int_{B_R(x_0)} \phi_{\xi, \Xi}^T dx \right|^{2q} \right]^{1/2q} \\ & + \mathbb{E} \left[\left| \int_{B_r(x_0)} \sigma_{\xi, \Xi}^T dx - \int_{B_R(x_0)} \sigma_{\xi, \Xi}^T dx \right|^{2q} \right]^{1/2q} \\ & \leq C(1 + |\xi|)^C |\Xi| q^C \mathbb{E} \left[\left(\frac{r_{*,T,\xi,\Xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q} \left(\sum_{l=0}^{\log_2 \frac{R}{r}} (2^l r)^2 \left(\frac{\varepsilon}{2^l r} \right)^d \right)^{1/2}. \end{aligned}$$

Furthermore, for any $R \geq \sqrt{T}$ we have the bound

$$\begin{aligned} (41) \quad & \mathbb{E} \left[\left| \int_{B_R(x_0)} \phi_{\xi, \Xi}^T dx \right|^q \right]^{1/q} + \mathbb{E} \left[\left| \int_{B_R(x_0)} \sigma_{\xi, \Xi}^T dx \right|^q \right]^{1/q} \\ & \leq C(1 + |\xi|)^C |\Xi| q^C \sqrt{T} \left(\frac{\varepsilon}{R} \right)^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi,\Xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q}. \end{aligned}$$

We then establish moment bounds on the minimal radius $r_{*,T,\xi,\Xi}$.

Lemma 23 (Moment bound on the minimal radius of the linearized equation). *Given the assumptions of Lemma 21, the minimal radius $r_{*,T,\xi,\Xi}$ has stretched exponential moments in the sense*

$$\mathbb{E} \left[\exp \left(\left(\frac{r_{*,T,\xi,\Xi}}{(1+|\xi|^C)\varepsilon} \right)^{1/C} \right) \right] \leq 2$$

for some constant $C = C(d, m, \lambda, \Lambda, \rho, \nu)$.

Note that under a different decorrelation assumption – namely, finite range of dependence as opposed to a spectral gap inequality – stochastic moment bounds for (basically) the quantity $r_{*,\infty,\xi,\Xi}$ have already been established in [2]. The estimates in [2] even achieve optimal stochastic integrability.

A necessary ingredient for the sensitivity estimates for functionals of the linearized corrector $\phi_{\xi,\Xi}^T$ – as derived in Lemma 21 – is the following regularity estimate, which is a consequence of our corrector estimates for the nonlinear problem and the small-scale regularity condition (R).

Lemma 24. *Let the assumptions (A1)–(A3) and (P1)–(P2) be satisfied. Suppose furthermore that the small-scale regularity condition (R) holds. Then there exist $\delta = \delta(d, m, \lambda, \Lambda, \rho, \nu) > 0$ and a stationary nonnegative random field $C_{reg,\xi}$ with uniformly bounded stretched exponential moments $\mathbb{E}[\exp(C_{reg,\xi}(x)^{1/C})] \leq 2$ such that the estimate*

$$(42) \quad \sup_{\tilde{x} \in B_{\varepsilon/2}(x)} |\Xi + \nabla \phi_{\xi,\Xi}^T(\tilde{x})| \leq C_{reg,\xi}(x)(1+|\xi|)^C |\Xi| \left(\frac{r_{*,T,\xi,\Xi}(x)}{\varepsilon} \right)^{(d-\delta)/2}$$

holds for any $x \in \mathbb{R}^d$.

Proof. We use the assumption of small-scale regularity (R) to yield by Proposition 41

$$\sup_{\tilde{x} \in B_{\varepsilon/2}(x)} |\Xi + \nabla \phi_{\xi,\Xi}^T(\tilde{x})| \leq C_{reg,\xi}(x)(1+|\xi|)^C \left(\int_{B_\varepsilon(x)} |\Xi + \nabla \phi_{\xi,\Xi}^T|^2 d\tilde{x} \right)^{1/2}$$

Using (89) to estimate the right-hand side in this equation, we obtain the desired bound. \square

The following result converts estimates on linear functionals of the gradient of a random field and estimates on the gradient of the random field into L^p -estimates for the random field.

Lemma 25 (Estimate on the L^p norm by a multiscale decomposition). *Let $\gamma > 0$, $\varepsilon > 0$, $m \geq 2$, and $K \geq 0$. Let $u = u(\omega_\varepsilon, x)$ be a random function subject to the estimates*

$$(43) \quad \mathbb{E} \left[\left(\int_{B_\varepsilon(x_0)} |\nabla u|^2 dx \right)^{m/2} \right]^{1/m} \leq K \quad \text{for all } x_0 \in \mathbb{R}^d$$

and

$$(44) \quad \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \nabla u \cdot g dx \right)^m \right]^{1/m} \leq K \left(\frac{\varepsilon}{r} \right)^\gamma$$

for all $r \geq 2\varepsilon$, all $x_0 \in \mathbb{R}^d$, and all vector fields $g : \mathbb{R}^d \rightarrow \mathbb{R}^d$ supported in $B_r(x_0)$ satisfying

$$\left(\int_{B_r(x_0)} |g|^{2+1/d} dx \right)^{1/(2+1/d)} \leq r^{-d}.$$

Then estimates of the form

$$\mathbb{E} \left[\left(\int_{B_r(x_0)} \left| u - \int_{B_r(x_0)} u \right|^2 dx \right)^{m/2} \right]^{1/m} \leq \begin{cases} C_3 K \varepsilon^\gamma r^{1-\gamma} & \text{for } \gamma < 1, \\ C_3 K \varepsilon \sqrt{\log(r/\varepsilon)} & \text{for } \gamma = 1, \\ C_3 K \varepsilon & \text{for } \gamma > 1, \end{cases}$$

and

$$\mathbb{E} \left[\left(\int_{B_r(x_0)} \left| u - \int_{B_r(x_0)} u \right|^p dx \right)^{m/p} \right]^{1/m} \leq \begin{cases} C_3 K \varepsilon^\gamma r^{1-\gamma} & \text{for } \gamma < 1, \\ C_3 K \varepsilon \log(r/\varepsilon) & \text{for } \gamma = 1, \\ C_3 K \varepsilon & \text{for } \gamma > 1, \end{cases}$$

hold for all $r \geq \varepsilon$ and all $2 \leq p \leq \frac{2d}{(d-2)_+}$ (as long as $p < \infty$), the constant C_3 depending on γ and p , but being independent of m .

Proof. The L^2 -version of the statement is shown e.g. in [13, Lemma 12]. The L^p version is proven analogously. \square

Finally, note that our main results rely on estimates for the correctors ϕ_ξ and σ_ξ (and not the localized approximations ϕ_ξ^T and σ_ξ^T). While all of our bounds on the localized correctors ϕ_ξ^T and σ_ξ^T are uniform with respect to the parameter $T \geq 2\varepsilon^2$, it remains to justify the passage to the limit $T \rightarrow \infty$.

Lemma 26 (Convergence of the localized correctors in the limit $T \rightarrow \infty$). *Let (A1)–(A3) and (P1)–(P2) be satisfied. Let ϕ_ξ^T and σ_ξ^T be the unique solutions to the localized corrector equations (12a) and (12c). For any $r > 0$, as $T \rightarrow \infty$ the stationary random fields $\nabla \phi_\xi^T$ and $\nabla \sigma_{\xi, jk}^T$ converge strongly in $L^2(\Omega \times B_r)$ to the corrector gradients $\nabla \phi_\xi$ and $\nabla \sigma_{\xi, jk}$, with the correctors ϕ_ξ and σ_ξ as defined in Definition 1.*

3.3. Quantitative two-scale approximation. The estimate on the homogenization error invokes a quantitative two-scale expansion of the homogenized equation. As indicated earlier, for technical reasons, we do not work with the usual expansion $u_{\text{hom}} + \nabla \phi_{\nabla u_{\text{hom}}}$, but consider the (easier) case of a piecewise affine approximation¹ of the form

$$u_{\text{hom}} + \sum_{k \in \frac{\delta}{2}\mathbb{Z}^d} \eta_k \phi_{\xi_k},$$

where $\{\eta_k\}$ denotes a partition of unity subordinate to a uniform cover of \mathbb{R}^d on a scale $\delta \gtrsim \varepsilon$, and $\{\xi_k\}$ denotes the associated piecewise-constant approximation of ∇u_{hom} . Depending on our application, we will choose either $\delta := \varepsilon$ or $\delta := \varepsilon^{1/2}$.

Proposition 27 (Error representation by the two-scale expansion). *Let $A_\varepsilon : \mathbb{R}^d \times \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ be a monotone operator subject to uniform ellipticity and boundedness conditions as in (A1) and (A2). For any $\xi \in \mathbb{R}^{m \times d}$, denote by ϕ_ξ a solution*

¹The authors are indebted to Gilles Francfort for this suggestion, which led to a simplification of the proof.

to the corrector equation (4) on \mathbb{R}^d in the sense of Definition 1 and denote by σ_ξ a solution to the equation for the flux corrector (10) on \mathbb{R}^d .

Let $\delta > 0$. Suppose that either $\mathcal{O} = \mathbb{R}^d$ or that $\mathcal{O} \subset \mathbb{R}^d$ is a bounded domain of class $C^{1,1}$; in the former case, set $K := \delta\mathbb{Z}^d$, and in the latter case set $K := \{k \in \delta\mathbb{Z}^d : k + ((1 - 1/2d)\delta, (1 - 1/2d)\delta)^d \cap \mathcal{O} \neq \emptyset\}$. In the case $\mathcal{O} \neq \mathbb{R}^d$, suppose in addition that $\delta \leq c(\mathcal{O})$ for some $c(\mathcal{O})$ determined in the proof below. Let η_k , $k \in K$, be a partition of unity on \mathcal{O} subordinate to the cover

$$\mathcal{O} \subset \bigcup_{k \in K} (k + (-\delta, \delta)^d)$$

with $0 \leq \eta_k \leq 1$, $\eta_k \equiv 1$ on $\mathcal{O} \cap (k + (-\frac{\delta}{2d}, \frac{\delta}{2d})^d)$, $\eta_k \geq c$ on $\mathcal{O} \cap (k + ((1 - 1/3d)\delta, (1 - 1/3d)\delta)^d)$, and $|\nabla \eta_k| \leq C\delta^{-1}$.

For any given $\bar{u} \in H^2(\mathcal{O})$ and any $k \in K$, introduce the weighted average ξ_k of $\nabla \bar{u}$ near k as

$$\xi_k := \frac{\int_{\mathcal{O}} \eta_k(x) \nabla \bar{u} \, dx}{\int_{\mathcal{O}} \eta_k(x) \, dx}.$$

Define the two-scale expansion for \bar{u} with a piecewise constant approximation for the gradient $\nabla \bar{u}$ as

$$\hat{u} := \bar{u} + \sum_{k \in K} \eta_k \phi_{\xi_k}.$$

Then the action of the operator $-\nabla \cdot (A_\varepsilon(x, \nabla \cdot))$ on \hat{u} is related to the action of the operator $-\nabla \cdot (A_{\text{hom}}(\nabla \cdot))$ on \bar{u} via

$$-\nabla \cdot (A_\varepsilon(x, \nabla \hat{u})) = -\nabla \cdot (A_{\text{hom}}(\nabla \bar{u})) + \nabla \cdot R$$

with a residual $R \in L^2(\mathcal{O}; \mathbb{R}^{m \times d})$ subject to the estimate

$$(45) \quad \int_{\mathcal{O}} |R|^2 \, dx \leq C\delta^2 \int_{\mathcal{O}} |\nabla^2 \bar{u}|^2 \, dx + \frac{C}{\delta^2} \sum_{k, \bar{k} \in K: |k - \bar{k}| \leq d\delta} \int_{B_{5d\delta}(k)} |\phi_{\xi_k} - \phi_{\xi_{\bar{k}}}|^2 + |\sigma_{\xi_k} - \sigma_{\xi_{\bar{k}}}|^2 \, dx$$

with $C = C(d, m, \lambda, \Lambda, \mathcal{O})$. Furthermore, the ξ_k satisfy the estimates

$$(46a) \quad \sum_{k, \bar{k} \in K: |k - \bar{k}| \leq d\delta} |\xi_k - \xi_{\bar{k}}|^2 \int_{B_{d\delta}(k)} 1 \, dx \leq C(\mathcal{O})\delta^2 \int_{\mathcal{O}} |\nabla^2 \bar{u}|^2 \, dx,$$

$$(46b) \quad \sum_{k \in K: |k - \bar{k}| \leq d\delta} |\xi_k|^p \int_{B_{d\delta}(k)} 1 \, dx \leq C(\mathcal{O}, p) \int_{\mathcal{O}} |\nabla \bar{u}|^p \, dx,$$

$$(46c) \quad \sum_{k \in K \cap (\partial\mathcal{O} + B_{5\delta})} |\xi_k|^2 \int_{k + B_{10d\delta}} 1 \, dx \leq C(\mathcal{O}) \sum_{k \in K \cap (\partial\mathcal{O} + B_{5\delta})} \int_{k + B_{10d\delta} \cap \mathcal{O}} |\nabla \bar{u}|^2 \, dx$$

for any $p \geq 2$.

3.4. The approximation of the homogenized operator by periodic representative volumes. In this section we outline the general strategy for the proof of the a priori estimates for the RVE approximation of the effective material law A_{hom} stated in Theorem 9. It is inspired by [30], where the first optimal result for periodic RVEs in a linear discrete setting has been established. In the following we focus on the situation where the small-scale regularity assumption (R) is satisfied, since in that case we can obtain better rates. The argument features additional subtleties (even compared to the linear case of [30]). We start with the observation that the total approximation error decomposes into a random and a systematic part

$$\begin{aligned} & A^{\text{RVE},L}(\xi) - A_{\text{hom}}(\xi) \\ &= \left(A^{\text{RVE},L}(\xi) - \mathbb{E}_L [A^{\text{RVE},L}(\xi)] \right) + \left(\mathbb{E}_L [A^{\text{RVE},L}(\xi)] - A_{\text{hom}}(\xi) \right). \end{aligned}$$

The random error (the first term on the RHS) is a random variable with vanishing expectation and corresponds to the fluctuations of the periodic RVE approximation around its expected value. Theorem 9a asserts that this error decays with the same rate as the fluctuations of a linear average of the random parameter field on scale L , i. e. like $(\frac{L}{\varepsilon})^{-\frac{d}{2}}$. The second term on the RHS is the systematic error. It captures the error coming from approximating the whole-space law \mathbb{P} by the L -periodic law \mathbb{P}_L . As in [30] we decompose the systematic error into different contributions. In particular, we introduce the following notion of localized RVE approximation.

Definition 28 (Localized RVE approximation). *Let A satisfy (A1)–(A3). Let $T \geq 2\varepsilon^2$, let η be a non-negative weight $\eta : \mathbb{R}^d \rightarrow \mathbb{R}$ with $\int_{\mathbb{R}^d} \eta dx = 1$, and let $\tilde{\omega} : \mathbb{R}^d \rightarrow \mathbb{H} \cap B_1$ be a parameter field. We then define the localized RVE approximation of size L with localization parameter $T \in [2\varepsilon^2, L]$ for the effective operator A_{hom} by the expression*

$$(47) \quad A^{\text{RVE},\eta,T}(\tilde{\omega}, \xi) \cdot \Xi := \int_{\mathbb{R}^d} \eta \left(A(\tilde{\omega}, \xi + \nabla \phi_\xi^T) \cdot \Xi - \frac{1}{T} \phi_\xi^T \phi_{\xi,\Xi}^{*,T} \right) dx$$

for any $\xi, \Xi \in \mathbb{R}^{m \times d}$, where $\phi_\xi^T = \phi_\xi^T(\tilde{\omega})$ and $\phi_{\xi,\Xi}^{*,T} = \phi_{\xi,\Xi}^{*,T}(\tilde{\omega})$ denote the localized corrector and the localized linearized adjoint corrector, i. e. ϕ_ξ^T is the unique solution in $H_{\text{uloc}}^1(\mathbb{R}^d; \mathbb{R}^m)$ to the equation (12a) and $\phi_{\xi,\Xi}^{*,T}$ is the unique solution in $H_{\text{uloc}}^1(\mathbb{R}^d; \mathbb{R}^m)$ to the equation (18a) but with a_ξ^T replaced by its transpose $a_\xi^{T,*}$.

If we choose in the previous definition a random field ω_ε with some probability distribution \mathbb{P} subject to (P1)–(P2), the localized RVE approximation $A^{\text{RVE},\eta,T}(\xi)$ converges almost surely for $T \rightarrow \infty$ to the effective material law $A_{\text{hom}}(\xi)$. To see this, one may e. g. use ergodicity and stationarity as well as the sublinear growth of correctors. However, in contrast to the periodic RVE approximation $A^{\text{RVE},L}$ to the effective material law or the homogenized material law A_{hom} itself, the localized RVE can be defined for all parameter fields $\tilde{\omega}$, since it only invokes the localized correctors. This allows to couple parameter fields sampled with \mathbb{P} and \mathbb{P}_L , respectively. More precisely, with the restriction map

$$(48) \quad (\pi_L \tilde{\omega})(x) := \begin{cases} \tilde{\omega}(x) & \text{if } x \in B_{\frac{L}{4}}, \\ 0 & \text{else,} \end{cases}$$

we note that if \mathbb{P}_L is an L -periodic approximation of \mathbb{P} in the sense of Definition 8 and if $\omega_{\varepsilon,L}$ and ω_ε denote random fields distributed according to \mathbb{P}_L respectively

\mathbb{P} , then the equality of laws

$$\pi_L \omega_\varepsilon \sim \pi_L \omega_{\varepsilon, L}$$

holds. Hence, if the weight η of the localized RVE is supported in $B_{\frac{L}{4}}$ and \mathbb{P}_L is a L -periodic approximation of \mathbb{P} , then

$$(49) \quad \mathbb{E}_L [A^{\text{RVE}, \eta, T}(\pi_L \omega_\varepsilon, \xi)] = \mathbb{E} [A^{\text{RVE}, \eta, T}(\pi_L \omega_\varepsilon, \xi)].$$

This identity couples localized approximations for the homogenized material law associated with \mathbb{P} and \mathbb{P}_L . Motivated by this we decompose the systematic error as

$$(50) \quad \begin{aligned} \mathbb{E}_L [A^{\text{RVE}, L}(\xi)] - A_{\text{hom}}(\xi) &= \mathbb{E}_L [A^{\text{RVE}, L}(\xi)] - \mathbb{E}_L [A^{\text{RVE}, \eta, T}(\xi)] \\ &\quad + \mathbb{E}_L [A^{\text{RVE}, \eta, T}(\xi)] - \mathbb{E}_L [A^{\text{RVE}, \eta, T}(\pi_L \omega_\varepsilon, \xi)] \\ &\quad + \mathbb{E} [A^{\text{RVE}, \eta, T}(\pi_L \omega_\varepsilon, \xi)] - \mathbb{E} [A^{\text{RVE}, \eta, T}(\xi)] \\ &\quad + \mathbb{E} [A^{\text{RVE}, \eta, T}(\xi)] - A_{\text{hom}}(\xi). \end{aligned}$$

The differences in the second and third row capture the error coming from replacing ω_ε with $\pi_L \omega_\varepsilon$. As our next lemma shows, this error becomes small upon choosing the localization parameter T suitably. At the heart of the proof of the lemma is the exponential locality of the localized corrector equations (12a) and (18c), see Lemma 36.

Lemma 29 (Estimate for the coupling error). *Let $A : \mathbb{R}^d \times \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ be a monotone operator subject to conditions (A1)-(A2) and subject to the Lipschitz estimate for $\partial_\xi A$ as in (R). Let $L \geq \sqrt{T} \geq 2\varepsilon$ and let η_L denote a non-negative weight supported in $B_{\frac{L}{8}}$ with $|\eta_L| \leq C(d)L^{-d}$ and $|\nabla \eta| \leq C(d)L^{-d-1}$. Then there exist $q = q(d, m, \lambda, \Lambda)$, $\gamma = \gamma(d, m, \lambda, \Lambda)$, and $C = C(d, m, \lambda, \Lambda)$ such that for all parameter fields $\tilde{\omega}$ and all $\xi, \Xi \in \mathbb{R}^{m \times d}$ we have*

$$\begin{aligned} &|A^{\text{RVE}, \eta_L, T}(\tilde{\omega}, \xi) \cdot \Xi - A^{\text{RVE}, \eta_L, T}(\pi_L \tilde{\omega}, \xi) \cdot \Xi| \\ &\leq C \exp\left(-\frac{\gamma}{64} \cdot \frac{L}{\sqrt{T}}\right) \left(|\xi| |\Xi| + (1 + |\xi|) |\xi| |\Xi| + \|\nabla \phi_{\xi, \Xi}^T\|_{q, L, T}\right), \end{aligned}$$

where

$$\begin{aligned} &\|\Xi + \nabla \phi_{\xi, \Xi}^T\|_{q, L, T} \\ &:= \frac{1}{\#X_{L, T}} \sum_{x_0 \in X_{L, T}} \left(\sqrt{T}^{-d} \int_{\mathbb{R}^d} |\Xi + \nabla \phi_{\xi, \Xi}^T(\tilde{\omega}, x)|^q \exp\left(-\frac{\gamma}{2} \frac{|x - x_0|}{\sqrt{T}}\right) dx \right)^{\frac{1}{q}}, \end{aligned}$$

and where $X_{L, T} \subset B_{\frac{L}{8}}$ denotes an arbitrary finite set with cardinality $\#X_{L, T} \leq C(d) \left(\frac{L}{\sqrt{T}}\right)^d$ and $\cup_{x_0 \in X_{L, T}} B_{\sqrt{T}}(x_0) \subset B_{\frac{L}{8}}$.

The differences in the first and last row of the right-hand side in (50) are the *systematic localization errors*, which originate from the localization with parameter T . The systematic localization error can be estimated as follows.

Proposition 30 (Systematic error of localized RVE). *Let $A : \mathbb{H} \times \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ satisfy (A1)-(A3). Let \mathbb{P} be stationary in the sense of assumption (P1). Then the following holds for all $\xi \in \mathbb{R}^{m \times d}$:*

(a) $\mathbb{E} [A^{\text{RVE}, \eta, T}(\xi)]$ as defined in (47) is independent of the weight η .

- (b) Suppose that \mathbb{P} satisfies a spectral gap estimate in the sense of assumption (P2). Assume furthermore that the small-scale regularity condition (R) holds. Then for all $T \geq 2\varepsilon^2$ and all $\xi, \Xi \in \mathbb{R}^{m \times d}$ we have

$$\begin{aligned} & |A_{\text{hom}}(\xi) \cdot \Xi - \mathbb{E} [A^{\text{RVE}, \eta, T}(\xi) \cdot \Xi] | \\ & \leq C(1 + |\xi|)^C |\xi| |\Xi| \left(\frac{\varepsilon}{\sqrt{T}} \right)^{d \wedge 4} \times \begin{cases} |\log(\sqrt{T}/\varepsilon)|^2 & \text{for } d = 2 \text{ and } d = 4, \\ 1 & \text{for } d = 3, d = 1, \text{ and } d \geq 5 \end{cases} \end{aligned}$$

with a constant $C = C(d, m, \lambda, \Lambda, \rho, \nu)$.

- (c) If \mathbb{P} concentrates on L -periodic parameter fields and satisfies a periodic spectral gap estimate in the sense of Definition 8b as well as the regularity condition (R), then we have for all $T \in [2\varepsilon^2, L^2]$ and all $\xi, \Xi \in \mathbb{R}^{m \times d}$

$$\begin{aligned} & |\mathbb{E} [A^{\text{RVE}, \eta}(\xi) \cdot \Xi] - \mathbb{E} [A^{\text{RVE}, \eta, T}(\xi) \cdot \Xi] | \\ & \leq C(1 + |\xi|)^C |\xi| |\Xi| \left(\frac{\varepsilon}{\sqrt{T}} \right)^{d \wedge 4} \times \begin{cases} |\log(\sqrt{T}/\varepsilon)|^2 & \text{for } d = 2 \text{ and } d = 4, \\ 1 & \text{for } d = 3, d = 1, \text{ and } d \geq 5. \end{cases} \end{aligned}$$

In order to prove this result, we need to quantify the systematic error on the level of the correctors. Note that this estimate is slightly pessimistic (by the logarithmic factor) for $d = 2$ and $d = 4$. Moreover, note that for $d \geq 5$ the estimate saturates, an effect that is also observed in the stochastic homogenization of linear elliptic PDEs, see [30].

Lemma 31 (Localization error in the corrector). *Let A satisfy the assumptions (A1)–(A3) and let \mathbb{P} satisfy the assumptions (P1)–P(2). Then for all $T \geq 2\varepsilon^2$ and all $x_0 \in \mathbb{R}^d$ we have*

$$\begin{aligned} & \left(\int_{B_{\sqrt{T}}(x_0)} \left(|\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T|^2 + \frac{1}{T} |\phi_\xi^{2T} - \phi_\xi^T|^2 \right) dx \right)^{\frac{1}{2}} \\ & \leq C |\xi| \left(\frac{\varepsilon}{\sqrt{T}} \right)^{\frac{d \wedge 4}{2}} \times \begin{cases} |\log(\sqrt{T}/\varepsilon)|^{1/2} & \text{for } d \in \{2, 4\}, \\ 1 & \text{for } d = 3, d = 1, \text{ and } d \geq 5. \end{cases} \end{aligned}$$

Here C denotes a random constant as in Proposition 14.

With help of Meyers' estimate we may upgrade the previous estimate to an L^p bound.

Corollary 32. *Consider the setting of Lemma 31. Then there exists a Meyer's exponent p with $p \in (2, 2 + c(d, m, \lambda, \Lambda)]$ such that the estimate*

$$\begin{aligned} & \mathbb{E} [|\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T|^p]^{\frac{1}{p}} \\ & \leq C |\xi| \left(\frac{\varepsilon}{\sqrt{T}} \right)^{\frac{d \wedge 4}{2}} \times \begin{cases} |\log(\sqrt{T}/\varepsilon)| & \text{for } d \in \{2, 4\}, \\ 1 & \text{for } d = 3, d = 1, \text{ and } d \geq 5 \end{cases} \end{aligned}$$

holds, where $C = C(d, m, \lambda, \Lambda, \rho)$.

We also require control of the localization error for the linearized corrector.

Lemma 33 (Localization error for linearized corrector). *Consider the setting of Lemma 31. Furthermore, assume that the small-scale regularity condition (R) holds.*

Then

$$\begin{aligned} & \mathbb{E} \left[\left| \nabla \phi_{\xi, \Xi}^T - \nabla \phi_{\xi, \Xi}^{2T} + \frac{1}{T} |\phi_{\xi, \Xi}^T - \phi_{\xi, \Xi}^{2T}|^2 \right]^{1/2} \\ & \leq C(1 + |\xi|)^C |\Xi| \left(\frac{\varepsilon}{\sqrt{T}} \right)^{\frac{d\Delta 4}{2}} \times \begin{cases} |\log(\sqrt{T}/\varepsilon)| & \text{for } d \in \{2, 4\}, \\ 1 & \text{for } d = 3, d = 1, \text{ and } d \geq 5, \end{cases} \end{aligned}$$

where $C = C(d, m, \lambda, \Lambda, \rho, \nu)$.

4. PROOF OF THE MAIN RESULTS

4.1. Ingredients from regularity theory. Our estimates crucially rely on three basic regularity estimates for elliptic PDEs, the first two being the Caccioppoli inequality and the hole-filling estimate for *nonlinear elliptic equations* (and systems) with monotone nonlinearity and the last one being a weighted Meyers' estimate for linear elliptic equations (and systems). We first state the Caccioppoli inequality and the hole-filling estimates in the nonlinear setting. The (standard) proofs are provided in Appendix A.

Lemma 34 (Caccioppoli inequality and hole-filling estimate for monotone systems). *Let $A(x, \xi)$ be a monotone operator subject to the assumptions (A1)–(A2). Let $0 < T \leq \infty$ and let u be a solution to the system of PDEs*

$$-\nabla \cdot (A(x, \nabla u)) + \frac{1}{T} u = \nabla \cdot g + \frac{1}{T} f$$

for some $f \in L^2(\mathbb{R}^d; \mathbb{R}^m)$. Then there exist constants $C > 0$ and $\delta > 0$ depending only on d, m, λ , and Λ with the following property: For any $R, r > 0$ with $R \geq r$ we have the Caccioppoli inequality

$$\begin{aligned} (51) \quad & \int_{B_{R/2}(x_0)} |\nabla u|^2 + \frac{1}{T} |u|^2 dx \\ & \leq \frac{C}{R^2} \int_{B_R(x_0)} |u - b|^2 dx + \frac{C}{T} |b|^2 + C \int_{B_R(x_0)} |g|^2 + \frac{1}{T} |f|^2 dx \end{aligned}$$

for any $b \in \mathbb{R}^m$ and the hole-filling estimate

$$\begin{aligned} (52) \quad & \int_{B_r(x_0)} |\nabla u|^2 dx + \frac{1}{T} \int_{B_r(x_0)} |u|^2 dx \\ & \leq C \left(\frac{r}{R} \right)^\delta \left(\int_{B_R(x_0)} |\nabla u|^2 dx + \frac{1}{T} \int_{B_R(x_0)} |u|^2 dx \right) \\ & \quad + C \int_{B_R(x_0)} \left(\frac{r}{r + |x - x_0|} \right)^\delta \left(|g|^2 + \frac{1}{T} |f|^2 \right) dx. \end{aligned}$$

We next state a weighted Meyers-type estimate for linear uniformly elliptic equations and systems. Its proof (which is provided in Appendix C) relies on the usual Meyers estimate, along with a duality argument and a hole-filling estimate for the adjoint operator. The details are provided in [13] for the case $T = \infty$; however, the proof applies verbatim to the case $T > 0$, as the only ingredients are the Meyers' estimate for the PDE and the hole-filling estimate for the adjoint PDE.

Lemma 35 (A weighted Meyers estimate for linear elliptic systems). *Let $a : \mathbb{R}^d \rightarrow (\mathbb{R}^{m \times d})^2$ be a uniformly elliptic and bounded coefficient field with ellipticity and boundedness constants λ and Λ . Let $r > 0$ be arbitrary. Let $v \in H^1(\mathbb{R}^d; \mathbb{R}^m)$ and $g \in L^2(\mathbb{R}^d; \mathbb{R}^{m \times d})$, $f \in L^2(\mathbb{R}^d; \mathbb{R}^m)$ be functions related through*

$$-\nabla \cdot (a \nabla v) + \frac{1}{T}v = \nabla \cdot g + \frac{1}{\sqrt{T}}f.$$

There exists a Meyers exponent $\bar{p} > 2$ and a constant $c > 0$, which both only depend on d, m, λ , and Λ , such that for all $2 \leq p < \bar{p}$, all $\alpha_1 < c$, and all $0 < \alpha_0 < \alpha_1$ we have

$$(53) \quad \left(\int_{\mathbb{R}^d} \left(|\nabla v|^p + \left| \frac{1}{\sqrt{T}}v \right|^p \right) \left(1 + \frac{|x|}{r} \right)^{\alpha_0} dx \right)^{\frac{1}{p}} \leq C \left(\int_{\mathbb{R}^d} (|g|^p + |f|^p) \left(1 + \frac{|x|}{r} \right)^{\alpha_1} dx \right)^{\frac{1}{p}},$$

where the constant C depends only on $d, m, \lambda, \Lambda, p, \alpha_0$, and α_1 .

The localization ansatz for the correctors relies crucially on the following elementary deterministic energy estimate with exponential localization. As the proof is short and elementary, we directly provide it here.

Lemma 36 (Exponential localization). *Suppose that $A : \mathbb{R}^d \times \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ is a monotone operator satisfying (A1) and (A2). Let $T > 0$ and $L \geq \sqrt{T}$. Consider $u \in H_{\text{loc}}^1(\mathbb{R}^d; \mathbb{R}^m)$ and $f \in L_{\text{loc}}^2(\mathbb{R}^d; \mathbb{R}^m)$, $F \in L_{\text{loc}}^2(\mathbb{R}^d; \mathbb{R}^{m \times d})$ related by*

$$-\nabla \cdot (A(x, \nabla u)) + \frac{1}{T}u = \nabla \cdot F + \frac{1}{T}f$$

in a distributional sense in \mathbb{R}^d . Suppose that u, f , and F have at most polynomial growth in the sense that

$$\exists k \in \mathbb{N} : \quad \limsup_{R \rightarrow \infty} R^{-k} \left(\int_{B_R} (|u| + |\nabla u| + |f| + |F|)^2 \right)^{\frac{1}{2}} = 0.$$

Then for $0 < \gamma \leq c(d, m, \lambda, \Lambda)$ we have

$$\begin{aligned} & \int_{\mathbb{R}^d} \left(|\nabla u|^2 + \frac{1}{T}|u|^2 \right) \exp(-\gamma|x|/L) dx \\ & \leq C(d, m, \lambda, \Lambda) \int_{\mathbb{R}^d} \left(|F|^2 + \frac{1}{T}|f|^2 \right) \exp(-\gamma|x|/L) dx. \end{aligned}$$

Proof. Set $\eta(x) := \exp(-\gamma|x|/L)$. We test the equation with $u\eta$ (which can be justified by approximation thanks to the polynomial growth assumption). By an integration by parts, the ellipticity and Lipschitz continuity of A , and using $|\nabla \eta| \leq \gamma/L\eta \leq \gamma/\sqrt{T}\eta$, we get

$$\begin{aligned} & \lambda \int_{\mathbb{R}^d} |\nabla u|^2 \eta + \frac{1}{T} \int_{\mathbb{R}^d} |u|^2 \eta \\ & \leq \gamma \int_{\mathbb{R}^d} \Lambda |\nabla u| \frac{1}{\sqrt{T}} |u| \eta dx + \int_{\mathbb{R}^d} |F| \left(|\nabla u| + \gamma \frac{1}{\sqrt{T}} |u| \right) \eta dx + \frac{1}{T} \int_{\mathbb{R}^d} |f| |u| \eta dx \end{aligned}$$

The claim now follows for $\gamma \leq c$ by absorbing the terms with u and ∇u on the right-hand side into the left-hand side with help of Young's inequality. \square

Remark 37. We frequently apply the exponential localization in the following form: Suppose that $A : \mathbb{R}^d \times \mathbb{R}^{m \times d} \rightarrow \mathbb{R}^{m \times d}$ is a monotone operator satisfying (A1) and (A2). Let $T > 0$ and $L \geq \sqrt{T}$. Consider $u_1, u_2 \in H_{\text{loc}}^1(\mathbb{R}^d; \mathbb{R}^m)$ and $f \in L_{\text{loc}}^2(\mathbb{R}^d; \mathbb{R}^m)$, $F \in L_{\text{loc}}^2(\mathbb{R}^d; \mathbb{R}^{m \times d})$, all with polynomial growth and related by

$$-\nabla \cdot (A(x, \nabla u_1) - A(x, \nabla u_2)) + \frac{1}{T}(u_1 - u_2) = \nabla \cdot F + \frac{1}{T}f$$

in a distributional sense in \mathbb{R}^d . Then for $0 < \gamma \leq c(d, m, \lambda, \Lambda)$ we have

$$\begin{aligned} & \int_{\mathbb{R}^d} \left(|\nabla u_1 - \nabla u_2|^2 + \frac{1}{T}|u_1 - u_2|^2 \right) \exp(-\gamma|x|/L) dx \\ & \leq C(d, m, \lambda, \Lambda) \int_{\mathbb{R}^d} \left(|F|^2 + \frac{1}{T}|f|^2 \right) \exp(-\gamma|x|/L) dx. \end{aligned}$$

Indeed, with $a(x) := \int_0^1 \partial_\xi A(x, \xi + (1-s)\nabla u_1(x) + s\nabla u_2(x)) ds$ and $\delta u := u_1 - u_2$, we have $-\nabla \cdot (a(x)\nabla \delta u) + \frac{1}{T}\delta u = \nabla \cdot F + \frac{1}{T}f$. Since A is a monotone operator satisfying (A1) and (A2), the derivative $\partial_\xi A(x, \xi)$ is a uniformly elliptic matrix field; hence, $a(x)$ is a uniformly elliptic coefficient field and the claimed estimate follows from Lemma 36.

4.2. The convergence rate of the solutions. We first provide the proof of the error estimate for $\|u_\varepsilon - u_{\text{hom}}\|_{L^2}$. It is based on a two-scale expansion with a piecewise constant approximation for the slope of the limiting solution u_{hom} , whose approximation properties are stated in Proposition 27.

Proof of Proposition 27. Note that a simple application of the Poincaré inequality implies an estimate on the local approximation error $\nabla \bar{u} - \xi_k$ and on the differences $\xi_k - \xi_{\tilde{k}}$ between nearby lattice sites $k \in K$ and $\tilde{k} \in K \cap (k + [-\delta, \delta]^d)$ of the form

$$(54) \quad \begin{aligned} & \sum_{\tilde{k} \in k + [-\delta, \delta]^d \cap K} |\xi_k - \xi_{\tilde{k}}|^2 + \int_{k + [-\delta, \delta]^d \cap \mathcal{O}} |\nabla \bar{u} - \xi_k|^2 dx \\ & \leq C\delta^2 \int_{k + (-4\delta, 4\delta)^d \cap \mathcal{O}} |\nabla^2 \bar{u}|^2 dx \end{aligned}$$

for each $k \in K$. From this estimate and a straightforward estimate, the bounds (46) are immediate. Furthermore, this enables us to relate the action $\nabla \cdot (A_\varepsilon(x, \nabla \hat{u}))$ of the monotone operator A_ε on the two-scale expansion \hat{u} to the action $\nabla \cdot (A_{\text{hom}}(\bar{u}))$ of the homogenized operator A_{hom} on the (sufficiently smooth) function \bar{u} . To see this, we first compute by adding zero

$$\begin{aligned} & \nabla \cdot (A_\varepsilon(x, \nabla \hat{u})) \\ & = \nabla \cdot \left(A_\varepsilon \left(x, \nabla \bar{u} + \sum_{k \in K} \eta_k \nabla \phi_{\xi_k} \right) \right) \\ & \quad + \nabla \cdot \left(A_\varepsilon \left(x, \nabla \bar{u} + \sum_{k \in K} \eta_k \nabla \phi_{\xi_k} + \sum_{k \in K} \phi_{\xi_k} \nabla \eta_k \right) \right) \\ & \quad - A_\varepsilon \left(x, \nabla \bar{u} + \sum_{k \in K} \eta_k \nabla \phi_{\xi_k} \right). \end{aligned}$$

Adding zero several times and using the fact that the η_k form a decomposition of unity (i. e. $\sum_k \eta_k = 1$), we get

$$\begin{aligned}
& \nabla \cdot (A_\varepsilon(x, \nabla \hat{u})) \\
&= \nabla \cdot (A_{\text{hom}}(\nabla \bar{u})) \\
&+ \nabla \cdot \left(\sum_{k \in K} \eta_k (A_{\text{hom}}(\xi_k) - A_{\text{hom}}(\nabla \bar{u})) \right) \\
&+ \nabla \cdot \left(\sum_{k \in K} \eta_k (A_\varepsilon(x, \xi_k + \nabla \phi_{\xi_k}) - A_{\text{hom}}(\xi_k)) \right) \\
&+ \nabla \cdot \left(\sum_{k \in K} \eta_k (A_\varepsilon(x, \nabla \bar{u} + \nabla \phi_{\xi_k}) - A_\varepsilon(x, \xi_k + \nabla \phi_{\xi_k})) \right) \\
&+ \nabla \cdot \left(A_\varepsilon \left(x, \nabla \bar{u} + \sum_{k \in K} \eta_k \nabla \phi_{\xi_k} \right) - \sum_{k \in K} \eta_k A_\varepsilon \left(x, \nabla \bar{u} + \nabla \phi_{\xi_k} \right) \right) \\
&+ \nabla \cdot \left(A_\varepsilon \left(x, \nabla \bar{u} + \sum_{k \in K} \eta_k \nabla \phi_{\xi_k} + \sum_{k \in K} \phi_{\xi_k} \nabla \eta_k \right) \right. \\
&\quad \left. - A_\varepsilon \left(x, \nabla \bar{u} + \sum_{k \in K} \eta_k \nabla \phi_{\xi_k} \right) \right).
\end{aligned}$$

This yields using the equation for the flux corrector $\nabla \cdot \sigma_{\xi_k} = A_\varepsilon(x, \xi_k + \nabla \phi_{\xi_k}) - A_{\text{hom}}(\xi_k)$ (see (10)) and the skew-symmetry of σ_{ξ_k} (which implies $\nabla \cdot (\eta \nabla \cdot \sigma_{\xi_k}) = -\nabla \cdot (\sigma_{\xi_k} \nabla \eta)$)

$$\begin{aligned}
& \nabla \cdot (A_\varepsilon(x, \nabla \hat{u})) \\
&= \nabla \cdot (A_{\text{hom}}(\nabla \bar{u})) + \nabla \cdot I + \nabla \cdot II + \nabla \cdot III
\end{aligned}$$

with

$$\begin{aligned}
I &:= \sum_{k \in K} \eta_k (A_{\text{hom}}(\xi_k) - A_{\text{hom}}(\nabla \bar{u})) \\
&+ \sum_{k \in K} \eta_k (A_\varepsilon(x, \nabla \bar{u} + \nabla \phi_{\xi_k}) - A_\varepsilon(x, \xi_k + \nabla \phi_{\xi_k}))
\end{aligned}$$

and

$$\begin{aligned}
II &:= - \sum_{k \in K} \sigma_{\xi_k} \nabla \eta_k \\
&+ A_\varepsilon \left(x, \nabla \bar{u} + \sum_{k \in K} \eta_k \nabla \phi_{\xi_k} + \sum_{k \in K} \phi_{\xi_k} \nabla \eta_k \right) - A_\varepsilon \left(x, \nabla \bar{u} + \sum_{k \in K} \eta_k \nabla \phi_{\xi_k} \right)
\end{aligned}$$

as well as

$$III := A_\varepsilon \left(x, \nabla \bar{u} + \sum_{k \in K} \eta_k \nabla \phi_{\xi_k} \right) - \sum_{k \in K} \eta_k A_\varepsilon \left(x, \nabla \bar{u} + \nabla \phi_{\xi_k} \right).$$

We next show that this relation may be simplified to

$$\nabla \cdot (A(x, \nabla \hat{u})) = \nabla \cdot (A_{\text{hom}}(\nabla \bar{u})) + \nabla \cdot R$$

with an error term R estimated by

$$(55) \quad \begin{aligned} & \int_{\mathcal{O}} |R|^2 dx \\ & \leq C\delta^2 \sum_{k \in K} \int_{k+(-2\delta, 2\delta)^d \cap \mathcal{O}} |\nabla^2 \bar{u}|^2 dx \\ & \quad + \frac{C}{\delta^2} \sum_{k \in K} \sum_{\bar{k} \in k+[-\delta, \delta]^d \cap K} \int_{k+(-3\delta, 3\delta)^d} |\phi_{\xi_k} - \phi_{\xi_{\bar{k}}}|^2 + |\sigma_{\xi_k} - \sigma_{\xi_{\bar{k}}}|^2 dx. \end{aligned}$$

To see this, first observe that $\int_{\mathcal{O}} |I|^2 dx$ may be bounded by the first term on the right-hand side of (55): Indeed, using (54), the assumption of Lipschitz continuity of A_ε in ξ (see (A2)), the Lipschitz continuity of A_{hom} (see Theorem 6a), the fact that $\delta \leq c(\mathcal{O})$ in case $\mathcal{O} \neq \mathbb{R}^d$, and the support property $\text{supp } \eta_k \subset k + (-\delta, \delta)^d$, one verifies that $\int_{\mathcal{O}} |I|^2 dx$ is of the desired order.

To see that the term $\int_{\mathcal{O}} |II|^2 dx$ may be bounded by the second term on the right-hand side of (55), one uses the fact that $\sum_k \nabla \eta_k = 0$ (as the η_k are a partition of unity) along with the estimate $|\nabla \eta| \leq C\delta^{-1}$ as well as the Lipschitz continuity of A_ε in ξ (see (A2)) to deduce $|II(x)| \leq C\delta^{-1} \max_{k, \bar{k} \in x+(-\delta, \delta)^d} (|\phi_{\xi_k} - \phi_{\xi_{\bar{k}}}| + |\sigma_{\xi_k} - \sigma_{\xi_{\bar{k}}}|)(x)$. This directly implies the estimate for the contribution of II in (55).

Finally, to estimate the third term III , we first need to bound the difference of two corrector gradients $\nabla \phi_{\xi_k}$ and $\nabla \phi_{\xi_l}$ for nearby grid points k and l with $l \in k + [-\delta, \delta]^d$. To this aim, we compute using the corrector equation (4)

$$-\nabla \cdot (A_\varepsilon(x, \xi_k + \nabla \phi_{\xi_k}) - A_\varepsilon(x, \xi_l + \nabla \phi_{\xi_l})) = 0$$

which yields with an energy estimate using the assumptions (A1) and (A2)

$$\int_{x+(-\delta, \delta)^d} |\nabla \phi_{\xi_k} - \nabla \phi_{\xi_l}|^2 dx \leq C|\xi_k - \xi_l|^2 + \frac{C}{\delta^2} \int_{x+(-2\delta, 2\delta)^d} |\phi_{\xi_k} - \phi_{\xi_l}|^2 dx.$$

In conjunction with the Lipschitz continuity of A_ε in ξ (see (A2)) and (54), this entails that $\int_{\mathcal{O}} |III|^2 dx$ may be bounded by the right-hand side of (55).

We then observe that (55) directly implies the desired bound (45). \square

We now establish our main theorems on the estimate of the homogenization error $\|u_\varepsilon - u_{\text{hom}}\|_{L^p}$. As the proofs are highly similar, we combine the proofs of Theorem 2, Theorem 3, and Theorem 4.

Proof of Theorem 2, Theorem 3, and Theorem 4. Step 1: Estimates on u_{hom} . We intend to derive the estimates on the homogenization error by making use of the error bound for the two-scale expansion from Proposition 27. For this reason, we need an estimate on the second derivative of u_{hom} . We first treat the case $d \geq 3$. Differentiating the effective equation $-\nabla \cdot (A_{\text{hom}}(\nabla u_{\text{hom}})) = f$ with respect to the spatial coordinate x_i , we deduce by adding zero (using the convention $u_{Dir} = 0$ in case of the full space $\mathcal{O} = \mathbb{R}^d$)

$$(56) \quad \begin{aligned} & -\nabla \cdot (\partial_\xi A_{\text{hom}}(\nabla u_{\text{hom}}) \nabla \partial_i (u_{\text{hom}} - u_{Dir})) \\ & = \partial_i f + \nabla \cdot (\partial_\xi A_{\text{hom}}(\nabla u_{\text{hom}}) \nabla \partial_i u_{Dir}) \end{aligned}$$

in the full space \mathbb{R}^d respectively in the interior of the domain \mathcal{O} . Note that the coefficient field $a(x) := \partial_\xi A_{\text{hom}}(\nabla u_{\text{hom}}(x))$ is uniformly elliptic and bounded by the

structure properties of A_{hom} stated in Theorem 6a. An energy estimate for the previous equation therefore yields for any $\tau > 0$

$$(57) \quad \|\nabla \partial_i u_{\text{hom}}\|_{L^2(\{x \in \mathcal{O} : \text{dist}(x, \partial \mathcal{O}) \geq \tau\})}^2 \leq C(d, m, \lambda, \Lambda, \tau, \mathcal{O})(\|f\|_{L^2(\mathcal{O})}^2 + \|\nabla^2 u_{Dir}\|_{L^2(\mathcal{O})}^2).$$

Note that while this argument requires local H^2 regularity of u_{hom} , it can easily be justified for $u_{\text{hom}} \in H^1$ by the difference quotient method (yielding $u_{\text{hom}} \in H^2$ locally as a result). By the usual argument – a local change of coordinates, differentiation at first only in tangential directions, and using the equation itself to estimate the second derivative in normal direction – the estimate can also be easily extended up to the boundary of \mathcal{O} , with a domain-dependent constant. In total, we obtain

$$(58) \quad \int_{\mathcal{O}} |\nabla^2 u_{\text{hom}}|^2 dx \leq C(d, m, \lambda, \Lambda, \mathcal{O})(\|f\|_{L^2(\mathcal{O})}^2 + \|\nabla^2 u_{Dir}\|_{L^2(\mathcal{O})}^2).$$

Furthermore, testing the homogenized PDE (2) with $u_{\text{hom}} - u_{Dir}$ and using the Sobolev embedding as well as Young's inequality yields (at least for $d \geq 3$) the bound

$$(59a) \quad \int_{\mathcal{O}} |\nabla u_{\text{hom}}|^2 dx \leq C(d, m, \lambda, \Lambda)(\|f\|_{L^{2d/(d+2)}(\mathcal{O})}^2 + \|\nabla u_{Dir}\|_{L^2(\mathcal{O})}^2).$$

In the cases $d = 2$ and $d = 1$, the estimate (58) is also valid, the proof being entirely analogous. Due to the massive term, for $d = 2$ and $d = 1$ a weighted energy estimate yields the bound

$$(59b) \quad \int_{\mathcal{O}} |\nabla u_{\text{hom}}|^2 (1 + |x|)^{2\kappa} dx \leq C(d, m, \lambda, \Lambda)(\|(1 + |x|)^\kappa f\|_{L^2(\mathcal{O})}^2 + \|\nabla u_{Dir}\|_{L^2(\mathcal{O})}^2).$$

Step 2: Proof of Theorem 4. In order to establish Theorem 4, we apply Proposition 27 with $\delta := \varepsilon^{1/2}$ in case $d \geq 3$ and $\delta := \varepsilon^{1/2} |\log \varepsilon|^{1/4}$ in case $d = 2$ as well as $\bar{u} := u_{\text{hom}}$ to deduce

$$(60) \quad \nabla \cdot (A(\omega_\varepsilon, \nabla u_\varepsilon) - A(\omega_\varepsilon, \nabla \hat{u})) = \nabla \cdot R$$

with R subject to the bound in (45). From Corollary 15 we deduce the estimate

$$(61) \quad \left(\int_{B_r(x_0)} |\phi_\xi|^2 + |\sigma_\xi|^2 dx \right)^{1/2} \leq \begin{cases} C|\xi|\varepsilon |\log \frac{1}{\varepsilon}|^{1/2} & \text{for } d = 2, \\ C|\xi|\varepsilon & \text{for } d \geq 3, \end{cases}$$

for any $r \geq \varepsilon$, any $x_0 \in \mathcal{O}$, and any $\xi \in \mathbb{R}^{m \times d}$. Plugging in this bound as well as (58) and (59a) into (45), we see that we have the bounds

$$(62) \quad \int_{\mathcal{O}} |R|^2 dx \leq \begin{cases} C\varepsilon |\log \varepsilon|^{1/2} (\|f\|_{L^2(\mathcal{O})}^2 + \|\nabla^2 u_{Dir}\|_{L^2(\mathcal{O})}^2) & \text{for } d = 2, \\ C\varepsilon (\|f\|_{L^2(\mathcal{O})}^2 + \|\nabla^2 u_{Dir}\|_{L^2(\mathcal{O})}^2) & \text{for } d \geq 3, \end{cases}$$

with a random constant C with uniformly bounded stretched exponential moments in the sense $\mathbb{E}[\exp(C^{1/C(d, m, \lambda, \Lambda, \rho)/C}(\mathcal{O}))] \leq 2$.

Testing the equation (60) with the difference $(u_\varepsilon - \hat{u} + (1 - \psi)(\hat{u} - \bar{u})) \in H_0^1(\mathcal{O})$ with some nonnegative cutoff ψ with $\psi \equiv 1$ in $\{x \in \mathcal{O} : \text{dist}(x, \partial \mathcal{O}) > \mu\}$, $\psi = 0$ on

$\partial\mathcal{O}$, and $|\nabla\psi| \leq C\mu^{-1}$, we get by the monotonicity property (A1) and the Lipschitz continuity (A2)

$$\begin{aligned} \int_{\mathcal{O}} \lambda |\nabla u_\varepsilon - \nabla \hat{u}|^2 dx &\leq - \int_{\mathcal{O}} R \cdot \nabla (u_\varepsilon - \hat{u}) dx \\ &\quad + \Lambda \int_{(\partial\mathcal{O} + B_\mu) \cap \mathcal{O}} |\nabla u_\varepsilon - \nabla \hat{u}| |\nabla((1-\psi)(\hat{u} - \bar{u}))| dx \\ &\quad - \int_{(\partial\mathcal{O} + B_\mu) \cap \mathcal{O}} R \cdot \nabla((1-\psi)(\hat{u} - \bar{u})) dx. \end{aligned}$$

This entails the estimate

$$\begin{aligned} \int_{\mathcal{O}} |\nabla u_\varepsilon - \nabla \hat{u} - \nabla((1-\psi)(\bar{u} - \hat{u}))|^2 dx &\leq C \int_{\mathcal{O}} |R|^2 dx + C \int_{\partial\mathcal{O} + B_\mu} |\nabla \hat{u} - \nabla \bar{u}|^2 dx \\ &\quad + C\mu^{-2} \int_{\partial\mathcal{O} + B_\mu} |\hat{u} - \bar{u}|^2 dx. \end{aligned}$$

This implies by the definition of \hat{u}

$$\begin{aligned} &\int_{\mathcal{O}} |\nabla u_\varepsilon - \nabla \hat{u} - \nabla((1-\psi)(\bar{u} - \hat{u}))|^2 dx \\ &\leq C \int_{\mathcal{O}} |R|^2 dx + C \int_{(\partial\mathcal{O} + B_\mu) \cap \mathcal{O}} \sum_{k \in K \cap x + [-\delta, \delta]^d} (|\nabla \phi_{\xi_k}|^2 + \mu^{-2} |\phi_{\xi_k}|^2) dx. \end{aligned}$$

We next observe that by the last estimate in (46), a trace-type inequality, and the estimates (59a) respectively (59b) we have

$$\begin{aligned} \sum_{k \in K \cap \partial\mathcal{O} + B_{2d\delta}} \delta^d |\xi_k|^2 &\leq C \int_{\partial\mathcal{O} + B_{5d\delta} \cap \mathcal{O}} |\nabla u_{\text{hom}}|^2 dx \\ &\leq C(\lambda, \Lambda, \mathcal{O}) \delta (\|f\|_{L^2(\mathcal{O})}^2 + \|\nabla^2 u_{Dir}\|_{L^2(\mathcal{O})}^2). \end{aligned}$$

Plugging in this estimate, the estimate on ϕ_ξ from (61), the Caccioppoli inequality (51) for ϕ_ξ , as well as the bound (62) into the previous estimate, we deduce upon choosing $\mu := \varepsilon^{1/2}$ in case $d \geq 3$ and $\mu := \varepsilon^{1/2} |\log \varepsilon|^{1/4}$ in case $d = 2$

$$\begin{aligned} &\int_{\mathcal{O}} |\nabla u_\varepsilon - \nabla \hat{u} - \nabla((1-\psi)(\bar{u} - \hat{u}))|^2 dx \\ &\leq \begin{cases} C\varepsilon |\log \varepsilon|^{1/2} (\|f\|_{L^2(\mathcal{O})}^2 + \|\nabla^2 u_{Dir}\|_{L^2(\mathcal{O})}^2) & \text{for } d = 2, \\ C\varepsilon (\|f\|_{L^2(\mathcal{O})}^2 + \|\nabla^2 u_{Dir}\|_{L^2(\mathcal{O})}^2) & \text{for } d \geq 3. \end{cases} \end{aligned}$$

Using the Poincaré inequality, we finally deduce

$$\begin{aligned} &\left\| u_\varepsilon - u_{\text{hom}} - \psi \sum_k \eta_k \phi_{\xi_k} \right\|_{L^2(\mathcal{O})} \\ &\leq \begin{cases} C\varepsilon^{1/2} |\log \varepsilon|^{1/4} (\|f\|_{L^2(\mathcal{O})} + \|\nabla^2 u_{Dir}\|_{L^2(\mathcal{O})}) & \text{for } d = 2, \\ C\varepsilon^{1/2} (\|f\|_{L^2(\mathcal{O})} + \|\nabla^2 u_{Dir}\|_{L^2(\mathcal{O})}) & \text{for } d \geq 3. \end{cases} \end{aligned}$$

Using again the estimate on ϕ_ξ from (61), the L^2 estimate for the ξ_k in (46), and the energy estimate (59a), we obtain the statement of Theorem 4.

Step 3: Proof of Theorem 2 and Theorem 3. We will crucially use the improved estimate on the difference of correctors $\phi_{\xi_k} - \phi_{\xi_k^-}$ and $\sigma_{\xi_k} - \sigma_{\xi_k^-}$ provided by Corollary 15; however, as these estimates grow with a factor of $(1 + |\xi_1|^C + |\xi_2|^C)$,

we first need an L^∞ estimate for the gradient ∇u_{hom} . In the scalar case, this may be obtained by using Moser iteration for the differentiated equation (56), while in the vector-valued case we have to appeal to the Uhlenbeck structure of the limiting equation (and hence, in essence, use the fact that $|\nabla u_{\text{hom}}|^2$ is a subsolution to a suitable elliptic PDE, which allows one to apply Moser iteration again). In the scalar setting $m = 1$, by applying [28, Theorem 8.15] to (56) we obtain the estimate

$$(63) \quad \|\partial_i u_{\text{hom}}\|_{L^\infty(\mathbb{R}^d)} \leq C(\|f\|_{L^p(\mathbb{R}^d)} + \|\partial_i u_{\text{hom}}\|_{L^2(\mathbb{R}^d)})$$

for $p > d$ and all i . In the vector-valued case $m \geq 2$, we may use the assumption (see (R)) that the effective operator A_{hom} is of Uhlenbeck structure; hence, $|\nabla u_{\text{hom}}|^2$ is the subsolution of an elliptic PDE (see [41]) and we obtain an analogous result. Finally, in the case $d \leq 2$ we may apply Meyers' estimate (Lemma 35) to obtain an L^p estimate for $\nabla^2 u_{\text{hom}}$ for some $p > 2 \geq d$ and therefore a uniform bound on ∇u_{hom} of the form (63) by Morrey's embedding.

Corollary 15 yields the estimate

$$(64) \quad \left(\int_{B_{5d\delta}(x_0)} |\phi_{\xi_k} - \phi_{\xi_l}|^2 + |\sigma_{\xi_k} - \sigma_{\xi_l}|^2 dx \right)^{1/2} \\ \leq \begin{cases} \mathcal{C}(1 + \sup_{\mathbb{R}^d} |\nabla u_{\text{hom}}|)^{\mathcal{C}} |\xi_k - \xi_l| \varepsilon^{1/2} & \text{for } d = 1, \\ \mathcal{C}(1 + \sup_{\mathbb{R}^d} |\nabla u_{\text{hom}}|)^{\mathcal{C}} |\xi_k - \xi_l| \varepsilon \left| \log \frac{\delta + |x_0|}{\varepsilon} \right|^{1/2} & \text{for } d = 2, \\ \mathcal{C}(1 + \sup_{\mathbb{R}^d} |\nabla u_{\text{hom}}|)^{\mathcal{C}} |\xi_k - \xi_l| \varepsilon & \text{for } d \geq 3 \end{cases}$$

for a random constant \mathcal{C} with uniformly bounded stretched exponential moments in the sense $\mathbb{E}[\exp(\mathcal{C}^{1/C(d,m,\lambda,\Lambda,\rho,\nu)})] \leq 2$. Using Proposition 27 with $\delta := \varepsilon$ in case $d \geq 3$ and $\delta := \varepsilon |\log \varepsilon|^{1/2}$ in case $d = 2$, we deduce

$$(65) \quad \nabla \cdot (A(\omega_\varepsilon, \nabla u_\varepsilon) - A(\omega_\varepsilon, \nabla \hat{u})) = \nabla \cdot R$$

for $d \geq 3$ respectively

$$(66) \quad \nabla \cdot (A(\omega_\varepsilon, \nabla u_\varepsilon) - A(\omega_\varepsilon, \nabla \hat{u})) + (u_\varepsilon - \hat{u}) = \nabla \cdot R - \sum_{k \in K} \eta_k \phi_{\xi_k}$$

for $d = 2$ or $d = 1$; here, R is estimated by the estimate in Proposition 27 as well as (64), (58), (59b), and (63) by (using also the abbreviation $\mu(s) := 1$ for $d \geq 3$ and $\mu(s) := |\log(s+2)|^{1/2}$ for $d = 2$)

$$(67) \quad \int_{\mathbb{R}^d} |R|^2 dx \leq C\delta^2 \int_{\mathbb{R}^d} |\nabla^2 \bar{u}|^2 dx + \frac{C}{\delta^2} \sum_{k \in \delta\mathbb{Z}^d} \mathcal{C}_k \varepsilon^2 \mu_\varepsilon^2 \delta^2 \int_{B_{d\delta}(k)} |\nabla^2 \bar{u}|^2 dx \\ \leq \begin{cases} \mathcal{C}(f)\varepsilon^2 & \text{in case } d \geq 3, \\ \mathcal{C}(f)\varepsilon^2 |\log \varepsilon| & \text{in case } d = 2 \end{cases}$$

for random constants \mathcal{C}_k with uniformly bounded stretched exponential moments in the sense $\mathbb{E}[\exp(\mathcal{C}_k^{1/C(d,m,\lambda,\Lambda,\rho,\nu)})] \leq 2$ and a random constant \mathcal{C} with stretched exponential moments $\mathbb{E}[\exp(\mathcal{C}^{1/C(d,m,\lambda,\Lambda,\rho,\nu)}/C(f))] \leq 2$.

An energy estimate for the PDEs (65) and (66) followed by the Sobolev embedding now yields in case $d \geq 3$

$$\left\| u_\varepsilon - u_{\text{hom}} - \sum_{k \in K} \eta_k \phi_{\xi_k} \right\|_{L^{2d/(d-2)}(\mathbb{R}^d)} \leq C \|\nabla u_\varepsilon - \nabla \hat{u}\|_{L^2(\mathbb{R}^d)} \leq \mathcal{C}\varepsilon,$$

while for $d = 2$ we directly obtain from the energy estimate

$$\left\| u_\varepsilon - u_{\text{hom}} - \sum_{k \in K} \eta_k \phi_{\xi_k} \right\|_{L^2(\mathbb{R}^d)} \leq \mathcal{C}\varepsilon.$$

Using again the estimate on ϕ_ξ from Corollary 15, the bound for the ξ_k from (46), and the energy estimate (59a), we obtain with $p := \frac{2d}{d-2}$ in case $d \geq 3$ and $p := 2$ for $d \leq 2$

$$\begin{aligned} \int_{\mathbb{R}^d} \left| \sum_{k \in K} \eta_k \phi_{\xi_k} \right|^p dx &\leq \sum_k \int_{k+(-\delta, \delta)^d} \mathcal{C}_k \varepsilon^p \mu_\varepsilon^p |\xi_k|^p dx \\ &\leq \mathcal{C} \varepsilon^p \mu_\varepsilon^p \int_{\mathbb{R}^d} |\nabla u_{\text{hom}}|^p dx \stackrel{(58), (59b)}{\leq} \mathcal{C} \varepsilon^p \mu_\varepsilon^p. \end{aligned}$$

Combining the previous three estimates, we obtain the statements of Theorem 2 and Theorem 3. \square

Proof of Lemma 26. By (12a), the difference $\phi_\xi^{2T} - \phi_\xi^T$ satisfies the equation

$$-\nabla \cdot (\hat{a} \nabla (\phi_\xi^{2T} - \phi_\xi^T)) + \frac{1}{2T} (\phi_\xi^{2T} - \phi_\xi^T) = \frac{1}{2T} \phi_\xi^T$$

with the $-$ by (A1) and (A2) uniformly elliptic and bounded $-$ coefficient field \hat{a} defined by

$$\hat{a}(x) := \int_0^1 \partial_\xi A(\omega_\varepsilon(x), (1-s)\nabla \phi_\xi^T(x) + s\nabla \phi_\xi^{2T}(x)) ds.$$

By Lemma 36, we deduce

$$\int_{\mathbb{R}^d} |\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T|^2 \exp\left(-\frac{\gamma|x|}{\sqrt{T}}\right) dx \leq C \int_{\mathbb{R}^d} \frac{1}{T} |\phi_\xi^{2T}|^2 \exp\left(-\frac{\gamma|x|}{\sqrt{T}}\right) dx.$$

Taking the expected value and plugging in the estimates (21) and (22a), we deduce by stationarity

$$(68) \quad \mathbb{E}[|\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T|^2]^{1/2} \leq \begin{cases} C|\xi|\varepsilon^{1/2}\sqrt{T}^{-1/2} & \text{for } d = 1, \\ C|\xi|\varepsilon\sqrt{T}^{-1}|\log \sqrt{T}/\varepsilon|^{1/2} & \text{for } d = 2, \\ C|\xi|\varepsilon\sqrt{T}^{-1} & \text{for } d \geq 3. \end{cases}$$

Thus, $\nabla \phi_\xi^T$ converges in $L^2(\Omega \times B_r)$ to a stationary gradient field Θ with finite second moments and zero expectation. It may thus be represented in the form $\Theta = \nabla \varphi$, where φ is potentially non-stationary, but satisfies $\int_{B_\varepsilon(0)} \varphi dx = 0$. By passing to the limit in the localized corrector equation (12a), we see that φ is a solution to (4) in the sense of Definition 1. By uniqueness, we conclude that $\varphi = \phi_\xi$.

We next treat σ_ξ . Taking the difference of the PDE (12c) for T and $2T$, we obtain

$$\begin{aligned} -\Delta(\sigma_{\xi, jk}^{2T} - \sigma_{\xi, jk}^T) + \frac{1}{2T}(\sigma_{\xi, jk}^{2T} - \sigma_{\xi, jk}^T) \\ = \frac{1}{2T} \sigma_{\xi, jk}^T + \nabla \times (A(\omega_\varepsilon, \xi + \nabla \phi_\xi^{2T}) - A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T)) \end{aligned}$$

where by slight abuse of notation we have written $\nabla \times G$ to abbreviate $\partial_j(G \cdot e_k) - \partial_k(G \cdot e_j)$. Testing this equation with $\sigma_{\xi, jk} \exp(-\gamma|x|/\sqrt{T})$, taking the expectation,

and using stationarity as well as (A2) and (68), we obtain

$$\mathbb{E}[|\nabla\sigma_{\xi,jk}^{2T} - \nabla\sigma_{\xi,jk}^T|^2]^{1/2} \leq \begin{cases} C|\xi|\varepsilon^{1/2}\sqrt{T}^{-1/2} & \text{for } d = 1, \\ C|\xi|\varepsilon\sqrt{T}^{-1}|\log\sqrt{T}/\varepsilon|^{1/2} & \text{for } d = 2, \\ C|\xi|\varepsilon\sqrt{T}^{-1} & \text{for } d \geq 3. \end{cases}$$

The conclusion now follows by an argument analogous to the argument for ϕ_ξ^T . We finally note that for $d \geq 3$ the argument extends to ϕ_ξ^T and σ_ξ^T itself. \square

4.3. Estimates on the random fluctuations of the RVE approximation for the effective material law. We next establish the estimates on fluctuations of the representative volume approximation for the effective material law stated in Theorem 9a.

Proof of Theorem 9a. Consider the random variable

$$F(\omega_{\varepsilon,L}) := A^{\text{RVE},L}(\omega_{\varepsilon,L}, \xi) \cdot \Xi = \int_{[0,L]^d} A(\omega_{\varepsilon,L}(x), \xi + \nabla\phi_\xi(\omega_{\varepsilon,L}, x)) \cdot \Xi \, dx.$$

Let $\delta\omega_{\varepsilon,L}$ denote a periodic infinitesimal perturbation in the sense of Definition 11b. Then for all L -periodic parameter fields $\omega_{\varepsilon,L}$ we have

$$(69) \quad \begin{aligned} \delta F(\omega_{\varepsilon,L}) &:= \lim_{t \rightarrow 0} \frac{F(\omega + t\delta\omega_L) - F(\omega)}{t} \\ &= \int_{[0,L]^d} \partial_\omega A(\omega_{\varepsilon,L}(x), \xi + \nabla\phi_\xi(\omega_{\varepsilon,L}, x)) \delta\omega_{\varepsilon,L}(x) \cdot \Xi \\ &\quad + a_\xi(x) \nabla\delta\phi_\xi(\omega_{\varepsilon,L}, x) \cdot \Xi \, dx, \end{aligned}$$

where

$$a_\xi(x) = \partial_\xi A(\omega_{\varepsilon,L}(x), \xi + \nabla\phi_\xi(\omega, x)),$$

and where $\delta\phi_\xi = \delta\phi_\xi(\omega_{\varepsilon,L}, \cdot)$ is the unique (L -periodic) solution with mean zero to

$$(70) \quad -\nabla \cdot (a_\xi \nabla\delta\phi_\xi) = \nabla \cdot (\partial_\omega A(\omega_{\varepsilon,L}(x), \xi + \nabla\phi_\xi(\omega_{\varepsilon,L}, x)) \delta\omega_{\varepsilon,L}(x)).$$

Introducing the unique L -periodic solution h with vanishing mean to the PDE

$$(71) \quad -\nabla \cdot (a_\xi^* \nabla h) = \nabla \cdot (a_\xi^*(x) \Xi),$$

we deduce by testing (71) with ϕ_ξ and testing (70) with h

$$\begin{aligned} \delta F(\omega_{\varepsilon,L}) &= \int_{[0,L]^d} \partial_\omega A(\omega_{\varepsilon,L}(x), \xi + \nabla\phi_\xi(\omega, x)) \delta\omega_{\varepsilon,L}(x) \cdot \Xi \, dx \\ &\quad + \int_{[0,L]^d} \partial_\omega A(\omega_{\varepsilon,L}(x), \xi + \nabla\phi_\xi(\omega_{\varepsilon,L}, x)) \delta\omega_{\varepsilon,L}(x) \cdot \nabla h \, dx. \end{aligned}$$

This establishes by (A3)

$$\left| \frac{\partial F(\omega_{\varepsilon,L})}{\partial \omega_{\varepsilon,L}} \right| \leq CL^{-d} |\xi + \nabla\phi_\xi| (|\Xi| + |\nabla h|)$$

which yields by the q -th moment version of the spectral gap inequality in Lemma 16

$$\mathbb{E}_L \left[|F - \mathbb{E}_L[F]|^{2q} \right]^{1/2q} \leq CqL^{-d} \mathbb{E}_L \left[\left(\int_{[0,L]^d} \int_{B_\varepsilon(x)} |\xi + \nabla\phi_\xi| (|\Xi| + |\nabla h|) \, d\tilde{x} \right)^q \right]^{1/2q}.$$

By Hölder's inequality, we infer for any $p > 2$

$$\begin{aligned} & \mathbb{E}_L \left[|F - \mathbb{E}_L[F]|^{2q} \right]^{1/2q} \\ & \leq CqL^{-d/2} \mathbb{E}_L \left[\left(\int_{[0,L]^d} \left| \int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi|^2 d\tilde{x} \right|^{p/(p-2)} dx \right)^{q(p-2)/p} \right. \\ & \quad \left. \times \left(\int_{[0,L]^d} (|\Xi| + |\nabla h|)^p dx \right)^{2q/p} \right]^{1/2q}. \end{aligned}$$

Bounding the last integral by $C|\Xi|$ by Meyers' estimate for (71) (see Lemma 35) and using the estimate (88), we obtain

$$\begin{aligned} & \mathbb{E}_L \left[|F - \mathbb{E}_L[F]|^{2q} \right]^{1/2q} \\ & \leq CqL^{-d/2} |\xi| |\Xi| \mathbb{E}_L \left[\left(\int_{[0,L]^d} |r_{*,L,\xi}(x)|^{(d-\delta)p/(p-2)} dx \right)^{q(p-2)/p} \right]^{1/2q}. \end{aligned}$$

Using stationarity of $r_{*,L,\xi}$ and the moment bound of Lemma 19 (which we prove explicitly for the probability distribution \mathbb{P} , but which may be established for \mathbb{P}_L analogously; furthermore, while the estimates of Lemma 19 are stated for finite $T < \infty$, they are uniform in $T \geq \varepsilon^2$ and therefore also hold in the limit $T \rightarrow \infty$), we deduce for q large enough

$$\mathbb{E}_L \left[|F - \mathbb{E}_L[F]|^{2q} \right]^{1/2q} \leq Cq^C L^{-d/2} |\xi| |\Xi|.$$

This is the assertion of Theorem 9a. \square

4.4. Estimates for the error introduced by localization. We next establish the estimates from Lemma 31, Corollary 32, and Lemma 33 for the error introduced in the correctors by the exponential localization on scale \sqrt{T} via the massive term. We then prove Proposition 30, which estimates the systematic error in the approximation for the effective coefficient $\mathbb{E}[A^{\text{RVE},\eta,T}]$ introduced by the finite localization parameter $T < \infty$.

Proof of Lemma 31. We will use the exponential weight $\eta(x) := \exp(-\gamma|x|/\sqrt{T})$ with $0 < \gamma \ll 1$. Note that

$$(72) \quad |\nabla \eta| \leq \frac{\gamma}{\sqrt{T}} \eta.$$

By the localized corrector equation (12a) we have

$$-\nabla \cdot (A(\omega_\varepsilon, \xi + \nabla \phi_\xi^{2T}) - A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T)) + \frac{1}{2T} (\phi_\xi^{2T} - \phi_\xi^T) = \frac{1}{2T} \phi_\xi^T.$$

Testing with $(\phi_\xi^{2T} - \phi_\xi^T)\eta$ and using the monotonicity and Lipschitz continuity of A (see (A1)-(A2)) as well as (72), we get

$$\begin{aligned} & \int_{\mathbb{R}^d} \left(\lambda |\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T|^2 + \frac{1}{2T} |\phi_\xi^{2T} - \phi_\xi^T|^2 \right) \eta dx \\ & \leq \frac{1}{2T} \int_{\mathbb{R}^d} \phi_\xi^T (\phi_\xi^{2T} - \phi_\xi^T) \eta dx + \gamma \int_{\mathbb{R}^d} \Lambda |\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T| \frac{1}{\sqrt{T}} |\phi_\xi^{2T} - \phi_\xi^T| \eta dx. \end{aligned}$$

Choosing $\gamma \leq c(d, m, \lambda, \Lambda)$, we may absorb the second term on the RHS into the LHS to obtain

$$(73) \quad \int_{\mathbb{R}^d} \left(|\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T|^2 + \frac{1}{T} |\phi_\xi^{2T} - \phi_\xi^T|^2 \right) \eta \, dx \leq \frac{C}{T} \int_{\mathbb{R}^d} \phi_\xi^T (\phi_\xi^{2T} - \phi_\xi^T) \eta \, dx.$$

In the following, we treat dimensions $d \leq 2$ and $d \geq 3$ separately. In the case $d \leq 2$, by Young's inequality and absorption the previous inequality yields

$$\int_{\mathbb{R}^2} \left(|\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T|^2 + \frac{1}{T} |\phi_\xi^{2T} - \phi_\xi^T|^2 \right) \eta \, dx \leq \frac{C}{T} \int_{\mathbb{R}^2} |\phi_\xi^T|^2 \eta \, dx.$$

Note that Proposition 13 (in connection with a dyadic decomposition of \mathbb{R}^2 into the ball $B_{\sqrt{T}}(0)$ and the annuli $\{2^i \sqrt{T} \leq |x| < 2^{i+1} \sqrt{T}\}$, $i = 0, 1, 2, \dots$) we obtain

$$\frac{1}{T} \int_{\mathbb{R}^2} |\phi_\xi^T|^2 \eta \, dx \leq \begin{cases} C |\xi|^2 \varepsilon \sqrt{T}^{-1} & \text{for } d = 1, \\ C |\xi|^2 \varepsilon^2 \sqrt{T}^{-2} \left| \log \frac{\sqrt{T}}{\varepsilon} \right| & \text{for } d = 2. \end{cases}$$

Since $\eta \geq \exp(-1)$ on $B_{\sqrt{T}}$, the claimed estimate follows for $d \leq 2$.

In the case $d \geq 3$ we can use in (73) the representation $\phi_\xi^T = \nabla \cdot (\theta_\xi^T - b)$ for any $b \in \mathbb{R}^{m \times d}$ (see (19b)). We obtain by an integration by parts and by the Cauchy-Schwarz inequality the estimate

$$\begin{aligned} & \frac{1}{T} \int_{\mathbb{R}^d} \phi_\xi^T (\phi_\xi^{2T} - \phi_\xi^T) \eta \, dx \\ & \leq \frac{1}{T} \int_{\mathbb{R}^d} |\theta_\xi^T - b| \left(|\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T| + \frac{\gamma}{\sqrt{T}} |\phi_\xi^{2T} - \phi_\xi^T| \right) \eta \, dx \end{aligned}$$

With Young's inequality we may absorb the second factor into the LHS of (73). We thus obtain

$$\int_{\mathbb{R}^d} \left(|\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T|^2 + \frac{1}{T} |\phi_\xi^{2T} - \phi_\xi^T|^2 \right) \eta \, dx \leq \frac{C}{T^2} \int_{\mathbb{R}^d} |\theta_\xi^T - b|^2 \eta \, dx.$$

By appealing to Proposition 13 (in connection with a dyadic decomposition of \mathbb{R}^d), and the fact that $\eta \geq \exp(-1)$ on $B_{\sqrt{T}}$, the claimed estimate follows for $d \geq 3$. \square

Proof of Corollary 32. Set $u := \phi_\xi^{2T} - \phi_\xi^T$ and note that with $\hat{a}(\hat{\xi}) := \int_0^1 \partial_\xi A(\omega_\varepsilon, \xi + (1-s)\nabla \phi_\xi^T + s\nabla \phi_\xi^{2T}) \, ds$ we have by (12a)

$$-\nabla \cdot (\hat{a} \nabla u) + \frac{1}{T} u = \frac{1}{2T} \phi_{2T}.$$

Applying Lemma 44 to this PDE – upon rewriting the right-hand side using (19b) in case $d \geq 3$ – we obtain

$$\begin{aligned} & \left(\int_{B_{\sqrt{T}}(x_0)} |\nabla(\phi_\xi^{2T} - \phi_\xi^T)|^p \, dx \right)^{1/p} \\ & \leq C(d, m, \lambda, \Lambda, p) \left(\int_{B_{2\sqrt{T}}(x_0)} |\nabla u|^2 + \left| \frac{1}{\sqrt{T}} u \right|^2 \, dx \right)^{1/2} \\ & \quad + \begin{cases} C(d, m, \lambda, \Lambda, p) \left(\int_{B_{2\sqrt{T}}(x_0)} \left| \frac{1}{\sqrt{T}} \phi_\xi^{2T} \right|^p \, dx \right)^{1/p} & \text{in case } d \leq 2, \\ C(d, m, \lambda, \Lambda, p) \left(\int_{B_{2\sqrt{T}}(x_0)} \left| \frac{1}{T} (\phi_\xi^{2T} - b) \right|^p \, dx \right)^{1/p} & \text{in case } d \geq 3. \end{cases} \end{aligned}$$

This implies by Proposition 13 and Lemma 31

$$\begin{aligned} & \left(\int_{B_{\sqrt{T}}(x_0)} |\nabla(\phi_\xi^{2T} - \phi_\xi^T)|^p dx \right)^{1/p} \\ & \leq C|\xi| \left(\frac{\varepsilon}{\sqrt{T}} \right)^{\frac{d \wedge 4}{2}} \begin{cases} |\log(\sqrt{T}/\varepsilon)| & \text{for } d \in \{2, 4\}, \\ 1 & \text{for } d = 3, d = 1, \text{ and } d \geq 5. \end{cases} \end{aligned}$$

Taking the p -th stochastic moment and using stationarity, we conclude. \square

Proof of Lemma 33. Again, we use the weight $\eta(x) := \exp(-\gamma|x|/\sqrt{T})$ for $0 < \gamma \ll 1$. We subtract the equations for $\phi_{\xi, \Xi}^{2T}$ and $\phi_{\xi, \Xi}^T$ (see (18a)) to obtain

$$-\nabla \cdot \left(a_\xi^{2T}(\Xi + \nabla \phi_{\xi, \Xi}^{2T}) - a_\xi^T(\Xi + \nabla \phi_{\xi, \Xi}^T) \right) + \frac{1}{T}(\phi_{\xi, \Xi}^{2T} - \phi_{\xi, \Xi}^T) = \frac{1}{2T}\phi_{\xi, \Xi}^{2T}.$$

By adding and subtracting $a_\xi^T(\Xi + \phi_{\xi, \Xi}^{2T})$ and by additionally appealing to the representation $\nabla \cdot (\theta_{\xi, \Xi}^{2T} - b) = \phi_{\xi, \Xi}^{2T}$ for any $b \in \mathbb{R}^{m \times d}$ (see (20b)) in the case $d \geq 3$, we get

$$\begin{aligned} & -\nabla \cdot \left(a_\xi^T(\nabla \phi_{\xi, \Xi}^{2T} - \nabla \phi_{\xi, \Xi}^T) \right) + \frac{1}{T}(\phi_{\xi, \Xi}^{2T} - \phi_{\xi, \Xi}^T) \\ (74) \quad & = \nabla \cdot \left((a_\xi^{2T} - a_\xi^T)(\Xi + \nabla \phi_{\xi, \Xi}^{2T}) \right) + \begin{cases} \frac{1}{2T}\phi_{\xi, \Xi}^{2T} & \text{for } d \leq 2, \\ \nabla \cdot \left(\frac{1}{2T}\theta_{\xi, \Xi}^{2T} - b \right) & \text{for } d \geq 3. \end{cases} \end{aligned}$$

Testing the equation with $(\phi_{\xi, \Xi}^{2T} - \phi_{\xi, \Xi}^T)\eta$ (with $0 < \gamma \ll 1$) yields the exponentially localized energy estimate

$$\begin{aligned} & \int_{\mathbb{R}^d} \left(\lambda |\nabla \phi_{\xi, \Xi}^{2T} - \nabla \phi_{\xi, \Xi}^T|^2 + \frac{1}{T} |\phi_{\xi, \Xi}^{2T} - \phi_{\xi, \Xi}^T|^2 \right) \eta dx \\ & \leq C \int_{\mathbb{R}^d} |a_\xi^{2T} - a_\xi^T|^2 |\Xi + \nabla \phi_{\xi, \Xi}^{2T}|^2 \eta dx + C \begin{cases} \frac{1}{T} \int_{\mathbb{R}^d} |\phi_{\xi, \Xi}^{2T}|^2 \eta dx & \text{for } d \leq 2, \\ \frac{1}{T^2} \int_{\mathbb{R}^d} |\theta_{\xi, \Xi}^{2T} - b|^2 \eta dx & \text{for } d \geq 3. \end{cases} \end{aligned}$$

By taking the expectation and exploiting stationarity of the LHS, we get

$$\begin{aligned} & \mathbb{E} \left[|\nabla \phi_{\xi, \Xi}^{2T} - \nabla \phi_{\xi, \Xi}^T|^2 + \frac{1}{T} |\phi_{\xi, \Xi}^{2T} - \phi_{\xi, \Xi}^T|^2 \right] \\ & \leq C \mathbb{E} \left[|a_\xi^{2T} - a_\xi^T|^2 |\Xi + \nabla \phi_{\xi, \Xi}^{2T}|^2 \right] + C \begin{cases} \frac{1}{T} \mathbb{E} [|\phi_{\xi, \Xi}^{2T}|^2] & \text{for } d \leq 2, \\ \frac{1}{T^2} \mathbb{E} \left[\int_{\mathbb{R}^d} |\theta_{\xi, \Xi}^{2T} - b|^2 \frac{\eta}{\sqrt{T}^d} dx \right] & \text{for } d \geq 3. \end{cases} \end{aligned}$$

The second term on the RHS can be estimated with help of Proposition 14.

We estimate the first term on the RHS: Since $\partial_\xi A$ is Lipschitz by assumption (R) and since $a_\xi^T(x) = A(\omega_\varepsilon(x), \xi + \nabla \phi_\xi^T)$, we have $|a_\xi^{2T} - a_\xi^T| \leq C |\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T|$. Hence, with Hölder's inequality with exponents p and $\frac{p}{p-1}$, the bound on $\Xi + \nabla \phi_{\xi, \Xi}^T$

from (42), and Corollary 32, we get

$$\begin{aligned}
& \mathbb{E} [|a_{\xi}^{2T} - a_{\xi}^T|^2 |\Xi + \nabla \phi_{\xi, \Xi}^{2T}|^2] \\
& \leq C \mathbb{E} [|\nabla \phi_{\xi}^{2T} - \nabla \phi_{\xi}^T|^{2p}]^{\frac{1}{p}} \mathbb{E} [|\Xi + \nabla \phi_{\xi, \Xi}^{2T}|^{2\frac{p-1}{p}}]^{\frac{p-1}{p}} \\
& \leq C |\Xi|^2 (1 + |\xi|)^{2C} |\xi|^2 \left(\frac{\varepsilon}{\sqrt{T}} \right)^{d \wedge 4} \times \begin{cases} |\log(\sqrt{T}/\varepsilon)|^2 & \text{for } d \in \{2, 4\}, \\ 1 & \text{for } d = 3, d = 1, \text{ and } d \geq 5. \end{cases}
\end{aligned}$$

□

Proof of Proposition 30. Step 1: Proof of (a). This is a direct consequence of stationarity of \mathbb{P} (see assumption (P1)) and stationarity of the random field $A(\omega_{\varepsilon}, \xi + \nabla \phi_{\xi}^T) \cdot \Xi - \frac{1}{T} \phi_{\xi}^T \phi_{\xi, \Xi}^{*,T}$, the latter of which is a consequence of the former.

Step 2: Proof of (b). First note that it suffices to prove for any $T \geq \varepsilon^2$ the estimate

$$\begin{aligned}
(75) \quad & | \mathbb{E} [A^{\text{RVE}, \eta, 2T}(\xi) \cdot \Xi] - \mathbb{E} [A^{\text{RVE}, \eta, T}(\xi) \cdot \Xi] | \\
& \leq C (1 + |\xi|)^{2C} |\xi| |\Xi| \left(\frac{\varepsilon}{\sqrt{T}} \right)^{(d \wedge 4)} \begin{cases} |\log(\sqrt{T}/\varepsilon)| & \text{for } d = 2 \text{ and } d = 4 \\ 1 & \text{for } d = 3, d = 1, \text{ and } d \geq 5. \end{cases}
\end{aligned}$$

Indeed, the claimed estimate then follows by rewriting the systematic localization error as a telescopic sum,

$$\begin{aligned}
& A_{\text{hom}}(\xi) \cdot \Xi - \mathbb{E} [A^{\text{RVE}, \eta, T}(\xi) \cdot \Xi] \\
& = \sum_{i=0}^{\infty} \left(\mathbb{E} [A^{\text{RVE}, \eta, 2^{i+1}T}(\xi) \cdot \Xi] - \mathbb{E} [A^{\text{RVE}, \eta, 2^i T}(\xi) \cdot \Xi] \right),
\end{aligned}$$

which holds since

$$\lim_{T \rightarrow \infty} \mathbb{E} [A^{\text{RVE}, \eta, T}(\xi) \cdot \Xi] = A_{\text{hom}} \cdot \Xi \quad \mathbb{P}\text{-almost surely.}$$

We present the argument for (75). In view of (a), we may assume without loss of generality that the weight η satisfies

$$\text{supp } \eta \subset B_{\sqrt{T}}, \quad \int_{\mathbb{R}^d} \eta \, dx = 1, \quad |\eta| + \sqrt{T} |\nabla \eta| \lesssim \sqrt{T}^{-d}.$$

Let $\phi_{\xi, \Xi}^{*,T}$ denote the localized, linearized, adjoint corrector (i.e. the T -localized homogenization corrector associated with the linear elliptic PDE with coefficient field $(a_{\xi}^T)^*$). The localized corrector equation (12a) yields

$$\begin{aligned}
(76) \quad & - \int \eta \left(A(\omega_{\varepsilon}, \xi + \nabla \phi_{\xi}^{2T}) - A(\omega_{\varepsilon}, \xi + \nabla \phi_{\xi}^T) \right) \cdot \nabla \phi_{\xi, \Xi}^{*,T} \, dx \\
& = \int \left(A(\omega_{\varepsilon}, \xi + \nabla \phi_{\xi}^{2T}) - A(\omega_{\varepsilon}, \xi + \nabla \phi_{\xi}^T) \right) \cdot (\phi_{\xi, \Xi}^{*,T} \nabla \eta) + \eta \left(\frac{1}{2T} \phi_{\xi}^{2T} - \frac{1}{T} \phi_{\xi}^T \right) \phi_{\xi, \Xi}^{*,T} \, dx.
\end{aligned}$$

Combined with the definition of the localized RVE approximation in Definition 28, we get

$$\begin{aligned}
& \left(A^{\text{RVE},\eta,2T}(\xi) - A^{\text{RVE},\eta,T}(\xi) \right) \cdot \Xi \\
&= \int \eta \left(\left(A(\omega_\varepsilon, \xi + \nabla \phi_\xi^{2T}) - A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \right) \cdot \Xi - \frac{1}{2T} \phi_\xi^{2T} \phi_{\xi,\Xi}^{*,2T} + \frac{1}{T} \phi_\xi^T \phi_{\xi,\Xi}^{*,T} \right) dx \\
&= \int \eta \left(A(\omega_\varepsilon, \xi + \nabla \phi_\xi^{2T}) - A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \right) \cdot (\Xi + \nabla \phi_{\xi,\Xi}^{*,T}) - \frac{1}{2T} \phi_\xi^{2T} \phi_{\xi,\Xi}^{*,2T} + \frac{1}{T} \phi_\xi^T \phi_{\xi,\Xi}^{*,T} \Big) dx \\
&\quad + \int \left(A(\omega_\varepsilon, \xi + \nabla \phi_\xi^{2T}) - A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \right) \cdot (\phi_{\xi,\Xi}^{*,T} \nabla \eta) + \eta \left(\frac{1}{2T} \phi_\xi^{2T} - \frac{1}{T} \phi_\xi^T \right) \phi_{\xi,\Xi}^{*,T} dx \\
&= \int \eta \left(A(\omega_\varepsilon, \xi + \nabla \phi_\xi^{2T}) - A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \right) \cdot (\Xi + \nabla \phi_{\xi,\Xi}^{*,T}) - \frac{1}{2T} \phi_\xi^{2T} (\phi_{\xi,\Xi}^{*,2T} - \phi_{\xi,\Xi}^{*,T}) \Big) dx \\
&\quad + \int \left(A(\omega_\varepsilon, \xi + \nabla \phi_\xi^{2T}) - A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \right) \cdot (\phi_{\xi,\Xi}^{*,T} \nabla \eta) dx.
\end{aligned}$$

We subtract the linearized corrector equation (18a) in its form for the adjoint coefficient field and the corresponding corrector

$$\int a_\xi^{*,T} (\Xi + \nabla \phi_{\xi,\Xi}^{*,T}) \cdot \nabla (\eta (\phi_\xi^{2T} - \phi_\xi^T)) + \frac{1}{T} \int \eta \phi_{\xi,\Xi}^{*,T} (\phi_\xi^{2T} - \phi_\xi^T) dx = 0,$$

where $a_\xi^T := \partial_\xi A(\xi + \nabla \phi_\xi^T)$. We get

$$\begin{aligned}
& (77) \\
& \left(A^{\text{RVE},\eta,2T}(\xi) - A^{\text{RVE},\eta,T}(\xi) \right) \cdot \Xi \\
&= \int \eta \left(A(\omega_\varepsilon, \xi + \nabla \phi_\xi^{2T}) - A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) - a_\xi^T (\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T) \right) \cdot (\Xi + \nabla \phi_{\xi,\Xi}^{*,T}) dx \\
&\quad + \int \left(A(\omega_\varepsilon, \xi + \nabla \phi_\xi^{2T}) - A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \right) \cdot \phi_{\xi,\Xi}^{*,T} \nabla \eta - a_\xi^T (\Xi + \nabla \phi_{\xi,\Xi}^{*,T}) \\
&\quad \quad \cdot (\phi_\xi^{2T} - \phi_\xi^T) \nabla \eta dx \\
&\quad - \int \eta \left(\frac{1}{2T} \phi_\xi^{2T} (\phi_{\xi,\Xi}^{*,2T} - \phi_{\xi,\Xi}^{*,T}) - \frac{1}{T} \phi_{\xi,\Xi}^{*,T} (\phi_\xi^{2T} - \phi_\xi^T) \right) dx.
\end{aligned}$$

We take the expectation of this identity and note that the expectation of the second integral on the right-hand side vanishes: Indeed, since it is of the form $\mathbb{E}[\int_{\mathbb{R}^d} B \nabla \eta]$ where B is a stationary random field and η is compactly supported, we have $\mathbb{E}[\int_{\mathbb{R}^d} B \nabla \eta] = \mathbb{E}[B] \int_{\mathbb{R}^d} \nabla \eta = 0$. Moreover, for the first term on the RHS of (77) we appeal to the uniform bound on $\partial_\xi^2 A$ from assumption (R) in form of (recall that $a_\xi^T = \partial_\xi A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T)$)

$$|A(\omega_\varepsilon, \xi + \nabla \phi_\xi^{2T}) - A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) - a_\xi^T (\xi + \nabla \phi_\xi^T)| \leq C |\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T|^2.$$

We thus get

$$\begin{aligned}
(78) \quad & \left| \mathbb{E} \left[\left(A^{\text{RVE}, \eta, 2T}(\xi) - A^{\text{RVE}, \eta, T}(\xi) \right) \cdot \Xi \right] \right| \\
& \leq C \mathbb{E} \left[\left| \nabla \phi_\xi^{2T} - \nabla \phi_\xi^T \right|^2 |\Xi| + \nabla \phi_{\xi, \Xi}^{*, T} \right] \\
& \quad + \left| \mathbb{E} \left[\int \eta \left(\frac{1}{2T} \phi_\xi^{2T} (\phi_{\xi, \Xi}^{*, 2T} - \phi_{\xi, \Xi}^{*, T}) - \frac{1}{T} \phi_{\xi, \Xi}^{*, T} (\phi_\xi^{2T} - \phi_\xi^T) \right) dx \right] \right| \\
& =: I_1 + I_2.
\end{aligned}$$

To estimate I_1 we first apply Hölder's inequality with exponents p and $\frac{p}{p-1}$ with $0 < p-1 \ll 1$, and then appeal to Corollary 32 and the moment bound on the linearized corrector from (42) as well as (23) to deduce

$$I_1 \leq C |\xi|^2 (1 + |\xi|)^C |\Xi| \left(\frac{\varepsilon}{\sqrt{T}} \right)^{d \wedge 4} \begin{cases} |\log(\sqrt{T}/\varepsilon)|^2 & \text{for } d \in \{2, 4\}, \\ 1 & \text{for } d = 3, d = 1, \text{ and } d \geq 5. \end{cases}$$

Regarding I_2 we distinguish the cases $d \leq 2$ and $d \geq 3$. In the case $d \leq 2$, we apply Lemma 33, Lemma 31, as well as Proposition 13 and Proposition 14 to obtain

$$\begin{aligned}
I_2 & \leq C \mathbb{E} \left[\frac{1}{T} |\phi_\xi^{2T}|^2 \right]^{\frac{1}{2}} \mathbb{E} \left[\frac{1}{T} |\phi_{\xi, \Xi}^{*, 2T} - \phi_{\xi, \Xi}^{*, T}|^2 \right]^{\frac{1}{2}} + \mathbb{E} \left[\frac{1}{T} |\phi_{\xi, \Xi}^{*, T}|^2 \right]^{\frac{1}{2}} \mathbb{E} \left[\frac{1}{T} |\phi_\xi^{2T} - \phi_\xi^T|^2 \right]^{\frac{1}{2}} \\
& \leq C (1 + |\xi|)^C |\xi| |\Xi| \varepsilon^2 \sqrt{T}^{-2} |\log(\sqrt{T}/\varepsilon)|^2,
\end{aligned}$$

and in case $d = 1$ we proceed similarly.

In the case $d \geq 3$ we appeal to the representation of ϕ_ξ^{2T} and $\phi_{\xi, \Xi}^{*, T}$ as $\nabla \cdot \theta_\xi^{2T}$ and $\nabla \cdot \theta_{\xi, \Xi}^{*, 2T}$ by (19b) and (20b), respectively. To shorten the notation, in the following we assume without loss of generality that $f_{B_{\sqrt{T}}} \theta_\xi^{2T} = f_{B_{\sqrt{T}}} \theta_{\xi, \Xi}^{*, T} = 0$. An integration by parts thus yields

$$\begin{aligned}
I_2 & = \left| \mathbb{E} \left[\int \eta \left(\frac{1}{2T} \theta_\xi^{2T} \cdot (\nabla \phi_{\xi, \Xi}^{*, 2T} - \nabla \phi_{\xi, \Xi}^{*, T}) - \frac{1}{T} \theta_{\xi, \Xi}^{*, T} \cdot (\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T) \right) dx \right. \right. \\
& \quad \left. \left. + \int \nabla \eta \cdot \left(\frac{1}{2T} \theta_\xi^{2T} (\phi_{\xi, \Xi}^{*, 2T} - \phi_{\xi, \Xi}^{*, T}) - \frac{1}{T} \theta_{\xi, \Xi}^{*, T} (\phi_\xi^{2T} - \phi_\xi^T) \right) dx \right] \right|.
\end{aligned}$$

With the properties of η (in particular, $|\nabla \eta| \lesssim \sqrt{T}^{-d-1}$ and $\text{supp } \eta \subset B_{\sqrt{T}}$), by the Cauchy-Schwarz inequality, and by stationarity of the localized correctors, we get

$$\begin{aligned}
I_2 & \leq \frac{C}{T} \mathbb{E} \left[\int_{B_{\sqrt{T}}} |\theta_\xi^{2T}|^2 dx \right]^{\frac{1}{2}} \mathbb{E} \left[|\nabla \phi_{\xi, \Xi}^{*, 2T} - \nabla \phi_{\xi, \Xi}^{*, T}|^2 + \frac{1}{T} |\phi_{\xi, \Xi}^{*, 2T} - \phi_{\xi, \Xi}^{*, T}|^2 \right]^{\frac{1}{2}} \\
& \quad + \frac{C}{T} \mathbb{E} \left[\int_{B_{\sqrt{T}}} |\theta_{\xi, \Xi}^{*, 2T}|^2 dx \right]^{\frac{1}{2}} \mathbb{E} \left[|\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T|^2 + \frac{1}{T} |\phi_\xi^{2T} - \phi_\xi^T|^2 \right]^{\frac{1}{2}}.
\end{aligned}$$

By appealing to Proposition 13 and Lemma 33 for the first term and to Proposition 14 and Lemma 31 for the second term, we obtain

$$I_2 \leq C(1 + |\xi|)^C |\xi| |\Xi| \left(\frac{\varepsilon}{\sqrt{T}} \right)^{d \wedge 4} \begin{cases} 1 & \text{for } d = 3 \text{ and } d \geq 5, \\ |\log(\sqrt{T}/\varepsilon)|^2 & \text{for } d = 4. \end{cases}$$

Plugging in the estimates on I_1 and I_2 into (78), this establishes the estimate on the localization error for the representative volume element method for \mathbb{P} .

Step 3: Proof of (c). For the periodized probability distribution \mathbb{P}_L , one may proceed analogously to (b), deriving an error estimate on $\mathbb{E}_L[A^{\text{RVE}, \eta, T}] - \mathbb{E}_L[A^{\text{RVE}, L}]$. \square

4.5. Coupling error for RVEs.

Proof of Lemma 29. To shorten the presentation we set $\widehat{\omega} := \pi_L \widetilde{\omega}$ and similarly mark quantities that are associated with $\widehat{\omega}$, i.e.,

$$\widehat{\phi}_\xi^T(x) = \phi_\xi^T(\widehat{\omega}, x), \quad \widehat{\phi}_{\xi, \Xi}^T(x) := \phi_{\xi, \Xi}^T(\widehat{\omega}, x), \quad \widehat{a}_\xi^T(x) := A(\widehat{\omega}, \xi + \nabla \widehat{\phi}_\xi^T).$$

Moreover, we shall use the following notation for the differences

$$\widehat{\delta}\phi_\xi^T := \phi_\xi^T - \widehat{\phi}_\xi^T, \quad \widehat{\delta}\phi_{\xi, \Xi}^T := \phi_{\xi, \Xi}^T - \widehat{\phi}_{\xi, \Xi}^T;$$

(we use the symbol $\widehat{\delta}$ to distinguish the quantity from the sensitivities considered in Sections 5 and 6). In the proof we make use of the exponential test-functions

$$\eta(x) := \exp(-\gamma|x|/\sqrt{T})$$

where $0 < \gamma \ll 1$ is chosen such that the exponential localization estimate Lemma 36 applies.

Step 1. Estimate for $\widehat{\delta}\phi_\xi^T$. We claim that

$$(79) \quad \sup_{x_0 \in B_{L/8}} \int_{\mathbb{R}^d} \left(|\nabla \widehat{\delta}\phi_\xi^T|^2 + \frac{1}{T} |\widehat{\delta}\phi_\xi^T|^2 \right) \eta(x - x_0) dx \leq C\sqrt{T}^d \exp(-\frac{\gamma}{16}L/\sqrt{T}) |\xi|^2.$$

Indeed, by subtracting the equations for ϕ_ξ^T and $\widehat{\phi}_\xi^T$, we find that

$$(80) \quad -\nabla \cdot (a(x) \nabla \widehat{\delta}\phi_\xi^T) + \frac{1}{T} \widehat{\delta}\phi_\xi^T = \nabla \cdot F,$$

where

$$a(x) := \int_0^1 \partial_\xi A(\widetilde{\omega}, \xi + (1-s)\nabla \phi_\xi^T + s\nabla \widehat{\phi}_\xi^T) ds, \\ F := A(\widehat{\omega}, \xi + \nabla \widehat{\phi}_\xi^T) - A(\widetilde{\omega}, \xi + \nabla \phi_\xi^T).$$

By the exponentially localized energy estimate of Lemma 36 we have

$$\int_{\mathbb{R}^d} \left(|\nabla \widehat{\delta}\phi_\xi^T|^2 + \frac{1}{T} |\widehat{\delta}\phi_\xi^T|^2 \right) \eta(x - x_0) dx \leq C \int_{\mathbb{R}^d} |F|^2 \eta(x - x_0) dx.$$

By construction F vanishes on $B_{L/4}$. Hence, since

$$(81) \quad |x| \geq \frac{L}{4}, \quad |x_0| \leq \frac{L}{8} \quad \Rightarrow \quad |x - x_0| \geq \frac{L}{16} + \frac{1}{2}|x - x_0| \\ \Rightarrow \quad \eta(x - x_0) \leq \exp(-\frac{\gamma}{16}L/\sqrt{T}) \exp(-\frac{\gamma}{2}|x - x_0|/\sqrt{T})$$

and since $\hat{\omega} = \tilde{\omega}$ for $|x| \leq \frac{L}{4}$, we obtain

$$\begin{aligned} \int_{\mathbb{R}^d} |F|^2 \eta(x - x_0) dx &\leq C \exp(-\frac{\gamma}{16} L/\sqrt{T}) \int_{\mathbb{R}^d} |\xi + \nabla \hat{\phi}_T|^2 \exp(-\frac{\gamma}{2} |x - x_0|/\sqrt{T}) dx \\ &\leq C \exp(-\frac{\gamma}{16} L/\sqrt{T}) \sqrt{T}^d |\xi|^2, \end{aligned}$$

where for the last estimate we appealed to the localized energy for $\hat{\phi}_\xi^T$, see Lemma 36. We conclude that (79) holds. For further reference, we note that we may similarly derive

$$(82) \quad \sup_{x_0 \in B_{L/8}} \int_{\mathbb{R}^d} \left(|\nabla \hat{\phi}_\xi^T|^p + \left| \frac{1}{\sqrt{T}} \hat{\phi}_\xi^T \right|^p \right) \eta(x - x_0) dx \leq C \sqrt{T}^d \exp(-\frac{\gamma}{16} L/\sqrt{T}) |\xi|^p$$

for some $p = p(d, m, \lambda, \Lambda) > 2$ by applying the Meyers estimate of Lemma 44 to (80) with the dyadic decomposition $\mathbb{R}^d = B_{\sqrt{T}} \cup \bigcup_{k=1}^{\infty} (B_{2^k \sqrt{T}} \setminus B_{2^{k-1} \sqrt{T}})$, the estimate (79), and the bound

$$\begin{aligned} \int_{\mathbb{R}^d} |F|^p \eta(x - x_0) dx &\leq C \exp(-\frac{\gamma}{16} L/\sqrt{T}) \int_{\mathbb{R}^d} |\xi + \nabla \hat{\phi}_T|^p \exp(-\frac{\gamma}{2} |x - x_0|/\sqrt{T}) dx \\ &\leq C \exp(-\frac{\gamma}{16} L/\sqrt{T}) \sqrt{T}^d |\xi|^p. \end{aligned}$$

Note that in the last step of of the last inequality we have again used the Meyers estimate of Lemma 44 together with the localized energy estimate of Lemma 36 and a dyadic decomposition.

Step 2. Estimate for $\hat{\phi}_{\xi, \Xi}^T$. We claim that there exists $q = q(d, m, \lambda, \Lambda)$ such that for all $x_0 \in B_{\frac{L}{8}}$ we have

$$(83) \quad \begin{aligned} &\int_{\mathbb{R}^d} \left(|\nabla \hat{\phi}_{\xi, \Xi}^T|^2 + \frac{1}{T} |\hat{\phi}_{\xi, \Xi}^T|^2 \right) \eta(x - x_0) dx \\ &\leq C \sqrt{T}^d \exp(-\frac{\gamma}{32} L/\sqrt{T}) (1 + |\xi|^2) \|\Xi + \nabla \phi_{\xi, \Xi}^T\|_{q, T, x_0}^2, \end{aligned}$$

where

$$\|\Xi + \nabla \phi_{\xi, \Xi}^T\|_{q, T, x_0} := \left(\sqrt{T}^{-d} \int_{\mathbb{R}^d} |\Xi + \nabla \phi_{\xi, \Xi}^T|^q \exp(-\frac{\gamma}{2} |x - x_0|/\sqrt{T}) dx \right)^{\frac{1}{q}}.$$

Indeed, by subtracting the equations (18a) for $\phi_{\xi, \Xi}^T$ and $\hat{\phi}_{\xi, \Xi}^T$, we get

$$-\nabla \cdot (\hat{a}_\xi^T \nabla \hat{\phi}_{\xi, \Xi}^T) + \frac{1}{T} \hat{\phi}_{\xi, \Xi}^T = \nabla \cdot ((a_\xi^T - \hat{a}_\xi^T)(\Xi + \nabla \phi_{\xi, \Xi}^T)).$$

Note that

$$\begin{aligned} a_\xi^T - \hat{a}_\xi^T &= \partial_\xi A(\tilde{\omega}, \xi + \nabla \phi_\xi^T) - \partial_\xi A(\tilde{\omega}, \xi + \nabla \hat{\phi}_\xi^T) \\ &= (\partial_\xi A(\tilde{\omega}, \xi + \nabla \phi_\xi^T) - \partial_\xi \hat{A}(\tilde{\omega}, \xi + \nabla \phi_\xi^T)) \\ &\quad + (\partial_\xi A(\tilde{\omega}, \xi + \nabla \phi_\xi^T) - \partial_\xi A(\tilde{\omega}, \xi + \nabla \hat{\phi}_\xi^T)) \\ &=: F_1 + F_2. \end{aligned}$$

By exponential localization in form of Lemma 36, the Lipschitz continuity of $\partial_\xi \hat{A}$ (see (R)), the fact that F_1 vanishes on $B_{\frac{L}{4}}$ by $\tilde{\omega} = \hat{\omega}$ on $B_{\frac{L}{4}}$, and the uniform bound

on $\partial_\xi A$ from (A2), we get

$$\begin{aligned}
(84) \quad & \int_{\mathbb{R}^d} \left(|\nabla \hat{\delta} \phi_{\xi, \Xi}^T|^2 + \frac{1}{T} |\hat{\delta} \phi_{\xi, \Xi}^T|^2 \right) \eta(x - x_0) dx \\
& \leq C \int_{\mathbb{R}^d} |F_1|^2 |\Xi + \nabla \phi_{\xi, \Xi}^T|^2 \eta(x - x_0) dx + C \int_{\mathbb{R}^d} |F_2|^2 |\Xi + \nabla \phi_{\xi, \Xi}^T|^2 \eta(x - x_0) dx \\
& \leq C \int_{\{|x| > \frac{L}{4}\}} |\Xi + \nabla \phi_{\xi, \Xi}^T|^2 \eta(x - x_0) dx \\
& \quad + C \int_{\mathbb{R}^d} |\nabla \hat{\delta} \phi_{\xi}^T|^2 |\Xi + \nabla \phi_{\xi, \Xi}^T|^2 \eta(x - x_0) dx.
\end{aligned}$$

We estimate the first term on the RHS by Hölder's inequality as

$$\begin{aligned}
& \int_{\{|x| > \frac{L}{4}\}} |\Xi + \nabla \phi_{\xi, \Xi}^T|^2 \eta(x - x_0) dx \\
& \leq C(d, m, \lambda, \Lambda, q) \sqrt{T}^d \exp\left(-\frac{\gamma}{16} L / \sqrt{T}\right) \|\Xi + \nabla \phi_{\xi, \Xi}^T\|_{q, T, x_0}^2.
\end{aligned}$$

Next, we estimate the second term on the RHS in (84). By Lipschitz continuity of $\partial_\xi A$ in the second variable (see (R)), Hölder's inequality with exponents $p/2$ and $q = \frac{p}{p-2}$ with $0 < p - 2 \ll 1$, Meyers' estimate in form of Lemma 44 for the PDE (80), and (82) from Step 1, we get

$$\begin{aligned}
& \int_{\mathbb{R}^d} |\nabla \hat{\delta} \phi_{\xi}^T|^2 |\Xi + \nabla \phi_{\xi, \Xi}^T|^2 \eta(x - x_0) dx \\
& \leq C \sqrt{T}^d \left(\sqrt{T}^{-d} \int_{\mathbb{R}^d} |\nabla \hat{\delta} \phi_{\xi}^T|^p \eta(x - x_0) dx \right)^{\frac{2}{p}} \|\Xi + \nabla \phi_{\xi, \Xi}^T\|_{q, T, x_0}^2 \\
& \leq C \sqrt{T}^d \exp\left(-\frac{\gamma}{32} L / \sqrt{T}\right) |\xi|^2 \|\Xi + \nabla \phi_{\xi, \Xi}^T\|_{q, T, x_0}^2.
\end{aligned}$$

This completes the argument for (83).

Step 3. Conclusion. Set

$$\zeta(\tilde{\omega}, x) := \left(A(\tilde{\omega}(x), \xi + \nabla \phi_{\xi}^T(\tilde{\omega}, x)) \cdot \Xi - \frac{1}{T} \phi_{\xi}^T(\tilde{\omega}, x) \phi_{\xi, \Xi}^{*, T}(\tilde{\omega}, x) \right).$$

Since η_L is supported in $B_{\frac{L}{8}}$ and $\eta_L \leq L^{-d}$, we have

$$I_1 := \left| (A^{\text{RVE}, \eta_L, T}(\tilde{\omega}, \xi) - A^{\text{RVE}, \eta_L, T}(\hat{\omega}, \xi)) \cdot \Xi \right| \leq CL^{-d} \int_{B_{\frac{L}{8}}} |\zeta(\tilde{\omega}, x) - \zeta(\hat{\omega}, x)| dx.$$

We cover $B_{\frac{L}{8}}$ by balls of radius $\sqrt{T} \leq L$ and centers in $B_{\frac{L}{8}}$; more precisely, there exists a set $X_{L, T} \subset B_{\frac{L}{8}}$ with $\#X_{L, T} \leq C(d)(L/\sqrt{T})^d$ and $\cup_{x_0 \in X_{L, T}} B_{\sqrt{T}}(x_0) \supset B_{\frac{L}{8}}$. Thus,

$$\begin{aligned}
I_1 & \leq CL^{-d} \sum_{x_0 \in X_{L, T}} \int_{B_{\sqrt{T}}(x_0) \cap B_{L/8}} |\zeta(\tilde{\omega}, x) - \zeta(\hat{\omega}, x)| dx \\
& \leq C \sqrt{T}^{-d} \left(\frac{1}{\#X_{L, T}} \sum_{x_0 \in X_{L, T}} \int_{B_{L/8}(x_0) \cap B_{L/8}} |\zeta(\tilde{\omega}, x) - \zeta(\hat{\omega}, x)| \eta(x - x_0) dx \right).
\end{aligned}$$

By the definition of ζ , the estimates in (79) and (83) from Step 1 and Step 2, and the deterministic exponentially localized bounds on $\frac{1}{\sqrt{T}} \phi_{\xi}^T$ and $\frac{1}{\sqrt{T}} \phi_{\xi, \Xi}^{*, T}$ (which are

a consequence of Lemma 36), and the Lipschitz continuity of A with respect to the second variable (see (A2)), we conclude for $x_0 \in X_{L,T} \subset B_{\frac{L}{8}}$ that

$$\begin{aligned}
& \int_{B_{L/8}(x_0) \cap B_{L/8}} |\zeta(\tilde{\omega}, x) - \zeta(\hat{\omega}, x)| \eta(x - x_0) dx \\
& \leq C \int_{\mathbb{R}^d} (|A(\tilde{\omega}, \xi + \nabla \phi_\xi^T) - A(\tilde{\omega}, \xi + \nabla \hat{\phi}_\xi^T)| |\Xi| \\
& \quad + \frac{1}{T} |\phi_\xi^T| |\hat{\delta} \phi_{\xi, \Xi}^{*,T}| + \frac{1}{T} |\hat{\delta} \phi_\xi^T| |\hat{\phi}_{\xi, \Xi}^{*,T}|) \eta(x - x_0) dx \\
& \leq C \left(\int_{\mathbb{R}^d} |\Xi|^2 \eta(x - x_0) dx \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^d} |\nabla \hat{\delta} \phi_\xi^T|^2 \eta(x - x_0) \right)^{\frac{1}{2}} \\
& \quad + C \left(\int_{\mathbb{R}^d} \frac{1}{T} |\phi_\xi^T|^2 \eta(x - x_0) \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^d} \frac{1}{T} |\hat{\delta} \phi_{\xi, \Xi}^{*,T}|^2 \eta(x - x_0) \right)^{\frac{1}{2}} \\
& \quad + C \left(\int_{\mathbb{R}^d} \frac{1}{T} |\hat{\delta} \phi_\xi^T|^2 \eta(x - x_0) \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^d} \frac{1}{T} |\hat{\phi}_{\xi, \Xi}^{*,T}|^2 \eta(x - x_0) \right)^{\frac{1}{2}} \\
& \leq C \sqrt{T}^d \left(\exp(-\frac{\gamma}{32} L / \sqrt{T}) \right)^{\frac{1}{2}} \left((1 + |\xi|^2) |\xi|^2 \|\Xi + \nabla \phi_{\xi, \Xi}^T\|_{q, T, x_0}^2 + |\xi|^2 |\Xi|^2 \right)^{\frac{1}{2}}.
\end{aligned}$$

In total, we have shown the desired deterministic estimate

$$\begin{aligned}
& |A^{\text{RVE}, \eta, T}(\tilde{\omega}, \xi) - A^{\text{RVE}, \eta, T}(\pi_L \tilde{\omega}, \xi)| \\
& \leq C \exp\left(-\frac{\gamma}{64} \cdot \frac{L}{\sqrt{T}}\right) \left(|\xi| |\Xi| + (1 + |\xi|) |\xi| \|\Xi + \nabla \phi_{\xi, \Xi}^T\|_{q, L, T} \right).
\end{aligned}$$

□

4.6. Estimate on the systematic error of the RVE method. We now estimate the systematic error of the RVE approximation for the effective material law. We begin with the case in the presence of the regularity condition (R).

Proof of Theorem 9b – the case with (R). By rescaling we may assume without loss of generality that $\varepsilon = 1$. In the following $\eta : \mathbb{R}^d \rightarrow \mathbb{R}$ denotes a non-negative weight supported in $B_{\frac{L}{8}}$ with $|\eta| \leq C(d)L^{-d}$ and $\int_{\mathbb{R}^d} \eta dx = 1$. Moreover, we consider a localization parameter T according to

$$(85) \quad \sqrt{T} = \frac{\gamma}{64} \frac{L}{\log((L/\varepsilon)^{d \wedge 4})}$$

for the $\gamma = \gamma(d, m, \lambda, \Lambda)$ from Lemma 29. Our starting point is the error decomposition (50), which we recall in the following form

$$\begin{aligned}
& \mathbb{E}_L [A^{\text{RVE}, L}(\xi)] - A_{\text{hom}}(\xi) \\
& = \mathbb{E}_L [A^{\text{RVE}, L}(\xi)] - \mathbb{E}_L [A^{\text{RVE}, \eta, T}(\xi)] \\
& \quad + \mathbb{E}_L [A^{\text{RVE}, \eta, T}(\xi)] - \mathbb{E}_L [A^{\text{RVE}, \eta, T}(\pi_L \omega_\varepsilon, \xi)] \\
& \quad + \mathbb{E} [A^{\text{RVE}, \eta, T}(\pi_L \omega_\varepsilon, \xi)] - \mathbb{E} [A^{\text{RVE}, \eta, T}(\xi)] \\
& \quad + \mathbb{E} [A^{\text{RVE}, \eta, T}(\xi)] - A_{\text{hom}}(\xi) \\
& =: I_1 + I_2 + I_3 + I_4.
\end{aligned}
\tag{86}$$

Note that in the above decomposition we already used the equality

$$\mathbb{E}_L [A^{\text{RVE},\eta,T}(\pi_L \omega_\varepsilon, \xi)] = \mathbb{E} [A^{\text{RVE},\eta,T}(\pi_L \omega_\varepsilon, \xi)],$$

which is valid since \mathbb{P}_L is assumed to be a L -periodic approximation of \mathbb{P} in the sense of Definition 8 (recall also (48)). The terms I_2 and I_3 are coupling errors that can be estimated deterministically with help of Lemma 29. Combined with the choice of T in (85) and the bound on high moments of $\nabla \phi_{\xi, \Xi}^T$ obtained by combining (42) and (23), we arrive at

$$|I_2| + |I_3| \leq C(1 + |\xi|^C) |\xi| |\Xi| \left(\frac{L}{\varepsilon}\right)^{-(d \wedge 4)}.$$

The terms I_1 and I_4 are systematic localization errors that can be estimated with help of Proposition 30b. We obtain using again (85)

$$|I_1| + |I_4| \leq C(1 + |\xi|)^C |\xi| |\Xi| \left(\frac{L}{\varepsilon}\right)^{-(d \wedge 4)} |\log(L/\varepsilon)|^{\alpha d}.$$

Having estimated all terms in (86), this establishes the first estimate in Theorem 9b upon taking the supremum with respect to Ξ , $|\Xi| \leq 1$. \square

We next establish the suboptimal estimate for the systematic error of the RVE method in the case without the small-scale regularity condition (R).

Proof of Theorem 9b – the case without (R). As in the case with small scale regularity (R), we denote by $\eta : \mathbb{R}^d \rightarrow \mathbb{R}$ a non-negative weight supported in $B_{\frac{L}{8}}$ with $|\eta| \leq C(d)L^{-d}$ and $\int_{\mathbb{R}^d} \eta = 1$. Moreover, we consider a localization parameter $\sqrt{T} \leq L$, whose relative scaling with respect to L will be specified below in Step 3. For any parameter field $\tilde{\omega}$ we consider the localized RVE-approximation

$$A^{\text{RVE},\eta,T}(\tilde{\omega}, \xi) := \int \eta A(\tilde{\omega}, \xi + \nabla \phi_\xi^T) dx.$$

Note that it has a simpler form compared to the quantity introduced in Definition 28. In particular, the above expression does not invoke the linearized corrector (for which we cannot derive suitable estimates without the small scale regularity condition (R)). As in the case with small scale regularity, the starting point is estimate (86), i. e. the decomposition of the systematic error

$$\mathbb{E}_L [A^{\text{RVE},L}(\xi)] - A_{\text{hom}}(\xi) = I_1 + I_2 + I_3 + I_4$$

into the two coupling errors

$$\begin{aligned} I_2 &:= \mathbb{E}_L [A^{\text{RVE},\eta,T}(\xi)] - \mathbb{E}_L [A^{\text{RVE},\eta,T}(\pi_L \omega_\varepsilon, \xi)], \\ I_3 &:= \mathbb{E} [A^{\text{RVE},\eta,T}(\pi_L \omega_\varepsilon, \xi)] - \mathbb{E} [A^{\text{RVE},\eta,T}(\xi)], \end{aligned}$$

and the two systematic localization errors

$$\begin{aligned} I_1 &:= \mathbb{E}_L [A^{\text{RVE},L}(\xi)] - \mathbb{E}_L [A^{\text{RVE},\eta,T}(\xi)], \\ I_4 &:= \mathbb{E} [A^{\text{RVE},\eta,T}(\xi)] - A_{\text{hom}}(\xi). \end{aligned}$$

Step 1. Estimate of the coupling errors. We claim that

$$|I_2| + |I_3| \leq C \exp\left(-\frac{\gamma}{32} L/\sqrt{T}\right) |\xi|.$$

Indeed, this can be seen by an argument similar to the proof of Lemma 29. In fact the argument is significantly simpler thanks to the absence of the linearized corrector in the definition of the localized RVE-approximation. We only discuss the argument for I_3 , since the one for I_2 is analogous. We first note that thanks to the assumptions on η and the Lipschitz continuity of A , see (A2), we have

$$(87) \quad \left| A^{\text{RVE},\eta,T}(\omega_\varepsilon, \xi) - A^{\text{RVE},\eta,T}(\pi_L \omega_\varepsilon, \xi) \right| \leq C \int_{B_{\frac{L}{8}}} |\nabla \hat{\delta} \phi_\xi^T|,$$

where $\hat{\delta} \phi_\xi^T$ is defined by (80). As in Section 4.6 we consider a minimal cover of $B_{\frac{L}{8}}$ by balls of radius \sqrt{T} ; more precisely, let $X_{L,T} \subset B_{\frac{L}{8}}$ denote a finite set of points such that $\#X_{L,T} \leq C(d)(L/\sqrt{T})^d$ and $\cup_{x_0 \in X_{L,T}} B_{\sqrt{T}}(x_0) \supset B_{\frac{L}{8}}$. Then

$$\begin{aligned} [\text{RHS of (87)}] &\leq C\sqrt{T}^{-d} \frac{1}{\#X_{L,T}} \sum_{x_0 \in X_{L,T}} \int_{B_{\frac{L}{8}}} |\nabla \hat{\delta} \phi_\xi^T| \exp(-\gamma|x - x_0|/\sqrt{T}) dx \\ &\leq C\sqrt{T}^{-\frac{d}{2}} \frac{1}{\#X_{L,T}} \sum_{x_0 \in X_{L,T}} \left(\int_{\mathbb{R}^d} |\nabla \hat{\delta} \phi_\xi^T|^2 \exp(-\gamma|x - x_0|/\sqrt{T}) dx \right)^{\frac{1}{2}} \end{aligned}$$

where $0 < \gamma \ll 1$ is chosen such that the exponential localization estimate of Lemma 36 applies. Combining the estimate with (79) thus yields

$$\begin{aligned} &\left| A^{\text{RVE},\eta,T}(\omega_\varepsilon, \xi) - A^{\text{RVE},\eta,T}(\pi_L \omega_\varepsilon, \xi) \right| \\ &\leq C\sqrt{T}^{-d/2} \sup_{x_0 \in B_{\frac{L}{8}}} \left(\int_{\mathbb{R}^d} |\nabla \hat{\delta} \phi_\xi^T|^2 \exp(-\gamma|x - x_0|/\sqrt{T}) \right)^{\frac{1}{2}} \\ &\leq C \exp\left(-\frac{\gamma}{32} L/\sqrt{T}\right) |\xi|. \end{aligned}$$

Step 2. Estimate of the systematic localization errors. We claim that

$$|I_1| + |I_4| \leq C|\xi| \left(\frac{\varepsilon}{\sqrt{T}} \right)^{\frac{d \wedge 4}{2}} \begin{cases} |\log \sqrt{T}|^{\frac{1}{2}} & \text{for } d \in \{2, 4\}, \\ 1 & \text{for } d = 3 \text{ and } d \geq 5. \end{cases}$$

We only discuss I_4 , since the argument for I_1 is similar. The Lipschitz continuity of A (see (A2)) and the localization error estimate for the corrector in form of Lemma 31 yield

$$\begin{aligned} &\left| \mathbb{E} [A^{\text{RVE},\eta,2T}(\xi) - A^{\text{RVE},\eta,T}(\xi)] \right| \\ &\leq C\mathbb{E} [|\nabla \phi_\xi^{2T} - \nabla \phi_\xi^T|^2]^{\frac{1}{2}} \\ &\leq C|\xi| \left(\frac{\varepsilon}{\sqrt{T}} \right)^{\frac{d \wedge 4}{2}} \times \begin{cases} |\log(\varepsilon/\sqrt{T})|^{\frac{1}{2}} & \text{for } d \in \{2, 4\}, \\ 1 & \text{for } d = 3 \text{ and } d \geq 5. \end{cases} \end{aligned}$$

The claimed estimate now follows by a telescopic sum argument similar to Step 2 of the proof of Proposition 30.

Step 3. Conclusion. The combination of the previous steps yields

$$\begin{aligned} & \frac{|I_1| + |I_2| + |I_3| + |I_4|}{|\xi|} \\ & \leq C \exp\left(-\frac{\gamma}{32} \frac{L}{\sqrt{T}}\right) + \left(\frac{\varepsilon}{\sqrt{T}}\right)^{\frac{d\wedge 4}{2}} \times \begin{cases} |\log(\sqrt{T}/\varepsilon)|^{\frac{1}{2}} & \text{for } d \in \{2, 4\}, \\ 1 & \text{otherwise.} \end{cases} \end{aligned}$$

With $\sqrt{T} = \frac{\gamma}{32} L (\log((L/\varepsilon)^{\frac{d\wedge 4}{2}}))^{-1}$ the RHS turns into

$$C \left(\frac{\varepsilon}{L}\right)^{\frac{d\wedge 4}{2}} + C \left(\frac{\varepsilon}{L}\right)^{\frac{d\wedge 4}{2}} (\log(L/\varepsilon))^{\frac{d\wedge 4}{2}} \times \begin{cases} |\log L|^{\frac{1}{2}} & \text{for } d \in \{2, 4\}, \\ 1 & \text{for } d = 3 \text{ and } d \geq 5. \end{cases}$$

This establishes the result. \square

5. CORRECTOR ESTIMATES FOR THE NONLINEAR PDE

5.1. Estimates on linear functionals of the corrector and the flux corrector. We now turn to the first key step, the estimate on the localized homogenization corrector ϕ_ξ^T and the localized flux corrector σ_ξ^T . For the proof of the estimate on linear functionals of the corrector and the flux corrector in Lemma 17, we need the following auxiliary lemma, which follows by a combination of the Caccioppoli inequality with hole-filling.

Lemma 38. *Let the assumptions (A1)–(A2) be satisfied, let $\varepsilon > 0$ be arbitrary, let $\tilde{\omega}$ be an arbitrary parameter field, let the correctors ϕ_ξ^T and $\phi_{\xi,\Xi}^T$ be defined as the unique solutions to the corrector equations (12a) and (18a) in H_{uloc}^1 , and let $r_{*,T,\xi}$ and $r_{*,T,\xi,\Xi}$ be as in (36) and (39). Let the parameter K_{mass} in (36), (39) be chosen as $K_{\text{mass}} \geq C(d, m, \lambda, \Lambda)$ for some C determined in the proof below. Then for any $x_0 \in \mathbb{R}^d$ the estimates*

$$(88) \quad \int_{B_\varepsilon(x_0)} |\xi + \nabla \phi_\xi^T|^2 dx \leq C |\xi|^2 \left(\frac{r_{*,T,\xi}(x_0)}{\varepsilon} \right)^{d-\delta}$$

and

$$(89) \quad \int_{B_\varepsilon(x_0)} |\Xi + \nabla \phi_{\xi,\Xi}^T|^2 dx \leq C |\Xi|^2 \left(\frac{r_{*,T,\xi,\Xi}(x_0)}{\varepsilon} \right)^{d-\delta}$$

hold true. Furthermore, we have

$$(90) \quad r_{*,T,\xi}(x_0) \leq C\sqrt{T},$$

$$(91) \quad r_{*,T,\xi,\Xi}(x_0) \leq C\sqrt{T}.$$

Proof. The estimates (90) and (91) are a simple consequence of the definitions (36) and (39), the assumed lower bound on K_{mass} , and the bounds

$$\begin{aligned} & \int_{B_R} \frac{1}{T} |\phi_\xi^T|^2 dx \leq C |\xi|^2 \\ & \int_{B_R} \frac{1}{T} |\phi_{\xi,\Xi}^T|^2 dx \leq C |\Xi|^2 \end{aligned}$$

valid for all $R \geq \sqrt{T}$ which are a consequence of Lemma 36 applied to the corrector equations (12a) and (18a).

To prove (88), we combine the Caccioppoli inequality (51) with the hole-filling estimate (52) for the function $u(x) := \xi \cdot (x - x_0) + \phi_\xi^T$, which by (12a) solves the PDE

$$-\nabla \cdot (A(\tilde{\omega}, \nabla u)) + \frac{1}{T}u = \frac{1}{T}\xi \cdot (x - x_0).$$

This yields

$$\begin{aligned} & \int_{B_\varepsilon(x_0)} |\xi + \nabla \phi_\xi^T|^2 dx \\ & \leq C \left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{d-\delta} \inf_{b \in \mathbb{R}^m} \left(\frac{1}{r_{*,T,\xi}^2} \int_{B_{r_{*,T,\xi}}(x_0)} |\xi \cdot (x - x_0) + \phi_\xi^T - b|^2 dx + \frac{1}{T}|b|^2 \right. \\ & \quad \left. + \int_{B_{r_{*,T,\xi}}(x_0)} \frac{1}{T} |\xi \cdot (x - x_0)|^2 dx \right) \\ & \quad + C \left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^d \int_{B_{r_{*,T,\xi}}(x_0)} \left(\frac{\varepsilon}{\varepsilon + |x - x_0|} \right)^\delta \frac{1}{T} |\xi \cdot (x - x_0)|^2 dx. \end{aligned}$$

Choosing $b := \int_{B_{r_{*,T,\xi}}(x_0)} \phi_\xi^T dx$ and using the definition of $r_{*,T,\xi}$ (36) as well as the fact that $r_{*,T,\xi} \leq C\sqrt{T}$ then entails (88). To show (89), one argues analogously. \square

We now prove the estimate in Lemma 17 for functionals of the localized corrector ϕ_ξ^T for the nonlinear elliptic PDE and the corresponding localized flux corrector σ_ξ^T . As the proof of Lemma 18 is very similar, we combine their proofs.

Proof of Lemma 17 and Lemma 18. Part a: Estimates for linear functionals of the homogenization corrector ϕ_ξ^T . Taking the derivative of the corrector equation (12a) with respect to a perturbation $\delta\omega_\varepsilon$ of the random field ω_ε , we see that the change $\delta\phi_\xi^T$ of the corrector under such an infinitesimal perturbation satisfies the linear elliptic PDE

$$(92) \quad -\nabla \cdot (\partial_\xi A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \nabla \delta\phi_\xi^T) + \frac{1}{T} \delta\phi_\xi^T = \nabla \cdot (\partial_\omega A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \delta\omega_\varepsilon).$$

Define the coefficient field $a_\xi^T(x) := \partial_\xi A(\omega_\varepsilon(x), \xi + \nabla \phi_\xi^T(x))$. By our assumptions (A1) and (A2), the coefficient field $a_\xi^T : \mathbb{R}^d \rightarrow (\mathbb{R}^{m \times d})^2$ exists and is uniformly elliptic and bounded in the sense $a_\xi^T v \cdot v \geq \lambda|v|^2$ and $|a_\xi^T v| \leq \Lambda|v|$ for any $v \in \mathbb{R}^{m \times d}$.

Now, consider a functional of the form $F := \int_{\mathbb{R}^d} g \cdot \nabla \phi_\xi^T dx$ for a deterministic compactly supported function g . Denoting by $h \in H^1(\mathbb{R}^d; \mathbb{R}^m)$ the (unique) solution to the dual equation

$$(93) \quad -\nabla \cdot (a_\xi^{T,*} \nabla h) + \frac{1}{T} h = \nabla \cdot g,$$

we deduce

$$\begin{aligned} \delta F &= \int_{\mathbb{R}^d} g \cdot \nabla \delta\phi_\xi^T dx \stackrel{(93)}{=} - \int_{\mathbb{R}^d} a_\xi^T \nabla \delta\phi_\xi^T \cdot \nabla h + \frac{1}{T} \delta\phi_\xi^T \cdot h dx \\ &\stackrel{(92)}{=} \int_{\mathbb{R}^d} \partial_\omega A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \delta\omega_\varepsilon \cdot \nabla h dx, \end{aligned}$$

i. e. we have

$$(94) \quad \frac{\partial F}{\partial \omega_\varepsilon} = \partial_\omega A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \cdot \nabla h.$$

By the assumption (A3), this implies

$$\begin{aligned} & \int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} \left| \frac{\partial F}{\partial \omega_\varepsilon} \right| d\tilde{x} \right|^2 dx \\ & \leq C \int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T| |\nabla h| d\tilde{x} \right|^2 dx. \end{aligned}$$

Using the version of the spectral gap inequality for the q -th moment (see Lemma 16), we deduce for any $q \geq 1$

$$\mathbb{E} \left[|F - \mathbb{E}[F]|^{2q} \right]^{1/2q} \leq C q \varepsilon^{d/2} \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T| |\nabla h| d\tilde{x} \right|^2 dx \right)^q \right]^{1/2q}.$$

As ϕ_ξ^T is a stationary vector field, we have $\mathbb{E}[F] = \int_{\mathbb{R}^d} g \cdot \nabla \mathbb{E}[\phi_\xi^T] dx = 0$. By (88), we have

$$\int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T|^2 d\tilde{x} \leq C |\xi|^2 \left(\frac{r_{*,T,\xi}(x_0)}{\varepsilon} \right)^{d-\delta} =: B^2(x).$$

An application of Lemma 39 therefore yields for any $0 < \tau < 1$, any $0 < \alpha_0 < c$, any $p \in (2, 2 + c(d, m, \lambda, \Lambda, \alpha_0))$, and any $q \geq C(d, m, \lambda, \Lambda, \tau, \alpha_0, p)$

$$(95) \quad \mathbb{E} [|F|^{2q}]^{1/2q} \leq C |\xi| q \varepsilon^{d/2} r^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q} \\ \times \mathbb{E} \left[\left(r^{-d} \int_{\mathbb{R}^d} |\nabla h|^p \left(1 + \frac{|x|}{r} \right)^{\alpha_0} dx \right)^{2q/p\tau} \right]^{\tau/2q}.$$

Applying the weighted Meyers estimate of Lemma 35 to the equation (93) and using our assumption $(\int_{B_r(x_0)} |g|^p dx)^{1/p} \leq r^{-d}$, the integral in the last term may be estimated to yield

$$\mathbb{E} [|F|^{2q}]^{1/2q} \leq C |\xi| q \left(\frac{\varepsilon}{r} \right)^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q}$$

for any $2 < p < 2 + c$, any $0 < \tau < 1$, and any $q \geq C(d, m, \lambda, \Lambda, p, \tau)$. This is the desired estimate in Lemma 17a.

Next, we establish (38). To this end, we consider the random variable $F := \int_{B_R(x_0)} \phi_\xi^T dx$. By $\mathbb{E}[\phi_\xi^T] = 0$ – which follows by testing the PDE (12a) with a test function, taking the expectation, and using stationarity – we see that we have $\mathbb{E}[F] = 0$. We may therefore repeat the previous computation to estimate the stochastic moments of F , up to the following changes: The equation (93) is replaced by the equation with non-divergence form right-hand side $-\nabla \cdot (a_\xi^{T,*} \nabla h) + \frac{1}{T} h = \frac{1}{|B_R|} \chi_{B_R(x_0)}$, and the estimate for ∇h deduced from Lemma 35 now reads

$$\begin{aligned} & \left(R^{-d} \int_{\mathbb{R}^d} |\nabla h|^p \left(1 + \frac{|x|}{R} \right)^{\alpha_0} dx \right)^{1/p} \\ & \leq C \sqrt{T} \left(R^{-d} \int_{\mathbb{R}^d} \left| \frac{1}{|B_R|} \chi_{B_R(x_0)} \right|^p dx \right)^{1/p}. \end{aligned}$$

In total, we deduce

$$\mathbb{E} \left[\left| \int_{B_R(x_0)} \phi_\xi^T dx \right|^q \right]^{1/q} \leq C |\xi| q \sqrt{T} \left(\frac{\varepsilon}{R} \right)^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q}.$$

This is precisely the desired estimate on the average of ϕ_ξ^T from (38).

Part b: Estimates for linear functionals of the flux corrector σ_ξ^T . Taking the derivative of the flux corrector equation (12c) with respect to a perturbation $\delta\omega_\varepsilon$ of the random field ω_ε , we see that the change $\delta\sigma_\xi^T$ of the corrector under such an infinitesimal perturbation satisfies the linear elliptic PDE

$$(96) \quad \begin{aligned} -\Delta \delta\sigma_{\xi,jk}^T + \frac{1}{T} \delta\sigma_{\xi,jk}^T &= \nabla \cdot (\partial_\omega A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T) \omega_\varepsilon \cdot (e_k \otimes e_j - e_j \otimes e_k)) \\ &\quad + \nabla \cdot (\partial_\xi A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T) \nabla\delta\phi_\xi^T \cdot (e_k \otimes e_j - e_j \otimes e_k)). \end{aligned}$$

For the sensitivity of the functional

$$F_\sigma = \int_{\mathbb{R}^d} g \cdot \nabla\sigma_{\xi,jk}^T dx,$$

we obtain by defining $\bar{h} \in H^1(\mathbb{R}^d; \mathbb{R}^{m \times d})$ as the unique decaying solution to the (system of) Poisson equation(s) with massive term

$$(97) \quad -\Delta \bar{h} + \frac{1}{T} \bar{h} = \nabla \cdot g$$

and defining $\hat{h} \in H^1(\mathbb{R}^d; \mathbb{R}^{m \times d})$ as the unique decaying solution to the uniformly elliptic PDE (with $a_\xi^T(x) := \partial_\xi A(\omega_\varepsilon(x), \xi + \nabla\phi_\xi^T)$)

$$(98) \quad -\nabla \cdot (a_\xi^{T,*} \nabla \hat{h}) + \frac{1}{T} \hat{h} = \nabla \cdot ((\partial_j \bar{h} e_k - \partial_k \bar{h} e_j) \cdot \partial_\xi A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T))$$

that

$$\begin{aligned} \delta F_\sigma &= \int_{\mathbb{R}^d} g \cdot \nabla \delta\sigma_{\xi,jk}^T dx \stackrel{(97)}{=} - \int_{\mathbb{R}^d} \nabla \bar{h} \cdot \nabla \delta\sigma_{\xi,jk}^T + \frac{1}{T} \bar{h} \delta\sigma_{\xi,jk}^T dx \\ &\stackrel{(96)}{=} \int_{\mathbb{R}^d} \nabla \bar{h} \cdot (\partial_\omega A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T) \delta\omega_\varepsilon \cdot (e_k \otimes e_j - e_j \otimes e_k)) dx \\ &\quad + \int_{\mathbb{R}^d} \nabla \bar{h} \cdot (\partial_\xi A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T) \nabla\delta\phi_\xi^T \cdot (e_k \otimes e_j - e_j \otimes e_k)) dx \\ &\stackrel{(98)}{=} \int_{\mathbb{R}^d} \nabla \bar{h} \cdot (\partial_\omega A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T) \delta\omega_\varepsilon \cdot (e_k \otimes e_j - e_j \otimes e_k)) dx \\ &\quad - \int_{\mathbb{R}^d} \partial_\xi A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T) \nabla\delta\phi_\xi^T \cdot \nabla \hat{h} + \frac{1}{T} \delta\phi_\xi^T \hat{h} dx \\ &\stackrel{(92)}{=} \int_{\mathbb{R}^d} \nabla \bar{h} \cdot (\partial_\omega A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T) \delta\omega_\varepsilon \cdot (e_k \otimes e_j - e_j \otimes e_k)) dx \\ &\quad + \int_{\mathbb{R}^d} \partial_\omega A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T) \delta\omega_\varepsilon \cdot \nabla \hat{h} dx. \end{aligned}$$

i. e. we have

$$\frac{\partial F}{\partial \omega_\varepsilon} = (\partial_j \bar{h} e_k - \partial_k \bar{h} e_j) \cdot \partial_\omega A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T) + \nabla \hat{h} \cdot \partial_\omega A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T).$$

Using again the version of the spectral gap inequality for the q -th moment from Lemma 16, the fact that $\mathbb{E}[F_\sigma] = \int_{\mathbb{R}^d} g \cdot \nabla \mathbb{E}[\sigma_{\xi,jk}^T] dx = 0$ by stationarity, and the bound $|\partial_\omega A(\omega, \xi)| \leq \Lambda|\xi|$ (recall (A1)-(A2)), we obtain

$$\mathbb{E}[|F_\sigma|^{2q}]^{1/2q} \leq Cq\varepsilon^{d/2} \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T| (|\nabla \bar{h}| + |\nabla \hat{h}|) d\tilde{x} \right|^2 dx \right)^q \right]^{1/2q}.$$

By (88), we have

$$\int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T|^2 d\tilde{x} \leq C|\xi|^2 \left(\frac{r_{*,T,\xi}(x_0)}{\varepsilon} \right)^{d-\delta} =: B^2(x).$$

By a combination of Hölder's and Jensen's inequality in form of Lemma 39 (see below) we deduce that for any $\alpha_0 \in (0, c)$, any $p \in (2, 2 + c(d, m, \lambda, \Lambda, \alpha_0))$, any $\tau \in (0, 1)$, and any $q \geq q(d, m, \lambda, \Lambda, \alpha_0, p)$

$$\begin{aligned} \mathbb{E}[|F_\sigma|^{2q}]^{1/2q} &\leq C|\xi|q\varepsilon^{d/2}r^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q} \\ &\quad \times \mathbb{E} \left[\left(r^{-d} \int_{\mathbb{R}^d} (|\nabla \bar{h}|^p + |\nabla \hat{h}|^p) \left(1 + \frac{|x|}{r} \right)^{\alpha_0} dx \right)^{2q/p\tau} \right]^{\tau/2q}. \end{aligned}$$

By the weighted Meyers estimate of Lemma 35 and the defining equation (98) for \hat{h} as well as the uniform bound on $\partial_\xi A$ inferred from (A2), we deduce for any $\alpha_{1/2} \in (\alpha_0, c)$

$$\begin{aligned} \mathbb{E}[|F_\sigma|^{2q}]^{1/2q} &\leq C|\xi|q\varepsilon^{d/2}r^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q} \\ &\quad \times \mathbb{E} \left[\left(r^{-d} \int_{\mathbb{R}^d} |\nabla \bar{h}|^p \left(1 + \frac{|x|}{r} \right)^{\alpha_{1/2}} dx \right)^{2q/p\tau} \right]^{\tau/2q}. \end{aligned}$$

Applying Lemma 35 to the equation (97) and using $(\int_{B_r(x_0)} |g|^p dx)^{1/p} \leq r^{-d}$, the integral in the last term may be estimated to yield

$$\mathbb{E}[|F_\sigma|^{2q}]^{1/2q} \leq C|\xi|q \left(\frac{\varepsilon}{r} \right)^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q}.$$

This is the desired estimate in Lemma 17b.

As in Part a, the estimate on the average of σ_ξ^T from (38) follows upon replacing the equation for \bar{h} by the PDE $-\Delta \bar{h} + \frac{1}{T} \bar{h} = \frac{1}{|B_R|} \chi_{B_R(x_0)}$ in the previous computation.

Part c: Estimates on linear functionals of the potential $\theta_{\xi,i}^T$. Taking the derivative of the equation (19a) with respect to a perturbation $\delta\omega_\varepsilon$ of the random field ω_ε , we get

$$(99) \quad \Delta \delta\theta_{\xi,i}^T = \partial_i \delta\phi_\xi^T.$$

Defining $\bar{g} \in H^1(\mathbb{R}^d; \mathbb{R}^m)$ as the unique decaying solution to the PDE

$$(100) \quad -\Delta \bar{g} = \nabla \cdot g,$$

we obtain

$$\delta F_\theta = \int_{\mathbb{R}^d} g \cdot \nabla \delta\theta_{\xi,i}^T dx \stackrel{(100)}{=} - \int_{\mathbb{R}^d} \nabla \bar{g} \cdot \nabla \delta\theta_{\xi,i}^T dx \stackrel{(99)}{=} \int_{\mathbb{R}^d} \bar{g} \cdot (e_i \cdot \nabla) \delta\phi_\xi^T dx.$$

In other words, we are back in the situation of Part a, the only difference being that the function g has been replaced by the function $\bar{g}e_i$. The function \bar{g} now no longer has compact support; rather, being the solution to the Poisson equation (100) in $d \geq 3$ dimensions, for any $0 < \alpha_1 < 1$ and any $p \geq 2$ it satisfies the weighted Calderon-Zygmund estimate

$$\left(\int_{B_R(x_0)} |\bar{g}|^p \left(1 + \frac{|x - x_0|}{r}\right)^{\alpha_1} dx \right)^{1/p} \leq Cr \left(\int_{B_r(x_0)} |g|^p dx \right)^{1/p}.$$

Now, we may follow the estimates from Part a leading to (95) line by line. In the next step, we again employ Lemma 35, which by the previous estimate on \bar{g} gives

$$\mathbb{E} [|F_\theta|^{2q}]^{1/2q} \leq Cr |\xi| q \left(\frac{\varepsilon}{r}\right)^{d/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon}\right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q}.$$

This is our desired estimate.

Part d: Proof of the estimate (37). Let $w \in H^1(\mathbb{R}^d)$ be arbitrary. Assuming for the moment $R = 2^N r$ for some $N \in \mathbb{N}$, we may write

$$\left| \int_{B_R(x_0)} w dx - \int_{B_r(x_0)} w dx \right|^2 = \sum_{n=1}^N |\langle f_n, w \rangle_{L^2}|^2$$

with $f_n := \frac{1}{|B_{2^n r}|} \chi_{B_{2^n r}(x_0)} - \frac{1}{|B_{2^{n-1} r}|} \chi_{B_{2^{n-1} r}(x_0)}$, where we have used the orthogonality of the functions f_n . Solving the PDEs $\Delta v = f_n$ on $B_{2^n r}(x_0)$ with Neumann boundary conditions, we see that $g_n := \nabla v \chi_{B_{2^n r}(x_0)}$ solves $\nabla \cdot g_n = f_n$ on \mathbb{R}^d and satisfies for any $p \geq 2$ the estimate $(\int_{B_{2^n r}(x_0)} |g_n|^p dx)^{1/p} \leq C(2^n r)^{1-d}$. This implies

$$\mathbb{E} \left[\left| \int_{B_R(x_0)} w dx - \int_{B_r(x_0)} w dx \right|^{2q} \right]^{1/q} \leq \sum_{n=1}^N \mathbb{E} \left[\left| \int_{\mathbb{R}^d} g_n \cdot \nabla w \right|^{2q} \right]^{1/q}.$$

Combining this estimate with the bounds on g_n and the estimates from Lemma 17, we deduce the estimate (37) if R is of the form $R = 2^N r$. If R is not of this form, we need one additional step to estimate the difference of the averages on the radius $2^{\lfloor \log_2 \frac{R}{r} \rfloor} r$ and R . □

The following estimate is a consequence of a straightforward application of Hölder's inequality and Jensen's inequality. Nevertheless, we state it to avoid repetition of the same computations, as it is repeatedly used in the proofs of Lemma 17 and Lemma 21.

Lemma 39. *Let $\varepsilon > 0$ and let b and B be stationary random fields with*

$$\int_{B_\varepsilon(x)} |b|^2 d\tilde{x} \leq |B(x)|^2.$$

Let h be a random field satisfying $\nabla h \in L^p(\mathbb{R}^d)$ almost surely. Let $0 < \alpha_0 \leq 1$ and $0 < \tau < 1$. Let $p > 2$ be close enough to 2 depending only on d and on (a lower bound on) α_0 , but otherwise arbitrary. Then for any $q \geq C(p, \alpha_0, \tau)$ and for any

$r \geq \varepsilon$ the estimate

$$\begin{aligned} & \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} |b| |\nabla h| d\tilde{x} \right|^2 dx \right)^q \right]^{1/2q} \\ & \leq C(d, \alpha_0, p, \tau) r^{d/2} \mathbb{E} \left[|B|^{2q/(1-\tau)} \right]^{(1-\tau)/2q} \\ & \quad \times \mathbb{E} \left[\left(r^{-d} \int_{\mathbb{R}^d} |\nabla h|^p \left(1 + \frac{|x|}{r} \right)^{\alpha_0} dx \right)^{2q/p\tau} \right]^{\tau/2q} \end{aligned}$$

holds true.

Proof of Lemma 39. Several applications of Hölder's inequality yield together with the bound on b

$$\begin{aligned} & \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} |b| |\nabla h| d\tilde{x} \right|^2 dx \right)^q \right]^{1/2q} \\ & \leq \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \int_{B_\varepsilon(x)} |b|^2 d\tilde{x} \int_{B_\varepsilon(x)} |\nabla h|^2 d\tilde{x} dx \right)^q \right]^{1/2q} \\ & \leq \mathbb{E} \left[\left(\int_{\mathbb{R}^d} |B(x)|^2 \int_{B_\varepsilon(x)} |\nabla h|^2 d\tilde{x} dx \right)^q \right]^{1/2q} \\ & \leq r^{d/2} \mathbb{E} \left[\left(r^{-d} \int_{\mathbb{R}^d} |B(x)|^{2p/(p-2)} \left(1 + \frac{|x|}{r} \right)^{-\alpha_0/(p-2)} dx \right)^{q(p-2)/p} \right. \\ & \quad \left. \times \left(r^{-d} \int_{\mathbb{R}^d} \left(\int_{B_\varepsilon(x)} |\nabla h|^2 d\tilde{x} \right)^{p/2} \left(1 + \frac{|x|}{r} \right)^{\alpha_0} dx \right)^{2q/p} \right]^{1/2q}. \end{aligned}$$

This implies using Hölder's inequality with exponents $\frac{1}{1-\tau}$ and $\frac{1}{\tau}$ and Jensen's inequality

$$\begin{aligned} & \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} |b| |\nabla h| d\tilde{x} \right|^2 dx \right)^q \right]^{1/2q} \\ & \leq r^{d/2} \mathbb{E} \left[\left(r^{-d} \int_{\mathbb{R}^d} |B(x)|^{2p/(p-2)} \left(1 + \frac{|x|}{r} \right)^{-\alpha_0/(p-2)} dx \right)^{q(p-2)/p(1-\tau)} \right]^{(1-\tau)/2q} \\ & \quad \times \mathbb{E} \left[\left(r^{-d} \int_{\mathbb{R}^d} \int_{B_\varepsilon(x)} |\nabla h|^p d\tilde{x} \left(1 + \frac{|x|}{r} \right)^{\alpha_0} dx \right)^{2q/p\tau} \right]^{\tau/2q}. \end{aligned}$$

Assuming that $q \geq p/(p-2)$ and $\alpha_0/(p-2) \geq 2d$, we obtain using Jensen's inequality in the first term (note that the integral $r^{-d} \int_{\mathbb{R}^d} \left(1 + \frac{|x|}{r} \right)^{\alpha_0/(p-2)} dx$ is bounded by $C(d, \alpha_0, p)$) as well as the fact that the supremum and the infimum of

the function $1 + \frac{|x|}{r}$ on an ε -ball differ by at most a factor of 2 in the second term

$$\begin{aligned} & \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} |b| |\nabla h| d\tilde{x} \right|^2 dx \right)^q \right]^{1/2q} \\ & \leq C(d, \alpha_0, p, \tau) r^{d/2} \mathbb{E} \left[r^{-d} \int_{\mathbb{R}^d} |B(x)|^{2q/(1-\tau)} \left(1 + \frac{|x|}{r} \right)^{-\alpha_0/(p-2)} dx \right]^{(1-\tau)/2q} \\ & \quad \times \mathbb{E} \left[\left(r^{-d} \int_{\mathbb{R}^d} |\nabla h|^p \left(1 + \frac{|x|}{r} \right)^{\alpha_0} dx \right)^{2q/p\tau} \right]^{\tau/2q}. \end{aligned}$$

By the stationarity of B , this yields the assertion of the lemma. \square

5.2. Estimates on the corrector for the monotone system. We will need the following technical lemma.

Lemma 40. *Let $R > 0$ and $K \in \mathbb{N}$. For any function $v \in H^1([-R, R]^d; \mathbb{R}^m)$ the estimate*

$$\begin{aligned} & \int_{[-R, R]^d} \left| v - \int_{[-R, R]^d} v d\tilde{x} \right|^2 dx \\ & \leq C \sum_{k \in \{-K, \dots, 0, \dots, K\}^d} R^2 \left| \int_{[-R, R]^d} \prod_{i=1}^d \cos \left(\frac{\pi k_i (x_i + R)}{2R} \right) \nabla \phi_\xi^T dx \right|^2 \\ & \quad + \frac{C}{K^2} R^2 \int_{[-R, R]^d} |\nabla v|^2 dx \end{aligned}$$

holds.

Proof. The proof is an elementary consequence of the Fourier series representation of v . \square

Combining the estimates on linear functionals of the corrector from Lemma 17 with the estimate on the corrector gradient (88) and the technical Lemma 40, we now derive stochastic moment bounds on the minimal radius $r_{*,T,\xi}$.

Proof of Lemma 19. In order to obtain a bound for the minimal radius $r_{*,T,\xi}$, we derive an estimate on the probability of the event $r_{*,T,\xi}(x_0) = R = 2^\ell \varepsilon$ for a fixed $x_0 \in \mathbb{R}^d$ and any $R = 2^\ell \varepsilon > \varepsilon$. In the case of this event, we have by the Caccioppoli inequality (51) applied to the function $\xi \cdot (x - x_0) + \phi_\xi^T$, which solves the PDE

$$-\nabla \cdot (A(\omega_\varepsilon, \nabla(\xi \cdot (x - x_0) + \phi_\xi^T))) + \frac{1}{T}(\xi \cdot (x - x_0) + \phi_\xi^T) = -\frac{1}{T}\xi \cdot (x - x_0),$$

the definition of $r_{*,T,\xi}(x_0)$ in (36), and the fact that $r_{*,T,\xi} \leq C\sqrt{T}$ (see (90))

$$(101) \quad \int_{B_{dR}(x_0)} |\nabla \phi_\xi^T| dx \leq C|\xi|^2.$$

Furthermore, in the event $r_{*,T,\xi}(x_0) = R > \varepsilon$ we also know by the definition (36) that at least one of the inequalities

$$\begin{aligned}
(102) \quad & \frac{1}{R^2} \int_{x_0+[-R,R]^d} \left| \phi_\xi^T - \int_{x_0+[-R,R]^d} \phi_\xi^T d\tilde{x} \right|^2 dx \\
&= \inf_{b \in \mathbb{R}^m} \frac{1}{R^2} \int_{x_0+[-R,R]^d} |\phi_\xi^T - b|^2 dx \\
&\geq c \inf_{b \in \mathbb{R}^m} \frac{1}{(R/2)^2} \int_{B_{R/2}(x_0)} |\phi_\xi^T - b|^2 dx \\
&> c|\xi|^2
\end{aligned}$$

or

$$(103) \quad \frac{1}{\sqrt{T}} \left| \int_{B_{R/2}(x_0)} \phi_\xi^T dx \right| > K_{mass} |\xi|$$

holds. We now distinguish these two cases.

Case 1: The estimate (102) holds. By Lemma 40, we have for any $\delta > 0$ for a sufficiently large $K = K(d, \delta)$

$$\begin{aligned}
(104) \quad & \frac{1}{R^2} \int_{x_0+[-R,R]^d} \left| \phi_\xi^T - \int_{x_0+[-R,R]^d} \phi_\xi^T d\tilde{x} \right|^2 dx \\
&\leq \delta \int_{x_0+[-R,R]^d} |\nabla \phi_\xi^T|^2 dx \\
&+ C \sum_{k \in \{-K, \dots, 0, \dots, K\}^d} \left| \int_{x_0+[-R,R]^d} \prod_{i=1}^d \cos\left(\frac{\pi k_i(x_i + R)}{2R}\right) \nabla \phi_\xi^T dx \right|^2.
\end{aligned}$$

Now let $g_{n,R}$ be the family of all functions $(2R)^{-d} \prod_{i=1}^d \cos\left(\frac{\pi k_i(x_i + R)}{2R}\right) e_l \otimes e_j$, $1 \leq l \leq m$, $1 \leq j \leq d$, and $k \in \{-K, \dots, K\}^d$. Note that all $g_{n,R}$ are supported in $B_{dR}(x_0)$ and satisfy for any $p \in [1, \infty]$

$$(105) \quad \left(\int_{B_{dR}(x_0)} |g_{n,R}|^p dx \right)^{1/p} \leq C(d, p)(dR)^{-d}.$$

Inserting both (102) and (101) in (104) and choosing $\delta > 0$ small enough (depending only on d and the constants c and C from (102) and (101)), we obtain

$$\frac{1}{2}c|\xi|^2 \leq C \sum_{k \in \{-K, \dots, 0, \dots, K\}^d} \left| \int_{x_0+[-R,R]^d} \prod_{i=1}^d \cos\left(\frac{\pi k_i(x_i + R)}{2R}\right) \nabla \phi_\xi^T dx \right|^2.$$

This implies that for at least one of the $N = N(d, K)$ functionals $\int_{\mathbb{R}^d} g_{n,R} \cdot \nabla \phi_\xi^T dx$ we have

$$\left| \int_{\mathbb{R}^d} g_{n,R} \cdot \nabla \phi_\xi^T dx \right| \geq c(d, m, \lambda, \Lambda, K) |\xi|.$$

Fixing δ and K depending only on d , m , λ , and Λ , this entails for any $q \geq 1$

$$\mathbb{P}[r_{*,T,\xi}(x_0) = R \text{ and (102) holds}] \leq \sum_{n=1}^N \frac{\mathbb{E} \left[\left| \int_{\mathbb{R}^d} g_{n,R} \cdot \nabla \phi_\xi^T dx \right|^q \right]}{c^q |\xi|^q}$$

with $c = c(d, m, \lambda, \Lambda)$. Estimating the weighted averages of $\nabla \phi_\xi^T$ by means of Lemma 17 – note that by (105) and the support property of the $g_{n,R}$ the lemma is indeed applicable to the function $\frac{g_{n,R}}{C(d,p)}$ – we obtain for any $0 < \tau < 1$ and any $q \geq C(d, \tau)$

$$(106) \quad \begin{aligned} & \mathbb{P}[r_{*,T,\xi}(x_0) = R \text{ and (102) holds}] \\ & \leq C^q q^q \left(\frac{\varepsilon}{R}\right)^{qd/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/2(1-\tau)} \right]^{(1-\tau)}. \end{aligned}$$

Case 2: The estimate (103) holds. We first observe that as a consequence of Lemma 36 and (12a) – rewritten in form of $-\nabla \cdot (A(\tilde{\omega}, \nabla \phi_\xi^T)) + \frac{1}{T} \phi_\xi^T = \nabla \cdot \hat{g}$ with $\hat{g} := A(\tilde{\omega}, \xi + \nabla \phi_\xi^T) - A(\tilde{\omega}, \nabla \phi_\xi^T)$ – we have

$$\left| \int_{B_{\sqrt{T}}(x_0)} \phi_\xi^T dx \right| \leq \left(\int_{B_{\sqrt{T}}(x_0)} |\phi_\xi^T|^2 dx \right)^{1/2} \leq C |\xi| \sqrt{T}.$$

Upon choosing $K_{mass} \geq C + 1$, (103) is seen to imply

$$\left| \int_{B_{R/2}(x_0)} \phi_\xi^T dx - \int_{B_{\sqrt{T}}(x_0)} \phi_\xi^T dx \right| \geq |\xi| \sqrt{T}.$$

In particular, we have for any $q \geq 1$

$$\begin{aligned} & \mathbb{P}[r_{*,T,\xi}(x_0) = R \text{ and (103) holds}] \\ & \leq \mathbb{E} \left[\left| \frac{1}{|\xi| \sqrt{T}} \int_{B_{R/2}(x_0)} \phi_\xi^T dx - \frac{1}{|\xi| \sqrt{T}} \int_{B_{\sqrt{T}}(x_0)} \phi_\xi^T dx \right|^q \right]^{1/q} \end{aligned}$$

Inserting the estimate (37), we deduce

$$(107) \quad \begin{aligned} & \mathbb{P}[r_{*,T,\xi}(x_0) = R \text{ and (103) holds}] \\ & \leq C^q q^q \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/2(1-\tau)} \right]^{(1-\tau)} \left(\sum_{l=0}^{\log_2 \frac{\sqrt{T}}{R}} \left(\frac{2^l R}{\sqrt{T}} \right)^2 \left(\frac{\varepsilon}{2^l R} \right)^d \right)^{q/2} \\ & \leq C^q q^q \left(\frac{\varepsilon}{R}\right)^{qd/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/2(1-\tau)} \right]^{(1-\tau)}. \end{aligned}$$

Conclusion: Estimates on the minimal radius. Taking the sum of (106) and (107) over all dyadic $R = 2^k \varepsilon > \varepsilon$ and using the fact that $r_{*,T,\xi} \geq \varepsilon$ by its definition (36), we deduce

$$\begin{aligned} & \mathbb{E} \left[\left(\frac{r_{*,T,\xi}(x_0)}{\varepsilon} \right)^{q(d-\delta/2)/2} \right] \\ & \leq 1 + \sum_{k=1}^{\infty} (2^k)^{q(d-\delta/2)/2} \mathbb{P}[r_{*,T,\xi}(x_0) = 2^k \varepsilon] \\ & \stackrel{(106),(107)}{\leq} 1 + C^q q^q \sum_{k=1}^{\infty} (2^k)^{q(d-\delta/2)/2} \left(\frac{\varepsilon}{\varepsilon 2^k}\right)^{qd/2} \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/2(1-\tau)} \right]^{(1-\tau)} \\ & \leq 1 + C^q q^q \mathbb{E} \left[\left(\frac{r_{*,T,\xi}}{\varepsilon} \right)^{(d-\delta)q/2(1-\tau)} \right]^{(1-\tau)}, \end{aligned}$$

where $C = C(d, m, \lambda, \Lambda, \rho, \delta, \tau)$ and where $q \geq C(d, m, \lambda, \Lambda, \delta)$. Now, fixing $\delta > 0$ small enough (depending only on d, m, λ , and Λ) and choosing $\tau \in (0, 1)$ such that $1 - \tau = (d - \delta)/(d - \delta/2)$, the estimate buckles and yields a bound of the form

$$\mathbb{E} \left[\left(\frac{r_{*,T,\xi}(x_0)}{\varepsilon} \right)^{(d-\delta/2)q/2} \right]^\tau \leq C(d, m, \lambda, \Lambda, \rho)^q q^q$$

for all $q \geq C(d, m, \lambda, \Lambda)$. This may be rewritten as

$$\mathbb{E} \left[\left(\frac{r_{*,T,\xi}(x_0)}{\varepsilon} \right)^{(d-\delta/2)q/2} \right]^{1/q} \leq C(d, m, \lambda, \Lambda, \rho) q^C,$$

which establishes Lemma 19. \square

We now derive the estimates on the corrector ϕ_ξ^T , the flux corrector σ_ξ^T , and the potential field θ_ξ^T . Note that the only required ingredients for the proof are the estimate for functionals of the corrector which we derived in Lemma 17, the estimate on the corrector gradient given by (88), as well as the general technical results of Lemma 40 and Lemma 25.

Proof of Proposition 13. We insert the estimate on $r_{*,T,\xi}$ from Lemma 19 into Lemma 17. For any $x_0 \in \mathbb{R}^d$, any $r \geq \varepsilon$, and any g with $\text{supp } g \subset B_r(x_0)$ and

$$\left(\int_{B_r(x_0)} |g|^p dx \right)^{1/p} \leq r^{-d},$$

this provides bounds of the form

$$(108) \quad \mathbb{E} \left[\left| \int_{\mathbb{R}^d} g \cdot \nabla \phi_\xi^T dx \right|^q \right]^{1/q} \leq C|\xi|q^{C(d,m,\lambda,\Lambda)} \left(\frac{\varepsilon}{r} \right)^{d/2}$$

and

$$(109) \quad \mathbb{E} \left[\left| \int_{\mathbb{R}^d} g \cdot \nabla \sigma_{\xi,jk}^T dx \right|^q \right]^{1/q} \leq C|\xi|q^{C(d,m,\lambda,\Lambda)} \left(\frac{\varepsilon}{r} \right)^{d/2}$$

as well as

$$(110) \quad \mathbb{E} \left[\left| \int_{\mathbb{R}^d} g \cdot \nabla \theta_\xi^T dx \right|^q \right]^{1/q} \leq Cr|\xi|q^{C(d,m,\lambda,\Lambda)} \left(\frac{\varepsilon}{r} \right)^{d/2}$$

for any $q \geq C(d, m, \lambda, \Lambda)$.

Plugging in the estimates on $r_{*,T,\xi}$ from Lemma 19 into (88), we deduce for any $x_0 \in \mathbb{R}^d$

$$(111) \quad \mathbb{E} \left[\left| \int_{B_\varepsilon(x_0)} |\nabla \phi_\xi^T|^2 dx \right|^{q/2} \right]^{1/q} \leq C|\xi|q^{C(d,m,\lambda,\Lambda)}$$

for any $q \geq C(d, m, \lambda, \Lambda)$. Plugging in this bound and (108) into the (spatially rescaled) multiscale estimate for the L^2 norm from Lemma 25, we obtain the bounds on ϕ_ξ^T stated in (21) as well as the corresponding result for the L^p norm.

The estimates on ϕ_ξ^T in (22a) and (22b) are shown by combining (21) with the estimate from Lemma 18 and (111) as well as using in case $d \geq 3$ the relation

$$\lim_{R \rightarrow \infty} \int_{B_R(x_0)} \phi_\xi^T dx = 0.$$

In view of this use of Lemma 25, in order to obtain the estimates on $\sigma_{\xi,jk}^T$ and θ_ξ^T stated in (21) and (23), we only need to establish estimates on $\int_{B_\varepsilon(x_0)} |\nabla \sigma_{\xi ij}^T|^2 dx$ and $\int_{B_\varepsilon(x_0)} |\nabla \theta_\xi^T|^2 dx$. The Caccioppoli inequality (51) for the defining equation of the flux corrector (12c) yields in conjunction with (A2)

$$\begin{aligned} \int_{x_0+[-\varepsilon,\varepsilon]^d} |\nabla \sigma_{\xi,jk}^T|^2 dx &\leq \frac{C}{\varepsilon^2} \int_{x_0+[-2\varepsilon,2\varepsilon]^d} \left| \sigma_{\xi,jk}^T - \int_{x_0+[-2\varepsilon,2\varepsilon]^d} \sigma_{\xi,jk}^T d\tilde{x} \right|^2 dx \\ &\quad + C \int_{x_0+[-2\varepsilon,2\varepsilon]^d} |\xi + \nabla \phi_\xi|^2 dx \\ &\quad + \frac{C}{T} \left| \int_{x_0+[-2\varepsilon,2\varepsilon]^d} \sigma_{\xi,jk}^T d\tilde{x} \right|^2. \end{aligned}$$

Estimating the first term on the right-hand side by Lemma 40 and taking stochastic moments, we deduce (where we use the family of functions g_n of the form $\varepsilon^{-d} \prod_{i=1}^d \cos(\pi k_i(x_i + 2\varepsilon)/4\varepsilon)$, $k \in \{-K, \dots, K\}^d$)

$$\begin{aligned} &\mathbb{E} \left[\left| \int_{x_0+[-2\varepsilon,2\varepsilon]^d} |\nabla \sigma_{\xi,jk}^T|^2 dx \right|^q \right]^{1/q} \\ &\leq \frac{C}{K^2} \mathbb{E} \left[\left| \int_{x_0+[-2\varepsilon,2\varepsilon]^d} |\nabla \sigma_{\xi,jk}^T|^2 dx \right|^q \right]^{1/q} \\ &\quad + \sum_{n=1}^{N(d,K)} C \mathbb{E} \left[\left| \int_{\mathbb{R}^d} g_n \cdot \nabla \sigma_{\xi,jk}^T dx \right|^q \right]^{1/q} \\ &\quad + C \mathbb{E} \left[\left| \int_{x_0+[-2\varepsilon,2\varepsilon]^d} |\xi + \nabla \phi_\xi^T|^2 dx \right|^q \right]^{1/q} \\ &\quad + \frac{C}{T} \mathbb{E} \left[\left| \int_{x_0+[-2\varepsilon,2\varepsilon]^d} \sigma_{\xi,jk}^T d\tilde{x} \right|^{2q} \right]^{1/q}. \end{aligned}$$

Choosing K large enough and using stationarity, we may absorb the first term on the right-hand side in the left-hand side. Estimating the linear functionals of $\nabla \sigma_{\xi,jk}^T$ by (109), bounding the third term on the right-hand side by (111), and estimating the last term by (37) and (38) (where we may replace the average over the box $x_0 + [-2\varepsilon, 2\varepsilon]^d$ by the average over the ball $B_\varepsilon(x_0)$ using (109)), we deduce

$$\mathbb{E} \left[\left| \int_{x_0+[-\varepsilon,\varepsilon]^d} |\nabla \sigma_{\xi,jk}^T|^2 dx \right|^{q\gamma} \right]^{1/2q} \leq C |\xi| q^C.$$

Together with (109) and Lemma 25, we deduce the bound for $\sigma_{\xi,jk}^T$ in (21).

The estimates on σ_ξ^T in (22a) and (22b) are again shown by combining (21) with the estimate from Lemma 18 and (111) as well as using in case $d \geq 3$ the relation

$$\lim_{R \rightarrow \infty} \int_{B_R(x_0)} \sigma_\xi^T dx = 0.$$

The estimate for the gradient of the potential field $\nabla \theta_\xi^T$ is analogous but even simpler (due to the lack of the massive regularization in (19a)). We obtain the

bound

$$\mathbb{E} \left[\left| \int_{x_0 + [-\varepsilon, \varepsilon]^d} |\nabla \theta_\xi^T|^2 dx \right|^q \right]^{1/2q} \leq C q^C \varepsilon |\xi|.$$

Using this estimate and (109) in Lemma 25, we obtain (23). \square

6. CORRECTOR ESTIMATES FOR THE LINEARIZED SYSTEM

6.1. Estimates on linear functionals of the corrector and the flux corrector for the linearized PDE.

Proof of Lemma 21 and Lemma 22. Part a: Estimates for linear functionals of the homogenization corrector $\phi_{\xi, \Xi}^T$. Without loss of generality we may assume in the following argument that $x_0 = 0$, i. e. g is supported in $B_r(0)$, and that $(\int_{B_r} |g|^{p^2/2} dx)^{2/p^2} \leq 1$ – otherwise replace in the following argument p by $\sqrt{2p}$.

The argument is similar to the case of the corrector ϕ_ξ^T . We first observe that the expectation $\mathbb{E}[F]$ vanishes. Indeed, $\phi_{\xi, \Xi}^T$ is easily seen to be a stationary random field, which entails $\mathbb{E}[F] = \int_{\mathbb{R}^d} g \cdot \nabla \mathbb{E}[\phi_\xi^T] dx = 0$. By Lemma 16, to obtain stochastic moment bounds for F it suffices to estimate the sensitivity of F with respect to changes in the random field ω_ε . Taking the derivative with respect to ω_ε in (18a), we obtain

$$\begin{aligned} & -\nabla \cdot (\partial_\xi A(\omega_\varepsilon(x), \xi + \nabla \phi_\xi^T) \nabla \delta \phi_{\xi, \Xi}^T) + \frac{1}{T} \delta \phi_{\xi, \Xi}^T \\ (112) \quad & = \nabla \cdot (\partial_\omega \partial_\xi A(\omega_\varepsilon(x), \xi + \nabla \phi_\xi^T) \delta \omega_\varepsilon(\Xi + \nabla \phi_{\xi, \Xi}^T)) \\ & + \nabla \cdot (\partial_\xi^2 A(\omega_\varepsilon(x), \xi + \nabla \phi_\xi^T) (\Xi + \nabla \phi_{\xi, \Xi}^T) \nabla \delta \phi_\xi^T). \end{aligned}$$

Denoting by h the unique solution in $H^1(\mathbb{R}^d; \mathbb{R}^m)$ to the auxiliary PDE

$$(113) \quad -\nabla \cdot (a_\xi^{T,*} \nabla h) + \frac{1}{T} h = \nabla \cdot g$$

(where we again used the abbreviation $a_\xi^{T,*} := \partial_\xi A(\omega_\varepsilon(x), \xi + \nabla \phi_\xi^T)$) and denoting by $\hat{h} \in H^1(\mathbb{R}^d; \mathbb{R}^m)$ the unique solution to the auxiliary PDE

$$\begin{aligned} (114) \quad & -\nabla \cdot (a_\xi^{T,*} \nabla \hat{h}) + \frac{1}{T} \hat{h} \\ & = \sum_{j=1}^d \sum_{l=1}^m \partial_j (\partial_\xi^2 A(\omega_\varepsilon(x), \xi + \nabla \phi_\xi^T) (\Xi + \nabla \phi_{\xi, \Xi}^T) (e_l \otimes e_j) \cdot \nabla h) e_l, \end{aligned}$$

we deduce

$$\begin{aligned}
\delta F &= \int_{\mathbb{R}^d} g \cdot \nabla \delta \phi_{\xi, \Xi}^T dx \stackrel{(113)}{=} - \int_{\mathbb{R}^d} a_{\xi}^T \nabla \delta \phi_{\xi, \Xi}^T \cdot \nabla h + \frac{1}{T} \delta \phi_{\xi, \Xi}^T h dx \\
&\stackrel{(112)}{=} \int_{\mathbb{R}^d} \partial_{\omega} \partial_{\xi} A(\omega_{\varepsilon}(x), \xi + \nabla \phi_{\xi}^T) \delta \omega_{\varepsilon}(\Xi + \nabla \phi_{\xi, \Xi}^T) \cdot \nabla h dx \\
&\quad + \int_{\mathbb{R}^d} \partial_{\xi}^2 A(\omega_{\varepsilon}(x), \xi + \nabla \phi_{\xi}^T) (\Xi + \nabla \phi_{\xi, \Xi}^T) \nabla \delta \phi_{\xi}^T \cdot \nabla h dx \\
&\stackrel{(114)}{=} \int_{\mathbb{R}^d} \partial_{\omega} \partial_{\xi} A(\omega_{\varepsilon}(x), \xi + \nabla \phi_{\xi}^T) \delta \omega_{\varepsilon}(\Xi + \nabla \phi_{\xi, \Xi}^T) \cdot \nabla h dx \\
&\quad - \int_{\mathbb{R}^d} a_{\xi}^{T,*} \nabla \hat{h} \cdot \nabla \delta \phi_{\xi}^T + \frac{1}{T} \hat{h} \phi_{\xi}^T dx \\
&\stackrel{(92)}{=} \int_{\mathbb{R}^d} \partial_{\omega} \partial_{\xi} A(\omega_{\varepsilon}(x), \xi + \nabla \phi_{\xi}^T) \delta \omega_{\varepsilon}(\Xi + \nabla \phi_{\xi, \Xi}^T) \cdot \nabla h dx \\
&\quad + \int_{\mathbb{R}^d} \partial_{\omega} A(\omega_{\varepsilon}(x), \xi + \nabla \phi_{\xi}^T) \delta \omega_{\varepsilon} \cdot \nabla \hat{h} dx.
\end{aligned}$$

In other words, we have the representation

$$\frac{\partial F}{\partial \omega_{\varepsilon}} = \partial_{\omega} \partial_{\xi} A(\omega_{\varepsilon}(x), \xi + \nabla \phi_{\xi}^T) (\Xi + \nabla \phi_{\xi, \Xi}^T) \cdot \nabla h + \partial_{\omega} A(\omega_{\varepsilon}(x), \xi + \nabla \phi_{\xi}^T) \cdot \nabla \hat{h}.$$

By (A3), this implies the sensitivity estimate

$$\begin{aligned}
&\int_{\mathbb{R}^d} \left| \int_{B_{\varepsilon}(x)} \left| \frac{\partial F}{\partial \omega_{\varepsilon}} \right| d\tilde{x} \right|^2 dx \\
&\leq C \int_{\mathbb{R}^d} \left| \int_{B_{\varepsilon}(x)} |\Xi + \nabla \phi_{\xi, \Xi}^T| |\nabla h| d\tilde{x} \right|^2 dx + C \int_{\mathbb{R}^d} \left| \int_{B_{\varepsilon}(x)} |\xi + \nabla \phi_{\xi}^T| |\nabla \hat{h}| d\tilde{x} \right|^2 dx.
\end{aligned}$$

Plugging this bound into the version of the spectral gap inequality for the q -th moment (see Lemma 16) and using $\mathbb{E}[F] = 0$, we deduce for any $q \geq 1$

$$\begin{aligned}
\mathbb{E}[|F|^{2q}]^{1/2q} &\leq Cq\varepsilon^{d/2} \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \left| \int_{B_{\varepsilon}(x)} |\Xi + \nabla \phi_{\xi, \Xi}^T| |\nabla h| d\tilde{x} \right|^2 dx \right)^q \right]^{1/2q} \\
&\quad + Cq\varepsilon^{d/2} \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \left| \int_{B_{\varepsilon}(x)} |\xi + \nabla \phi_{\xi}^T| |\nabla \hat{h}| d\tilde{x} \right|^2 dx \right)^q \right]^{1/2q}.
\end{aligned}$$

By Lemma 39 and (89) as well as (88), this entails for any $\tau, \bar{\tau} \in (0, 1)$

$$\begin{aligned}
(115) \quad &\mathbb{E}[|F|^{2q}]^{1/2q} \\
&\leq C|\Xi|q\varepsilon^{d/2}r^{d/2}\mathbb{E}\left[\left(\frac{r_{*,T,\xi,\Xi}}{\varepsilon}\right)^{(d-\delta)q/(1-\tau)}\right]^{(1-\tau)/2q} \\
&\quad \times \mathbb{E}\left[\left(r^{-d}\int_{\mathbb{R}^d}|\nabla h|^p\left(1+\frac{|x|}{r}\right)^{\alpha_0}dx\right)^{2q/p\tau}\right]^{\tau/2q} \\
&\quad + C|\xi|q\varepsilon^{d/2}r^{d/2}\mathbb{E}\left[\left(\frac{r_{*,T,\xi}}{\varepsilon}\right)^{(d-\delta)q/(1-\bar{\tau})}\right]^{(1-\bar{\tau})/2q}
\end{aligned}$$

$$\times \mathbb{E} \left[\left(r^{-d} \int_{\mathbb{R}^d} |\nabla \hat{h}|^p \left(1 + \frac{|x|}{r} \right)^{\alpha_0} dx \right)^{2q/p\bar{\tau}} \right]^{\bar{\tau}/2q}$$

with $C = C(d, m, \lambda, \Lambda, \rho, \alpha_0, p, \tau, \bar{\tau})$. By the weighted Meyers' estimate in Lemma 35 – applied to (114) – and the uniform bound $|\partial_\xi^2 A| \leq \Lambda$ from (R), we infer for any $\alpha_{1/2} > \alpha_0$

$$(116) \quad \mathbb{E} \left[\left(r^{-d} \int_{\mathbb{R}^d} |\nabla \hat{h}|^p \left(1 + \frac{|x|}{r} \right)^{\alpha_0} dx \right)^{2q/p\bar{\tau}} \right]^{\bar{\tau}/2q} \\ \leq C \mathbb{E} \left[\left(r^{-d} \int_{\mathbb{R}^d} |\nabla h|^p |\Xi + \nabla \phi_{\xi, \Xi}^T|^p \left(1 + \frac{|x|}{r} \right)^{\alpha_{1/2}} dx \right)^{2q/p\bar{\tau}} \right]^{\bar{\tau}/2q}$$

with $C = C(d, m, \lambda, \Lambda, \rho, \alpha_0, \alpha_{1/2}, p, \tau, \bar{\tau})$. Inserting the bound (42) into (116), we get

$$\mathbb{E} \left[\left(r^{-d} \int_{\mathbb{R}^d} |\nabla \hat{h}|^p \left(1 + \frac{|x|}{r} \right)^{\alpha_0} dx \right)^{2q/p\bar{\tau}} \right]^{\bar{\tau}/2q} \\ \leq C q^C (1 + |\xi|)^C |\Xi| \\ \times \mathbb{E} \left[\left(\int_{\mathbb{R}^d} r^{-d} \mathcal{C}_{reg, \xi}^p(x) \left(\frac{r_{*, T, \xi, \Xi}(x)}{\varepsilon} \right)^{(d-\delta)p/2} |\nabla h|^p \left(1 + \frac{|x|}{r} \right)^{\alpha_{1/2}} dx \right)^{2q/p\bar{\tau}} \right]^{\bar{\tau}/2q}$$

where C now additionally depends on ν . To proceed further, we write the second factor on the right-hand side in the form

$$\mathbb{E} \left[\left(\int_{\mathbb{R}^d} v_1 v_2 dx \right)^{2q/p\bar{\tau}} \right]^{\bar{\tau}/2q}$$

with

$$v_1 = \mathcal{C}_{reg, \xi}^p \left(\frac{r_{*, T, \xi, \Xi}}{\varepsilon} \right)^{(d-\delta)p/2}, \quad v_2 = r^{-d} |\nabla h|^p \left(1 + \frac{|x|}{r} \right)^{\alpha_{1/2}}.$$

By first smuggling in the weight

$$\varphi(x) = r^{-d} \left(1 + \frac{|x|}{r} \right)^{-(d+1)},$$

via Hölder's inequality with exponents $\frac{p}{p-2}$ and $\frac{p}{2}$ in space, and next by Hölder's inequality with exponents $\frac{\bar{\tau}}{\bar{\tau}-\tau/2}$ and $\frac{2\bar{\tau}}{\tau}$ in probability, we get

$$\mathbb{E} \left[\left(\int_{\mathbb{R}^d} v_1 v_2 dx \right)^{2q/p\bar{\tau}} \right]^{\bar{\tau}/2q} \\ \leq \mathbb{E} \left[\left(\int_{\mathbb{R}^d} v_1^{\frac{p}{p-2}} \varphi dx \right)^{\frac{p-2}{p} \frac{2q}{p(\bar{\tau}-\tau/2)}} \right]^{\frac{\bar{\tau}-\tau/2}{2q}} \mathbb{E} \left[\left(\int_{\mathbb{R}^d} v_2^{\frac{p}{2}} \varphi^{-\frac{p-2}{2}} dx \right)^{\frac{8q}{p^2\bar{\tau}}} \right]^{\frac{\tau}{4q}}.$$

By Jensen's inequality for the integral $\int_{\mathbb{R}^d} f \varphi dx$ – using that φ has mass of order unity and assuming also that $q \geq C(p, \tau, \bar{\tau})$ – and by exploiting the fact that v_1

is a stationary random field, we deduce that the right-hand side is bounded from above by

$$C\mathbb{E}\left[v_1^{\frac{2q}{p(\bar{\tau}-\tau/2)}}\right]^{\frac{\bar{\tau}-\tau/2}{2q}}\mathbb{E}\left[\left(\int_{\mathbb{R}^d}v_2^{\frac{p}{2}}\varphi^{-\frac{p-2}{2}}dx\right)^{\frac{8q}{p^2\tau}}\right]^{\frac{\tau}{4q}}.$$

By combining the previous estimates and plugging in the definitions of v_1, v_2 and φ , we thus arrive at

$$\begin{aligned} & \mathbb{E}\left[\left(r^{-d}\int_{\mathbb{R}^d}|\nabla\hat{h}|^p\left(1+\frac{|x|}{r}\right)^{\alpha_0}dx\right)^{2q/p\bar{\tau}}\right]^{\bar{\tau}/2q} \\ & \leq Cq^C(1+|\xi|)^C|\Xi|\mathbb{E}\left[|\mathcal{C}_{reg,\xi}|^{2q/(\bar{\tau}-\tau/2)}\left(\frac{r_{*,T,\xi,\Xi}}{\varepsilon}\right)^{(d-\delta)q/(\bar{\tau}-\tau/2)}\right]^{(\bar{\tau}-\tau/2)/2q} \\ & \quad \times \mathbb{E}\left[\left(\int_{\mathbb{R}^d}r^{-d}|\nabla h|^{p^2/2}\left(1+\frac{|x|}{r}\right)^{\alpha_{1/2}p/2+(d+1)(p-2)/2}dx\right)^{8q/p^2\tau}\right]^{\tau/4q}. \end{aligned}$$

Applying once more Hölder's inequality to the first expected value on the right-hand side, choosing $\bar{\tau} \in (0, 1)$ close enough to 1, choosing $\tau > 0$ small enough, and using the stretched exponential moment bounds for $\mathcal{C}_{reg,\xi}$, we deduce

$$\begin{aligned} (117) \quad & \mathbb{E}\left[\left(r^{-d}\int_{\mathbb{R}^d}|\nabla\hat{h}|^p\left(1+\frac{|x|}{r}\right)^{\alpha_0}dx\right)^{2q/p\bar{\tau}}\right]^{\bar{\tau}/2q} \\ & \leq Cq^C(1+|\xi|)^C|\Xi|\mathbb{E}\left[\left(\frac{r_{*,T,\xi,\Xi}}{\varepsilon}\right)^{(d-\delta)q/(1-\tau)}\right]^{(1-\tau)/2q} \\ & \quad \times \mathbb{E}\left[\left(\int_{\mathbb{R}^d}r^{-d}|\nabla h|^{p^2/2}\left(1+\frac{|x|}{r}\right)^{\alpha_{1/2}p/2+(d+1)(p-2)/2}dx\right)^{8q/p^2\tau}\right]^{\tau/4q}. \end{aligned}$$

Plugging in the resulting estimate into (115) and estimating $r_{*,T,\xi}$ in (115) via Lemma 19 yields

$$\begin{aligned} & \mathbb{E}\left[|F|^{2q}\right]^{1/2q} \\ & \leq Cq^C(1+|\xi|)^C|\Xi|\varepsilon^{d/2}r^{d/2}\mathbb{E}\left[\left(\frac{r_{*,T,\xi,\Xi}}{\varepsilon}\right)^{(d-\delta)q/(1-\tau)}\right]^{(1-\tau)/2q} \\ & \quad \times \mathbb{E}\left[\left(\int_{\mathbb{R}^d}r^{-d}|\nabla h|^{p^2/2}\left(1+\frac{|x|}{r}\right)^{\alpha_{1/2}p/2+(d+1)(p-2)/2}dx\right)^{8q/p^2\tau}\right]^{\tau/4q}. \end{aligned}$$

Choosing p close enough to 2 and choosing α_0 and α_1 small enough (all depending only on d, m, λ , and Λ), we may estimate the last factor by applying the weighted Meyers' estimate from Lemma 35 to the PDE (113). This yields by our assumed bound on g

$$(118) \quad \mathbb{E}\left[|F|^{2q}\right]^{1/2q} \leq Cq^C(1+|\xi|)^C|\Xi|\left(\frac{\varepsilon}{r}\right)^{d/2}\mathbb{E}\left[\left(\frac{r_{*,T,\xi,\Xi}}{\varepsilon}\right)^{(d-\delta)q/(1-\tau)}\right]^{(1-\tau)/2q}$$

for any $q \geq C$ with $C = C(d, m, \lambda, \Lambda, \rho, \nu, \tau)$ and any $\tau \leq c(d, m, \lambda, \Lambda)$.

Part b: Estimates for linear functionals of the flux corrector $\sigma_{\xi,\Xi}^T$. Differentiating the equation (18c), we see that the infinitesimal perturbation $\delta\sigma_{\xi,\Xi}^T$ caused by an infinitesimal perturbation $\delta\omega_\varepsilon$ in the random field ω_ε satisfies the PDE

$$(119) \quad \begin{aligned} & -\Delta\delta\sigma_{\xi,\Xi,jk}^T + \frac{1}{T}\sigma_{\xi,\Xi,jk}^T \\ & = \nabla \cdot (\partial_\xi A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T)\nabla\delta\phi_{\xi,\Xi}^T \cdot (e_k \otimes e_j - e_j \otimes e_k)) \\ & \quad + \nabla \cdot (\partial_\xi^2 A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T)(\Xi + \nabla\phi_{\xi,\Xi}^T)\nabla\delta\phi_\xi^T \cdot (e_k \otimes e_j - e_j \otimes e_k)) \\ & \quad + \nabla \cdot (\partial_\omega\partial_\xi A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T)\delta\omega_\varepsilon(\Xi + \nabla\phi_{\xi,\Xi}^T) \cdot (e_k \otimes e_j - e_j \otimes e_k)). \end{aligned}$$

Introducing the solution $\bar{h} \in H^1(\mathbb{R}^d; \mathbb{R}^m)$ to the equation

$$(120) \quad -\Delta\bar{h} + \frac{1}{T}\bar{h} = \nabla \cdot g,$$

the solution $h_2 \in H^1(\mathbb{R}^d; \mathbb{R}^m)$ to the equation

$$(121) \quad \begin{aligned} & -\nabla \cdot (a_\xi^{T,*}\nabla h_2) + \frac{1}{T}h_2 \\ & = \sum_{i=1}^d \sum_{l=1}^m \partial_i(\partial_\xi A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T)(e_l \otimes e_i) \cdot (e_k \otimes e_j - e_j \otimes e_k) \cdot \nabla\bar{h})e_l \end{aligned}$$

(where $a_\xi^T(x) := \partial_\xi A(\omega_\varepsilon(x), \xi + \nabla\phi_\xi^T(x))$), and the solution $h_3 \in H^1(\mathbb{R}^d; \mathbb{R}^m)$ to the equation

$$(122) \quad \begin{aligned} & -\nabla \cdot (a_\xi^{T,*}\nabla h_3) + \frac{1}{T}h_3 \\ & = \sum_{i=1}^d \sum_{l=1}^m \partial_i(\partial_\xi^2 A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T)(\Xi + \nabla\phi_{\xi,\Xi}^T)(e_l \otimes e_i) \cdot (e_k \otimes e_j - e_j \otimes e_k) \cdot \nabla\bar{h})e_l \\ & \quad + \sum_{i=1}^d \sum_{l=1}^m \partial_i(\partial_\xi^2 A(\omega_\varepsilon(x), \xi + \nabla\phi_\xi^T)(\Xi + \nabla\phi_{\xi,\Xi}^T)(e_l \otimes e_i) \cdot \nabla h_2)e_l, \end{aligned}$$

we may compute the sensitivity of linear functionals of the form F_σ : Indeed, we have

$$\begin{aligned} \delta F_\sigma & = \int_{\mathbb{R}^d} g \cdot \nabla\delta\sigma_{\xi,\Xi,jk}^T dx \stackrel{(120)}{=} - \int_{\mathbb{R}^d} \nabla\bar{h} \cdot \nabla\delta\sigma_{\xi,\Xi,jk}^T dx - \frac{1}{T} \int_{\mathbb{R}^d} \bar{h} \delta\sigma_{\xi,\Xi,jk}^T dx \\ & \stackrel{(119)}{=} \int_{\mathbb{R}^d} \partial_\xi A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T)\nabla\delta\phi_{\xi,\Xi}^T \cdot (e_k \otimes e_j - e_j \otimes e_k) \cdot \nabla\bar{h} dx \\ & \quad + \int_{\mathbb{R}^d} \partial_\xi^2 A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T)(\Xi + \nabla\phi_{\xi,\Xi}^T)\nabla\delta\phi_\xi^T \cdot (e_k \otimes e_j - e_j \otimes e_k) \cdot \nabla\bar{h} dx \\ & \quad + \int_{\mathbb{R}^d} \partial_\omega\partial_\xi A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T)\delta\omega_\varepsilon(\Xi + \nabla\phi_{\xi,\Xi}^T) \cdot (e_k \otimes e_j - e_j \otimes e_k) \cdot \nabla\bar{h} dx \\ & \stackrel{(121)}{=} - \int_{\mathbb{R}^d} \partial_\xi A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T)\nabla\delta\phi_{\xi,\Xi}^T \cdot \nabla h_2 + \frac{1}{T}\delta\phi_{\xi,\Xi}^T h_2 dx \\ & \quad + \int_{\mathbb{R}^d} \partial_\xi^2 A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T)(\Xi + \nabla\phi_{\xi,\Xi}^T)\nabla\delta\phi_\xi^T \cdot (e_k \otimes e_j - e_j \otimes e_k) \cdot \nabla\bar{h} dx \\ & \quad + \int_{\mathbb{R}^d} \partial_\omega\partial_\xi A(\omega_\varepsilon, \xi + \nabla\phi_\xi^T)\delta\omega_\varepsilon(\Xi + \nabla\phi_{\xi,\Xi}^T) \cdot (e_k \otimes e_j - e_j \otimes e_k) \cdot \nabla\bar{h} dx \end{aligned}$$

$$\begin{aligned}
&\stackrel{(112)}{=} \int_{\mathbb{R}^d} \partial_\omega \partial_\xi A(\omega_\varepsilon(x), \xi + \nabla \phi_\xi^T) \delta \omega_\varepsilon (\Xi + \nabla \phi_{\xi, \Xi}^T) \cdot \nabla h_2 \, dx \\
&\quad + \int_{\mathbb{R}^d} \partial_\xi^2 A(\omega_\varepsilon(x), \xi + \nabla \phi_\xi^T) (\Xi + \nabla \phi_{\xi, \Xi}^T) \nabla \delta \phi_\xi^T \cdot \nabla h_2 \, dx \\
&\quad + \int_{\mathbb{R}^d} \partial_\xi^2 A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) (\Xi + \nabla \phi_{\xi, \Xi}^T) \nabla \delta \phi_\xi^T \cdot (e_k \otimes e_j - e_j \otimes e_k) \cdot \nabla \bar{h} \, dx \\
&\quad + \int_{\mathbb{R}^d} \partial_\omega \partial_\xi A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \delta \omega_\varepsilon (\Xi + \nabla \phi_{\xi, \Xi}^T) \cdot (e_k \otimes e_j - e_j \otimes e_k) \cdot \nabla \bar{h} \, dx.
\end{aligned}$$

Inserting (122) and (92), we obtain

$$\begin{aligned}
\delta F_\sigma &= \int_{\mathbb{R}^d} \partial_\omega \partial_\xi A(\omega_\varepsilon(x), \xi + \nabla \phi_\xi^T) \delta \omega_\varepsilon (\Xi + \nabla \phi_{\xi, \Xi}^T) \cdot \nabla h_2 \, dx \\
&\quad + \int_{\mathbb{R}^d} \partial_\omega A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \delta \omega_\varepsilon \cdot \nabla h_3 \, dx \\
&\quad + \int_{\mathbb{R}^d} \partial_\omega \partial_\xi A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \delta \omega_\varepsilon (\Xi + \nabla \phi_{\xi, \Xi}^T) \cdot (e_k \otimes e_j - e_j \otimes e_k) \cdot \nabla \bar{h} \, dx.
\end{aligned}$$

This identification of $\frac{\partial F}{\partial \omega_\varepsilon}$ together with assumption (A3) gives rise to the sensitivity estimate

$$\begin{aligned}
&\mathbb{E} \left[\left(\int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} \left| \frac{\partial F_\sigma}{\partial \omega_\varepsilon} \right| d\tilde{x} \right|^2 dx \right)^q \right]^{1/2q} \\
&\leq C \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} |\Xi + \nabla \phi_{\xi, \Xi}^T| (|\nabla \bar{h}| + |\nabla h_2|) d\tilde{x} \right|^2 dx \right)^q \right]^{1/2q} \\
&\quad + C \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T| |\nabla h_3| d\tilde{x} \right|^2 dx \right)^q \right]^{1/2q}.
\end{aligned}$$

Plugging in this bound into Lemma 16 and using $\mathbb{E}[F_\sigma] = \int_{\mathbb{R}^d} g \cdot \nabla \mathbb{E}[\sigma_{\xi, \Xi, jk}^T] \, dx = 0$ by stationarity of $\sigma_{\xi, \Xi, jk}^T$, we infer

$$\begin{aligned}
\mathbb{E} \left[|F_\sigma|^{2q} \right]^{1/2q} &\leq C q \varepsilon^{d/2} \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} |\Xi + \nabla \phi_{\xi, \Xi}^T| (|\nabla \bar{h}| + |\nabla h_2|) d\tilde{x} \right|^2 dx \right)^q \right]^{1/2q} \\
&\quad + C q \varepsilon^{d/2} \mathbb{E} \left[\left(\int_{\mathbb{R}^d} \left| \int_{B_\varepsilon(x)} |\xi + \nabla \phi_\xi^T| |\nabla h_3| d\tilde{x} \right|^2 dx \right)^q \right]^{1/2q}.
\end{aligned}$$

By Lemma 39 and (89) as well as (88), we obtain for $\tau, \bar{\tau} \in (0, 1)$ and for p close enough to 2

$$\begin{aligned}
&\mathbb{E} \left[|F_\sigma|^{2q} \right]^{1/2q} \\
&\leq C |\Xi| q \varepsilon^{d/2} r^{d/2} \mathbb{E} \left[\left(\frac{r_{*, T, \xi, \Xi}}{\varepsilon} \right)^{(d-\delta)q/(1-\tau)} \right]^{(1-\tau)/2q} \\
&\quad \times \mathbb{E} \left[\left(r^{-d} \int_{\mathbb{R}^d} (|\nabla \bar{h}|^p + |\nabla h_2|^p) \left(1 + \frac{|x|}{r} \right)^{\alpha_0} dx \right)^{2q/p\tau} \right]^{\tau/2q}
\end{aligned}$$

$$\begin{aligned}
& + C|\xi|q\varepsilon^{d/2}r^{d/2}\mathbb{E}\left[\left(\frac{r_{*,T,\xi}}{\varepsilon}\right)^{(d-\delta)q/(1-\bar{\tau})}\right]^{(1-\bar{\tau})/2q} \\
& \quad \times \mathbb{E}\left[\left(r^{-d}\int_{\mathbb{R}^d}|\nabla h_3|^p\left(1+\frac{|x|}{r}\right)^{\alpha_0}dx\right)^{2q/p\bar{\tau}}\right]^{\bar{\tau}/2q}.
\end{aligned}$$

By the weighted Meyers' estimate in Lemma 35 applied to (122) and the uniform bound on $\partial_\xi^2 A$ from (R), we infer for any $\alpha_{1/2} > \alpha_0$

$$\begin{aligned}
& \mathbb{E}\left[|F_\sigma|^{2q}\right]^{1/2q} \\
& \leq C|\Xi|q\varepsilon^{d/2}r^{d/2}\mathbb{E}\left[\left(\frac{r_{*,T,\xi,\Xi}}{\varepsilon}\right)^{(d-\delta)q/(1-\bar{\tau})}\right]^{(1-\bar{\tau})/2q} \\
& \quad \times \mathbb{E}\left[\left(r^{-d}\int_{\mathbb{R}^d}(|\nabla \bar{h}|^p + |\nabla h_2|^p)\left(1+\frac{|x|}{r}\right)^{\alpha_0}dx\right)^{2q/p\bar{\tau}}\right]^{\bar{\tau}/2q} \\
& \quad + C|\xi|q\varepsilon^{d/2}r^{d/2}\mathbb{E}\left[\left(\frac{r_{*,T,\xi}}{\varepsilon}\right)^{(d-\delta)q/(1-\bar{\tau})}\right]^{(1-\bar{\tau})/2q} \\
& \quad \times \mathbb{E}\left[\left(r^{-d}\int_{\mathbb{R}^d}(|\nabla \bar{h}|^p + |\nabla h_2|^p)|\Xi + \nabla \phi_{\xi,\Xi}^T|^p\left(1+\frac{|x|}{r}\right)^{\alpha_{1/2}}dx\right)^{2q/p\bar{\tau}}\right]^{\bar{\tau}/2q}.
\end{aligned}$$

Arguing analogously to the derivation of (118) from (115) but using also the estimate

$$\begin{aligned}
& \int_{\mathbb{R}^d} r^{-d}|\nabla h_2|^{p^2/2}\left(1+\frac{|x|}{r}\right)^{\alpha_{1/2}p/2+(d+1)(p-2)/2}dx \\
& \leq C \int_{\mathbb{R}^d} r^{-d}|\nabla \bar{h}|^{p^2/2}\left(1+\frac{|x|}{r}\right)^{\alpha_{3/4}}dx
\end{aligned}$$

for $\alpha_{3/4} > \alpha_{1/2}p/2+(d+1)(p-2)/2$ (which follows from (121), (A2), and Lemma 35), we deduce

$$\mathbb{E}\left[|F_\sigma|^{2q}\right]^{1/2q} \leq C(1+|\xi|)^C|\Xi|q^C\left(\frac{\varepsilon}{r}\right)^{d/2}\mathbb{E}\left[\left(\frac{r_{*,T,\xi,\Xi}}{\varepsilon}\right)^{(d-\delta)q/(1-\bar{\tau})}\right]^{(1-\bar{\tau})/2q}$$

for any $0 < \tau \leq c$ and any $q \geq C$.

Part c: Estimates for linear functionals of the potential field $\theta_{\xi,\Xi}^T$. The argument of reducing the estimate to the estimates in Part a is exactly the same as in part c of the proof of Lemma 17, as the relation between $\theta_{\xi,\Xi}^T$ and $\phi_{\xi,\Xi}^T$ is exactly the same as the one between θ_ξ^T and $\phi_{\xi,\Xi}^T$. As in the proof of Lemma 17, we introduce \bar{g} as the solution to the PDE

$$-\Delta \bar{g} = \nabla \cdot g.$$

We then obtain an estimate for F_θ of the form (115), but with h solving the PDE $-\nabla \cdot (a_\xi^{T,*} \nabla h) + \frac{1}{T}h = \nabla \cdot (\bar{g}e_i)$ and \hat{h} solving the same PDE but with the new h . Inserting the Calderon-Zygmund bounds on \bar{g} and the modified bound on $\nabla \hat{h}$ in the steps leading to (116) and (118), we deduce the desired estimate

$$\mathbb{E}\left[|F_\theta|^{2q}\right]^{1/2q} \leq Cr(1+|\xi|)^C|\Xi|q^C\left(\frac{\varepsilon}{r}\right)^{d/2}\mathbb{E}\left[\left(\frac{r_{*,T,\xi,\Xi}}{\varepsilon}\right)^{(d-\delta)q/(1-\bar{\tau})}\right]^{(1-\bar{\tau})/2q}$$

for any $0 < \tau \leq c$ and any $q \geq C$.

Part d: Proof of Lemma 22. The proof of the estimates (40) and (41) is analogous to the proof of (37) and (38). \square

6.2. Estimates on the corrector. We first establish the moment bounds on $r_{*,T,\xi,\Xi}$.

Proof of Lemma 23. The proof is entirely analogous to the proof of Lemma 19: Recall that the only required ingredients for the proof of moment bounds for $r_{*,T,\xi}$ in Lemma 19 were the estimate for functionals of the corrector ϕ_ξ^T from Lemma 17, the estimate on corrector averages from Lemma 18, the estimate on the corrector gradient given by (88), as well as the general technical results of Lemma 40 and Lemma 25. Lemma 21 provides bounds on the stochastic moments of linear functionals of the corrector $\phi_{\xi,\Xi}^T$ for the linearized PDE that are essentially analogous (up to the prefactor $q^C(1+|\xi|)^C$) to the bounds for functionals of ϕ_ξ^T in Lemma 17. Similarly, Lemma 22 provides estimates on averages of the linearized corrector $\phi_{\xi,\Xi}^T$ that are (again up to the prefactor $q^C(1+|\xi|)^C$) analogous to the bounds on averages of ϕ_ξ^T from Lemma 18. Furthermore, the estimate (89) for the gradient of $\phi_{\xi,\Xi}^T$ is completely analogous to the estimate (88) for the gradient of ϕ_ξ^T . In conclusion, by the same arguments as the proof of Lemma 19 (up to replacing ϕ_ξ^T by $\phi_{\xi,\Xi}^T$, ξ by Ξ , $r_{*,T,\xi}$ by $r_{*,T,\xi,\Xi}$, and including an additional prefactor $q^C(1+|\xi|)^C$ in the bounds), we obtain

$$\mathbb{E} \left[\left(\frac{r_{*,T,\xi}(x_0)}{\varepsilon} \right)^{(d-\delta/2)q/2} \right]^{1/q} \leq C(d, m, \lambda, \Lambda, \rho, \nu) q^C (1 + |\xi|)^C.$$

\square

Proof of Proposition 14. By the same arguments as in the proof of Proposition 13, we obtain the desired bounds on $\phi_{\xi,\Xi}^T$, $\sigma_{\xi,\Xi}^T$, and $\theta_{\xi,\Xi}^T$. Note that we simply need to replace the use of Lemma 17 by Lemma 21, the use of Lemma 19 by Lemma 23, and the use of (88) by (89). \square

Proof of Corollary 15. To establish the estimate (27), we simply pass to the limit $T \rightarrow \infty$ in the L^p version of the estimate (22a) for ϕ_ξ^T and σ_ξ^T . By combining (21) with Lemma 18 and Lemma 19, we deduce

$$\begin{aligned} & \left(\int_{B_r(x_0)} \left| \phi_\xi^T - \int_{B_\varepsilon(0)} \phi_\xi^T(\tilde{x}) d\tilde{x} \right|^2 + \left| \sigma_\xi^T - \int_{B_\varepsilon(0)} \sigma_\xi^T(\tilde{x}) d\tilde{x} \right|^2 dx \right)^{1/2} \\ & \leq C\varepsilon\mu(|x_0| + r)/\varepsilon \end{aligned}$$

with $\mu(s) := 1$ for $d \geq 3$, $\mu(s) := (\log(2+s))^{1/2}$ for $d = 2$, and $\mu(s) := (1+s)^{1/2}$ for $d = 1$. Passing to the limit $T \rightarrow \infty$, the quantities $\phi_\xi^T - \int_{B_\varepsilon(0)} \phi_\xi^T(\tilde{x}) d\tilde{x}$ converge to a solution ϕ_ξ to the corrector equation with vanishing average in $B_\varepsilon(0)$ subject to the bound (28) (and similarly for σ_ξ).

It remains to show that the limit σ_ξ satisfies the PDE (10), as passing to the limit in the equation (12c) yields only

$$-\Delta\sigma_{\xi,jk} = \partial_j(A(\omega_\varepsilon, \xi + \nabla\phi_\xi) \cdot e_k) - \partial_k((A(\omega_\varepsilon, \xi + \nabla\phi_\xi) \cdot e_j).$$

However, as we have $\nabla \cdot (A(\omega_\varepsilon, \xi + \nabla \phi_\xi)) = 0$ by the equation for ϕ_ξ , we deduce

$$-\Delta \sum_k \partial_k \sigma_{\xi, jk} = -\Delta (A(\omega_\varepsilon, \xi + \nabla \phi_\xi) \cdot e_j).$$

By the sublinear growth of $\sigma_{\xi, jk}$, it follows that

$$\nabla \cdot \sigma_\xi = A(\omega_\varepsilon, \xi + \nabla \phi_\xi) - \mathbb{E}[A(\omega_\varepsilon, \xi + \nabla \phi_\xi)] = A(\omega_\varepsilon, \xi + \nabla \phi_\xi) - A_{\text{hom}}(\xi),$$

which is the flux corrector equation (10).

In order to derive (29), we first write

$$\phi_{\xi_2}^T - \phi_{\xi_1}^T = \int_0^1 \phi_{(1-s)\xi_1 + s\xi_2, \Xi}^T ds$$

with $\Xi := \xi_2 - \xi_1$. This entails in case $d \geq 3$ using Proposition 14

$$\mathbb{E} \left[\left(\int_{B_r(x_0)} |\phi_{\xi_2}^T - \phi_{\xi_1}^T|^2 dx \right)^{q/2} \right]^{\frac{1}{q}} \leq Cq^C (1 + |\xi_1| + |\xi_2|)^C |\xi_2 - \xi_1| \varepsilon$$

respectively in case $d \leq 2$ using Proposition 14, Lemma 22, and Lemma 23

$$\begin{aligned} & \mathbb{E} \left[\left(\int_{B_r(x_0)} |\phi_{\xi_2}^T - \phi_{\xi_1}^T - \int_{B_\varepsilon(0)} \phi_{\xi_2}^T - \phi_{\xi_1}^T dx|^2 dx \right)^{q/2} \right]^{\frac{1}{q}} \\ & \leq Cq^C (1 + |\xi_1| + |\xi_2|)^C |\xi_2 - \xi_1| \varepsilon \mu((r + |x_0|)/\varepsilon). \end{aligned}$$

We may then pass to the limit $T \rightarrow \infty$ in these estimates to deduce (29). The corresponding bound for $\sigma_{\xi_2} - \sigma_{\xi_1}$ is derived analogously. \square

7. STRUCTURAL PROPERTIES OF THE EFFECTIVE EQUATION

We finally establish the structural properties of the effective (homogenized) equation, as stated in Theorem 6.

Proof of Theorem 6. Part a. The fact that the homogenized material law A_{hom} inherits the monotonicity properties of the law of the random material $A(\omega, \cdot)$ has already been established in [17, 18] for Euler-Lagrange equations associated with convex integral functionals. Nevertheless, we shall briefly recall the (standard) proof of this result. The result $A_{\text{hom}}(0) = 0$ is immediate from the definition $A_{\text{hom}}(\xi) := \mathbb{E}[A(\omega, \xi + \nabla \phi_\xi)]$ and (A2), as $\phi_0 \equiv 0$ is the (up to additive constants unique) solution to the corrector equation for $\xi = 0$. We next have

$$\begin{aligned} (123) \quad |A_{\text{hom}}(\xi_2) - A_{\text{hom}}(\xi_1)| &= |\mathbb{E}[A(\omega_\varepsilon, \xi_2 + \nabla \phi_{\xi_2})] - \mathbb{E}[A(\omega_\varepsilon, \xi_1 + \nabla \phi_{\xi_1})]| \\ &\stackrel{(A2)}{\leq} \Lambda \mathbb{E}[|\xi_2 + \nabla \phi_{\xi_2} - \xi_1 - \nabla \phi_{\xi_1}|]. \end{aligned}$$

Subtracting the corrector equations (4) for ξ_1 and ξ_2 from each other and testing the resulting equation with $(\phi_{\xi_2} - \phi_{\xi_1})\eta^2(x/r)$ for some cutoff η with $\text{supp } \eta \subset B_2$ and $\eta \equiv 1$ in B_1 , we deduce

$$\begin{aligned} & \int_{\mathbb{R}^d} (A(\omega_\varepsilon, \xi_2 + \nabla \phi_{\xi_2}) - A(\omega_\varepsilon, \xi_1 + \nabla \phi_{\xi_1})) \cdot (\nabla \phi_{\xi_2} - \nabla \phi_{\xi_1}) \eta^2 \left(\frac{x}{r} \right) dx \\ & \leq \int_{\mathbb{R}^d} |A(\omega_\varepsilon, \xi_2 + \nabla \phi_{\xi_2}) - A(\omega_\varepsilon, \xi_1 + \nabla \phi_{\xi_1})| \frac{1}{r} |\phi_{\xi_2} - \phi_{\xi_1}| |\nabla \eta^2| \left(\frac{x}{r} \right) dx. \end{aligned}$$

This implies by (A1), (A2), and Young's inequality

$$\begin{aligned} & \int_{\mathbb{R}^d} |\nabla\phi_{\xi_2} - \nabla\phi_{\xi_1}|^2 \eta^2\left(\frac{x}{r}\right) dx \\ & \leq \frac{4\Lambda^2}{\lambda^2} |\xi_1 - \xi_2|^2 \int_{\mathbb{R}^d} \eta^2\left(\frac{x}{r}\right) dx \\ & \quad + \frac{C}{r^2} \int_{\mathbb{R}^d} |\phi_{\xi_2} - \phi_{\xi_1}|^2 |\nabla\eta|^2\left(\frac{x}{r}\right) dx. \end{aligned}$$

Dividing by $\int_{\mathbb{R}^d} \eta^2(\frac{x}{r}) dx$, taking the expectation, using stationarity, and passing to the limit $r \rightarrow \infty$ (using the fact that the correctors ϕ_ξ grow sublinearly), we deduce $\mathbb{E}[|\nabla\phi_{\xi_1} - \nabla\phi_{\xi_2}|^2] \leq C(d) \frac{\Lambda^2}{\lambda^2} |\xi_1 - \xi_2|^2$. Inserting this into (123), we deduce the Lipschitz estimate in Theorem 6a.

Concerning the monotonicity property, we deduce by testing the corrector equations (4) for ξ_1 and ξ_2 with $(\phi_{\xi_1} - \phi_{\xi_2})\eta(x/r)$

$$\begin{aligned} & \int_{\mathbb{R}^d} \eta\left(\frac{x}{r}\right) (A(\omega_\varepsilon, \xi_2 + \nabla\phi_{\xi_2}) - A(\omega_\varepsilon, \xi_1 + \nabla\phi_{\xi_1})) \cdot (\xi_2 - \xi_1) dx \\ & = \int_{\mathbb{R}^d} \eta\left(\frac{x}{r}\right) (A(\omega_\varepsilon, \xi_2 + \nabla\phi_{\xi_2}) - A(\omega_\varepsilon, \xi_1 + \nabla\phi_{\xi_1})) \cdot (\xi_2 + \nabla\phi_{\xi_2} - \xi_1 - \nabla\phi_{\xi_1}) dx \\ & \quad + \frac{1}{r} \int_{\mathbb{R}^d} (\phi_{\xi_2} - \phi_{\xi_1}) (\nabla\eta)\left(\frac{x}{r}\right) \cdot (A(\omega_\varepsilon, \xi_2 + \nabla\phi_{\xi_2}) - A(\omega_\varepsilon, \xi_1 + \nabla\phi_{\xi_1})) dx \\ & \geq \lambda \int_{\mathbb{R}^d} \eta\left(\frac{x}{r}\right) |\xi_2 + \nabla\phi_{\xi_2} - \xi_1 - \nabla\phi_{\xi_1}|^2 dx \\ & \quad + \frac{1}{r} \int_{\mathbb{R}^d} (\phi_{\xi_2} - \phi_{\xi_1}) (\nabla\eta)\left(\frac{x}{r}\right) \cdot (A(\omega_\varepsilon, \xi_2 + \nabla\phi_{\xi_2}) - A(\omega_\varepsilon, \xi_1 + \nabla\phi_{\xi_1})) dx \end{aligned}$$

which after dividing by $\int_{\mathbb{R}^d} \eta(\frac{x}{r}) dx$, taking the expectation and using stationarity, and passing to the limit $r \rightarrow \infty$ yields by the sublinear growth of the ϕ_ξ

$$\begin{aligned} & (A_{\text{hom}}(\xi_2) - A_{\text{hom}}(\xi_1)) \cdot (\xi_2 - \xi_1) \\ & = \mathbb{E}[(A(\omega, \xi_2 + \nabla\phi_{\xi_2}) - A(\omega, \xi_2 + \nabla\phi_{\xi_2})) \cdot (\xi_2 - \xi_1)] \\ & \geq \lambda |\xi_1 - \xi_2|^2. \end{aligned}$$

Part b. We next show that frame-indifference of the material law – in the sense that $A(\omega_\varepsilon, O\xi) = O^{-1}A(\omega_\varepsilon, \xi)$ for all $x \in \mathbb{R}^d$, almost every ω_ε , all $\xi \in \mathbb{R}^{m \times d}$, and all $O \in \text{SO}(m)$ – is preserved under homogenization. Indeed, if ϕ_ξ solves the corrector equation $\nabla \cdot (A(\omega_\varepsilon, \xi + \nabla\phi_\xi)) = 0$ and if $O \in \text{SO}(m)$, then $O\phi_\xi$ solves the corrector equation $\nabla \cdot (A(\omega_\varepsilon, O\xi + \nabla O\phi_\xi)) = 0$. Using the uniqueness of the corrector up to additive constants, we obtain $\nabla O\phi_\xi = \nabla\phi_{O\xi}$. This entails

$$\begin{aligned} A_{\text{hom}}(O\xi) & = \mathbb{E}[A(\omega_\varepsilon, O\xi + \nabla\phi_{O\xi})] \\ & = \mathbb{E}[A(\omega_\varepsilon, O\xi + \nabla O\phi_\xi)] = \mathbb{E}[O^{-1}A(\omega_\varepsilon, \xi + \nabla\phi_\xi)] \\ & = O^{-1}A_{\text{hom}}(\xi). \end{aligned}$$

Part c. We next show that isotropy of the probability distribution of the material law implies isotropy of the homogenized material law. Let $V \in \text{SO}(d)$. In this case, if ϕ_ξ solves the corrector equation $\nabla \cdot (A(\omega_\varepsilon(x), \xi + \nabla\phi_\xi(x))) = 0$, then the rotated function $\phi_\xi(V \cdot)$ solves the corrector equation $\nabla \cdot (A(\omega_\varepsilon(Vx), \xi VV^{-1} +$

$\nabla(\phi_\xi(Vx))V^{-1})V^{-1}) = 0$ for a rotated monotone operator, i. e. $\tilde{\phi}_{\xi V}(x) := \phi_\xi(Vx)$ is the corrector associated with the rotated operator

$$A_\varepsilon^V(x, \tilde{\xi}) := A(\omega_\varepsilon(Vx), \tilde{\xi}V^{-1})V^{-1}$$

and slope ξV and we have $\nabla\tilde{\phi}_{\xi V}(x) = \nabla\phi_\xi(Vx)V$. This entails by the assumed equality of the laws of $A_\varepsilon(x, \tilde{\xi}) := A(\omega_\varepsilon(x), \tilde{\xi})$ and A_ε^V

$$\begin{aligned} A_{\text{hom}}(\xi V) &= \mathbb{E}[A_\varepsilon(x, \xi V + \nabla\phi_{\xi V}(x))] \\ &= \mathbb{E}[A_\varepsilon^V(x, \xi V + \nabla\tilde{\phi}_{\xi V}(x))] \\ &= \mathbb{E}[A_\varepsilon^V(x, \xi VV^{-1} + \nabla(\phi_\xi(Vx))V^{-1})V^{-1}] \\ &= \mathbb{E}[A_\varepsilon(\omega_\varepsilon(Vx), \xi + \nabla\phi_\xi(Vx))V^{-1}] \\ &= A_{\text{hom}}(\xi)V^{-1}. \end{aligned}$$

□

APPENDIX A. AUXILIARY RESULTS FROM REGULARITY THEORY

We now provide the (standard) proof of the Caccioppoli inequality and the hole-filling estimate for nonlinear elliptic PDEs with monotone nonlinearity from Lemma 34.

Proof of Lemma 34. Let $R > 0$ and let η be a standard cutoff with $\eta \equiv 0$ outside of B_R , $\eta \equiv 1$ in $B_{R/2}$, and $|\nabla\eta| \leq CR^{-1}$. Testing the equation with $\eta^2(u - b)$ for some $b \in \mathbb{R}^m$ to be chosen, we obtain by (A1)–(A2)

$$\begin{aligned} &\lambda \int_{\mathbb{R}^d} \eta^2 |\nabla u|^2 dx + \frac{1}{T} \int_{\mathbb{R}^d} \eta^2 |u|^2 dx \\ &\stackrel{(A1)}{\leq} \int_{\mathbb{R}^d} A(x, \nabla u) \cdot \eta^2 \nabla u dx + \frac{1}{T} \int_{\mathbb{R}^d} \eta^2 |u|^2 dx \\ &\leq - \int_{\mathbb{R}^d} 2\eta(A(x, \nabla u) + g) \cdot (u - b) \otimes \nabla \eta dx + \frac{1}{T} \int_{\mathbb{R}^d} \eta^2 ub dx \\ &\quad + \int_{\mathbb{R}^d} \eta^2 \left(-g \cdot \nabla u + \frac{1}{T} f(u - b) \right) dx \\ &\stackrel{(A2)}{\leq} \frac{C\Lambda}{R} \int_{B_R(x_0) \setminus B_{R/2}(x_0)} \eta(|\nabla u| + |g|)|u - b| dx + \frac{1}{T} \int_{\mathbb{R}^d} \eta^2 ub dx \\ &\quad + \int_{\mathbb{R}^d} \eta^2 \left(-g \cdot \nabla u + \frac{1}{T} f(u - b) \right) dx. \end{aligned}$$

Young's inequality and an absorption argument yields the Caccioppoli-type inequality

$$\begin{aligned} &\int_{B_{R/2}(x_0)} |\nabla u|^2 + \frac{1}{T} |u|^2 dx \\ &\leq \frac{C}{R^2} \int_{B_R(x_0) \setminus B_{R/2}(x_0)} |u - b|^2 dx + \frac{C}{T} \int_{B_R(x_0)} |b|^2 dx + C \int_{B_R(x_0)} |g|^2 + \frac{1}{T} |f|^2 dx \end{aligned}$$

which directly implies (51).

An application of the Poincaré inequality on the annulus $B_R(x_0) \setminus B_{R/2}(x_0)$ in the previous estimate gives upon choosing $b := \int_{B_R(x_0) \setminus B_{R/2}(x_0)} u dx$, using also

Jensen's inequality and the fact that $|B_R(x_0)| \sim |B_R(x_0) \setminus B_{R/2}(x_0)|$ to estimate the second term on the right-hand side,

$$\begin{aligned} & \int_{B_{R/2}(x_0)} |\nabla u|^2 dx + \frac{1}{T} \int_{B_{R/2}(x_0)} |u|^2 dx \\ & \leq C \int_{B_R(x_0) \setminus B_{R/2}(x_0)} |\nabla u|^2 dx + \frac{C}{T} \int_{B_R(x_0) \setminus B_{R/2}(x_0)} |u|^2 dx \\ & \quad + C \int_{B_R(x_0)} |g|^2 + \frac{1}{T} |f|^2 dx. \end{aligned}$$

This in turn yields by the hole-filling argument

$$\begin{aligned} & \int_{B_{R/2}(x_0)} |\nabla u|^2 dx + \frac{1}{T} \int_{B_{R/2}(x_0)} |u|^2 dx \\ & \leq \theta \left(\int_{B_R(x_0)} |\nabla u|^2 dx + \frac{1}{T} \int_{B_R(x_0)} |u|^2 dx \right) + C \int_{B_R(x_0)} |g|^2 + \frac{1}{T} |f|^2 dx \end{aligned}$$

for $\theta = \frac{C}{C+1}$. Iterating this estimate with R replaced by $2^{-k}R$, we deduce our desired estimate (52) if r is of the form $r = 2^{-K}R$. For other values of r , we may simply use the already-established inequality for the next bigger radius of the form $r = 2^{-K}R$ and increase the constant C if necessary. \square

We next provide a small-scale Hölder regularity result for the linearized corrector $\phi_{\xi, \Xi}^T$.

Proposition 41. *Let the assumptions (A1)–(A3) and (P1)–(P2) as well as (R) be in place. Then there exists $\alpha > 0$ such that for all $\xi, \Xi \in \mathbb{R}^{m \times d}$ and all $T \geq \varepsilon^2$ the gradient of the linearized corrector $\phi_{\xi, \Xi}^T$ is subject to a Hölder regularity estimate of the form*

$$\begin{aligned} & |(\Xi + \nabla \phi_{\xi, \Xi}^T)(x) - (\Xi + \nabla \phi_{\xi, \Xi}^T)(y)| \\ & \leq \mathcal{C}(x_0, \xi, T) (1 + |\xi|)^C \left(\frac{|x - y|}{\varepsilon} \right)^\alpha \left(\int_{B_\varepsilon(x_0)} |\Xi + \nabla \phi_{\xi, \Xi}^T|^2 dx + |\Xi|^2 \right)^{1/2} \end{aligned}$$

for all $x, y \in B_{\varepsilon/2}(x_0)$, where $\mathcal{C}(x_0, \xi, T)$ denotes a stationary random field with stretched exponential moment bounds $\mathbb{E}[\exp(\mathcal{C}^{1/C})] \leq 2$ for some constant $C = C(d, m, \lambda, \Lambda, \rho, \nu)$.

Proof. The result is a straightforward consequence of classical Schauder theory (see e. g. the proof of [27, Theorem 5.19]) applied to the equation

$$-\nabla \cdot (\partial_\xi A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T)(\Xi + \nabla \phi_{\xi, \Xi}^T)) + \frac{1}{T} (\Xi \cdot (x - x_0) + \phi_{\xi, \Xi}^T) = \frac{1}{T} \Xi \cdot (x - x_0),$$

which is possible by the Hölder continuity of the coefficient $\partial_\xi A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T)$ on $B_\varepsilon(x_0)$, which in turn may be deduced from Proposition 42, our regularity assumptions on ω_ε (see (R)), and the Lipschitz dependence of $\partial_\xi A$ on both variables (see (A3) and (R)). \square

Proposition 42. *Let the assumptions (A1)–(A3) and (P1)–(P2) as well as (R) be in place. Then there exists $\alpha > 0$ such that for all $\xi \in \mathbb{R}^{m \times d}$ and all $T \geq \varepsilon^2$ the*

gradient of the corrector ϕ_ξ^T is subject to a Hölder regularity estimate of the form

$$|(\xi + \nabla \phi_\xi^T)(x) - (\xi + \nabla \phi_\xi^T)(y)| \leq C(x_0, \xi, T)(1 + |\xi|) \left(\frac{|x - y|}{\varepsilon} \right)^\alpha$$

for all $x, y \in B_{\varepsilon/2}(x_0)$, where C denotes a stationary random field with stretched exponential moment bounds $\mathbb{E}[\exp(C^{1/C})] \leq 2$ for some $C = C(d, m, \lambda, \Lambda, \rho, \nu)$.

Proof. We differentiate the equation $-\nabla \cdot (A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T)) + \frac{1}{T} \phi_\xi^T = 0$. This yields

$$(124) \quad -\nabla \cdot (\partial_\xi A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \nabla \partial_i \phi_\xi^T) + \frac{1}{T} \partial_i \phi_\xi^T = \nabla \cdot (\partial_\omega \partial_\xi A(\omega_\varepsilon, \xi + \nabla \phi_\xi^T) \partial_i \omega_\varepsilon).$$

By (A1)–(A3), this is a linear elliptic system for the derivative $\partial_i \phi_\xi^T$ with uniformly elliptic and bounded coefficient field.

Case a: Two-dimensional systems with smooth coefficients. Meyers' estimate for the PDE (124) in the form of Lemma 44 with $T := \infty$ and the condition (A3) imply for some $p > 2$ and any $b \in \mathbb{R}^m$

$$\begin{aligned} & \left(\int_{B_{\varepsilon/2}(x_0)} |\nabla \partial_i \phi_\xi^T|^p dx \right)^{1/p} \\ & \leq C \left(\int_{B_\varepsilon(x_0)} |\nabla \partial_i \phi_\xi^T|^2 dx \right)^{1/2} + C \left(\int_{B_\varepsilon(x_0)} |\nabla \omega_\varepsilon|^p + \left| \frac{1}{T} (\phi_\xi^T - b) \right|^p dx \right)^{1/p} \end{aligned}$$

Using the Caccioppoli inequality (51) with $T = \infty$ for the PDE (124), choosing $p - 2 > 0$ small enough, and using the Poincaré inequality, we get by $T \geq \varepsilon^2$

$$\begin{aligned} & \left(\int_{B_{\varepsilon/2}(x_0)} |\nabla \partial_i \phi_\xi^T|^p dx \right)^{1/p} \\ & \leq \frac{C}{\varepsilon} \left(\int_{B_{2\varepsilon}(x_0)} |\nabla \phi_\xi^T|^2 dx \right)^{1/2} + C \left(\int_{B_{2\varepsilon}(x_0)} |\nabla \omega_\varepsilon|^p dx \right)^{1/p}. \end{aligned}$$

By our estimate (88), Lemma 19, and the bound on $\nabla \omega_\varepsilon$ in (R), the right-hand side may be bounded by $C\varepsilon^{-1}(|\xi| + 1)$ for some random constant C with stretched exponential moments. By Morrey's embedding, we obtain the desired estimate.

Case b: Scalar equations and systems with Uhlenbeck structure with smooth coefficients.

In the case of a scalar equation, we infer the desired Hölder continuity of $\partial_i \phi_\xi^T$ from De Giorgi-Nash-Moser theory: Applying [28, Theorem 8.24] to the equation (124), we deduce

$$\begin{aligned} & \varepsilon^{\alpha-1} \sup_{x_1, x_2 \in B_{\varepsilon/2}(x_0)} \frac{|\partial_i \phi_\xi^T(x_1) - \partial_i \phi_\xi^T(x_2)|}{|x_1 - x_2|^\alpha} \\ & \leq C\varepsilon^{-1} \left(\int_{B_\varepsilon(x_0)} |\partial_i \phi_\xi^T|^2 dx \right)^{1/2} + C \left(\int_{B_\varepsilon(x_0)} |\nabla \omega_\varepsilon|^{2d} dx \right)^{1/2d}. \end{aligned}$$

Using our regularity assumption on ω_ε from (R) and again (88) and Lemma 19, we conclude.

In the systems' case, one replaces the De Giorgi-Nash-Moser theory by Uhlenbeck's regularity result [41]. \square

APPENDIX B. QUALITATIVE DIFFERENTIABILITY OF CORRECTORS

We now provide the proof of the qualitative differentiability results for the homogenization correctors that we have used throughout the present work.

Proof of Lemma 20. Let $h > 0$. Subtracting the equations for $\phi_{\xi+h\Xi}^T$ and ϕ_{ξ}^T , we obtain

$$(125) \quad -\nabla \cdot ((A(\tilde{\omega}, \xi + h\Xi + \nabla\phi_{\xi+h\Xi}^T) - A(\tilde{\omega}, \xi + \nabla\phi_{\xi}^T)) + \frac{1}{T}(\phi_{\xi+h\Xi}^T - \phi_{\xi}^T)) = 0$$

which we may rewrite as

$$\begin{aligned} & -\nabla \cdot ((A(\tilde{\omega}, \xi + \nabla\phi_{\xi+h\Xi}^T) - A(\tilde{\omega}, \xi + \nabla\phi_{\xi}^T)) + \frac{1}{T}(\phi_{\xi+h\Xi}^T - \phi_{\xi}^T)) \\ & = \nabla \cdot (A(\tilde{\omega}, \xi + h\Xi + \nabla\phi_{\xi+h\Xi}^T) - A(\tilde{\omega}, \xi + \nabla\phi_{\xi+h\Xi}^T)). \end{aligned}$$

Note that by (A2) we have $|A(\tilde{\omega}, \xi + h\Xi + \nabla\phi_{\xi+h\Xi}^T) - A(\tilde{\omega}, \xi + \nabla\phi_{\xi+h\Xi}^T)| \leq C|\xi|$. Lemma 36 yields

$$\begin{aligned} & \int_{\mathbb{R}^d} \left(|\nabla\phi_{\xi+h\Xi}^T - \nabla\phi_{\xi}^T|^2 + \frac{1}{T}|\phi_{\xi+h\Xi}^T - \phi_{\xi}^T|^2 \right) \exp(-c|x|/\sqrt{T}) dx \\ & \leq C \int_{\mathbb{R}^d} h^2 |\Xi|^2 \exp(-c|x|/\sqrt{T}) dx \leq Ch^2 |\Xi|^2 \sqrt{T}^d \end{aligned}$$

and Meyers' estimate (see Lemma 44) with a dyadic decomposition of \mathbb{R}^d into $B_{\sqrt{T}}$ and $B_{2^k\sqrt{T}} \setminus B_{2^{k-1}\sqrt{T}}$ for $k \in \mathbb{N}$ upgrades this to

$$\int_{\mathbb{R}^d} |\nabla\phi_{\xi+h\Xi}^T - \nabla\phi_{\xi}^T|^{2p} \exp(-c|x|/\sqrt{T}) dx \leq Ch^{2p} |\Xi|^{2p} \sqrt{T}^d$$

for some $p > 1$.

Adding a multiple of the PDE for the linearized corrector $\phi_{\xi,\Xi}^T$ to (125), we deduce

$$\begin{aligned} & -\nabla \cdot ((A(\tilde{\omega}, \xi + h\Xi + \nabla\phi_{\xi+h\Xi}^T) - A(\tilde{\omega}, \xi + \nabla\phi_{\xi}^T)) - h\partial_{\xi}A(\tilde{\omega}, \xi + \nabla\phi_{\xi}^T)\nabla\phi_{\xi,\Xi}^T) \\ & \quad + \frac{1}{T}(\phi_{\xi+h\Xi}^T - \phi_{\xi}^T - h\phi_{\xi,\Xi}^T) = 0. \end{aligned}$$

Using Taylor expansion, the uniform bound $|\partial_{\xi}^2 A| \leq \Lambda$ from (R), and the Lipschitz estimate for A from (A2), we obtain for any $\delta \in (0, 1]$

$$-\nabla \cdot (\partial_{\xi}A(\tilde{\omega}, \xi + \nabla\phi_{\xi}^T)(h\Xi + \nabla\phi_{\xi+h\Xi}^T - \nabla\phi_{\xi}^T)) + \frac{1}{T}(\phi_{\xi+h\Xi}^T - \phi_{\xi}^T) = \nabla \cdot R$$

with $|R| \leq Ch^{1+\delta}|\Xi|^{1+\delta} + C|\nabla\phi_{\xi+h\Xi}^T - \nabla\phi_{\xi}^T|^{1+\delta}$. Choosing δ such that $1 + \delta < p$ and applying Lemma 36, this finally yields the estimate

$$\int_{\mathbb{R}^d} |\nabla\phi_{\xi+h\Xi}^T - \nabla\phi_{\xi}^T - h\nabla\phi_{\xi,\Xi}^T|^2 \exp(-c|x|/\sqrt{T}) dx \leq Ch^{2+2\delta} \sqrt{T}^d.$$

The proof of the corresponding result for σ_{ξ} is even easier, as the equation for σ_{ξ} is linear in $q_{\xi}^T = A(\tilde{\omega}, \xi + \nabla\phi_{\xi}^T)$. \square

APPENDIX C. MEYERS' ESTIMATE FOR ELLIPTIC EQUATIONS WITH MASSIVE TERM

We recall Gehring's lemma in the following form.

Lemma 43 (see e.g. [27, Theorem 6.38]). *Let $K > 0$, $m \in (0, 1)$, $s > 1$ and $B = B_R(x_0)$ for some $x_0 \in \mathbb{R}^d$ and $R > 0$ be given. Suppose that $f \in L^1(B)$ and $g \in L^s(B)$ are such that for every $z \in \mathbb{R}^d$ and $r > 0$ with $B_r(z) \subset B$ it holds that*

$$\int_{B_{r/2}(z)} |f| dx \leq K \left(\int_{B_r(z)} |f|^m dx \right)^{1/m} + \int_{B_r(z)} |g| dx.$$

Then there exist $q = q(K, m) \in (1, s]$ and $C = C(K, m) \in [1, \infty)$ such that $f \in L^q(B_{R/2}(x_0))$ and the estimate

$$\left(\int_{B_{R/2}(z)} |f|^q dx \right)^{1/q} \leq C \int_{B_R(z)} |f| dx + C \left(\int_{B_R(z)} |g|^q dx \right)^{1/q}.$$

holds.

Lemma 44 (Meyers estimate for PDEs with massive term). *Let $d, m \in \mathbb{N}$, $T > 0$, and let $a : \mathbb{R}^d \rightarrow (\mathbb{R}^{m \times d} \otimes \mathbb{R}^{m \times d})$ be a uniformly elliptic and bounded coefficient field with ellipticity and boundedness constants $0 < \lambda \leq \Lambda < \infty$. Let $f, g \in L^2(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$ and let $v \in H^1(\mathbb{R}^d; \mathbb{R}^m)$ be the (unique) weak solution to the linear system*

$$-\nabla \cdot (a \nabla v) + \frac{1}{T} v = \nabla \cdot g + \frac{1}{\sqrt{T}} f \quad \text{on } \mathbb{R}^d.$$

Then there exists $p_0 = p_0(d, m, \lambda, \Lambda) > 2$ such that for all $2 \leq p < p_0$, any $x_0 \in \mathbb{R}^d$, and any $R > 0$ we have

$$\begin{aligned} & \int_{B_R(x_0)} |\nabla v|^p + \left| \frac{1}{\sqrt{T}} v \right|^p dx \\ & \leq C(d, m, \lambda, \Lambda, p) \int_{B_{2R}(x_0)} |g|^p + |f|^p dx \\ & \quad + C(d, m, \lambda, \Lambda, p) \left(\int_{B_{2R}(x_0)} |\nabla v|^2 + \left| \frac{1}{\sqrt{T}} v \right|^2 dx \right)^{p/2}. \end{aligned}$$

Proof. Since a is uniformly elliptic and bounded, we deduce by the Caccioppoli inequality (51) for any $b \in \mathbb{R}^m$, and $r > 0$, and any $z \in \mathbb{R}^d$

$$\int_{B_{r/2}(z)} |\nabla v|^2 + \frac{1}{T} |v|^2 dx \leq C(d, m, \lambda, \Lambda) \int_{B_r(z)} r^{-2} |v - b|^2 + \frac{1}{T} |b|^2 + |g|^2 + |f|^2 dx.$$

Choosing $b := \int_{B_r} v dx$ and using the Poincaré-Sobolev inequality as well as Jensen's inequality, we deduce

$$\begin{aligned} & \int_{B_{r/2}(z)} |\nabla v|^2 + \frac{1}{T} |v|^2 dx \\ & \leq C(d, m, \lambda, \Lambda) \left(\int_{B_r(z)} |\nabla v|^{2d/(d+2)} + \left| \frac{1}{\sqrt{T}} v \right|^{2d/(d+2)} dx \right)^{(d+2)/d} \\ & \quad + C(d, m, \lambda, \Lambda) \int_{B_r(z)} |g|^2 + |f|^2 dx. \end{aligned}$$

Lemma 43 now yields the desired estimate. \square

Proof of Lemma 35. Let $R > 0$. We split v as $v = v_{out} + \sum_{k=1}^{\infty} v_{in,k}$, where $v_{in,k} \in H^1(\mathbb{R}^d; \mathbb{R}^m)$ is the unique weak solution to the PDE

$$(126a) \quad -\nabla \cdot (a \nabla v_{in,k}) + \frac{1}{T} v_{in,k} = \nabla \cdot (g \chi_{B_{2^{-k}R}(x_0) \setminus B_{2^{-k-1}R}(x_0)}) + \frac{1}{\sqrt{T}} g \chi_{B_{2^{-k}R}(x_0) \setminus B_{2^{-k-1}R}(x_0)})$$

and where $v_{out} \in H^1(\mathbb{R}^d; \mathbb{R}^m)$ is the unique weak solution to the PDE

$$(126b) \quad -\nabla \cdot (a \nabla v_{out}) + \frac{1}{T} v_{out} = \nabla \cdot (g \chi_{\mathbb{R}^d \setminus B_{R/2}(x_0)}) + \frac{1}{\sqrt{T}} g \chi_{\mathbb{R}^d \setminus B_{R/2}(x_0)}.$$

Passing to the limit $R \rightarrow \infty$ in the hole-filling estimate (52) for v_{out} and inserting $4R$ in place of r in the resulting bound, we deduce for any $\delta > 0$ with $\delta < c$

$$(127) \quad \int_{B_{4R}(x_0)} |\nabla v_{out}|^2 + \frac{1}{T} |v_{out}|^2 dx \leq C \int_{\mathbb{R}^d \setminus B_R(x_0)} \left(\frac{R}{R + |x - x_0|} \right)^\delta (|g|^2 + |f|^2) dx.$$

This yields by dividing by $c(d)R^d$ and applying Hölder's inequality

$$\begin{aligned} & \int_{B_{4R}(x_0)} |\nabla v_{out}|^2 + \frac{1}{T} |v_{out}|^2 dx \\ & \leq C(d, m, \lambda, \Lambda, p, \gamma) \left(\int_{\mathbb{R}^d \setminus B_R(x_0)} R^{-d} \left(\frac{R}{R + |x - x_0|} \right)^{\delta p/2 - \gamma} (|g|^p + |f|^p) dx \right)^{2/p} \end{aligned}$$

for any γ with $d(p-2)/2 < \gamma < \delta$. Plugging this bound into Lemma 44, we obtain

$$(128) \quad \begin{aligned} & \int_{B_{2R}(x_0)} |\nabla v_{out}|^p + \left| \frac{1}{\sqrt{T}} v_{out} \right|^p dx \\ & \leq C(d, m, \lambda, \Lambda, p) \int_{B_{4R}(x_0) \setminus B_R(x_0)} |g|^p + |f|^p dx \\ & \quad + C(d, m, \lambda, \Lambda, p, \gamma) \int_{\mathbb{R}^d \setminus B_R(x_0)} \left(\frac{R}{R + |x - x_0|} \right)^{\delta p/2 - \gamma} (|g|^p + |f|^p) dx \\ & \leq C(d, m, \lambda, \Lambda, p, \gamma) \int_{\mathbb{R}^d \setminus B_R(x_0)} \left(\frac{R}{R + |x - x_0|} \right)^{\delta p/2 - \gamma} (|g|^p + |f|^p) dx. \end{aligned}$$

We next estimate the contributions of $v_{in,k}$. We have

$$\int_{B_{4R}(x_0) \setminus B_{R/2}(x_0)} |\nabla v_{in,k}|^2 + \frac{1}{T} |v_{in,k}|^2 dx \leq 2 \left| \sup_{\tilde{g}, \tilde{f}} \int_{\mathbb{R}^d} \tilde{g} \cdot \nabla v_{in,k} + \frac{1}{\sqrt{T}} v_{in,k} \tilde{f} dx \right|^2$$

where the supremum runs over all functions \tilde{g} and \tilde{f} with $\text{supp } \tilde{g} \cup \text{supp } \tilde{f} \subset B_{2R}(x_0) \setminus B_R(x_0)$ and $\int_{\mathbb{R}^d} |\tilde{g}|^2 + |\tilde{f}|^2 dx \leq 1$. Denoting by $w \in H^1(\mathbb{R}^d; \mathbb{R}^m)$ the unique solution to the dual PDE

$$-\nabla \cdot (a^* \nabla w) + \frac{1}{T} w = -\nabla \cdot \tilde{g} + \frac{1}{\sqrt{T}} \tilde{f},$$

we obtain

$$\begin{aligned}
& \int_{B_{4R}(x_0) \setminus B_{R/2}(x_0)} |\nabla v_{in,k}|^2 + \frac{1}{T} |v_{in,k}|^2 dx \\
& \leq 2 \left| \sup_{\tilde{g}, \tilde{f}} \int_{\mathbb{R}^d} a \nabla v_{in,k} \cdot \nabla w + \frac{1}{T} v_{in,k} \cdot w dx \right|^2 \\
& \leq 2 \sup_{\tilde{g}, \tilde{f}} \left| \int_{B_{2^{-k}R}(x_0) \setminus B_{2^{-k-1}R}(x_0)} g \cdot \nabla w - \frac{1}{\sqrt{T}} f w dx \right|^2 \\
& \leq 2 \int_{B_{2^{-k}R}(x_0) \setminus B_{2^{-k-1}R}(x_0)} |g|^2 + |f|^2 dx \\
& \quad \times \sup_{\tilde{g}, \tilde{f}} \int_{B_{2^{-k}R}(x_0) \setminus B_{2^{-k-1}R}(x_0)} |\nabla w|^2 + \frac{1}{T} |w|^2 dx.
\end{aligned}$$

By the hole-filling estimate for w in the form analogous to (127), we deduce

$$\begin{aligned}
& \int_{B_{2^{-k}R}(x_0) \setminus B_{2^{-k-1}R}(x_0)} |\nabla w|^2 + \frac{1}{T} |w|^2 dx \\
& \leq C \int_{\mathbb{R}^d} \left(\frac{2^{-k}R}{2^{-k}R + |x - x_0|} \right)^\delta (|\tilde{g}|^2 + |\tilde{f}|^2) dx \leq C
\end{aligned}$$

which entails by the bounds on \tilde{g} and \tilde{f}

$$\begin{aligned}
& \int_{B_{4R}(x_0) \setminus B_{R/2}(x_0)} |\nabla v_{in,k}|^2 + \frac{1}{T} |v_{in,k}|^2 dx \\
& \leq C(2^{-k})^\delta \int_{B_{2^{-k}R}(x_0) \setminus B_{2^{-k-1}R}(x_0)} |g|^2 + |f|^2 dx.
\end{aligned}$$

An application of Lemma 44 to (126a) yields by the preceding estimate and Jensen's inequality

$$\begin{aligned}
& \int_{B_{2R}(x_0) \setminus B_R(x_0)} |\nabla v_{in,k}|^p + \frac{1}{T} |v_{in,k}|^p dx \\
& \leq C(2^{-k})^{\delta p/2 - d(p-2)} \int_{B_{2^{-k}R}(x_0) \setminus B_{2^{-k-1}R}(x_0)} |g|^p + |f|^p dx.
\end{aligned}$$

Taking the sum in k and adding (128), we deduce by choosing $\gamma > 0$ small enough and by requiring p to be close enough to 2 (depending on $\tilde{\delta}$)

$$\begin{aligned}
& \int_{B_{4R}(x_0) \setminus B_{R/2}(x_0)} |\nabla v|^p + \left| \frac{1}{\sqrt{T}} v \right|^p dx \\
& \leq C \int_{B_R(x_0)} \left(\frac{|x - x_0|}{R} \right)^{p\tilde{\delta}/3} (|g|^p + |f|^p) dx \\
& \quad + C \int_{\mathbb{R}^d \setminus B_R(x_0)} \left(\frac{R}{|x - x_0|} \right)^{p\tilde{\delta}/3} (|g|^p + |f|^p) dx.
\end{aligned}$$

Multiplying both sides by $(\frac{R}{r})^{\alpha_0}$, taking the sum over all dyadic $R = 2^l r$, and using a standard Meyers estimate on the ball $B_r(x_0)$, we obtain

$$\begin{aligned} & \int_{\mathbb{R}^d} \left(|\nabla v_{in}|^2 + \frac{1}{T} |v_{in}|^2 \right) \left(1 + \frac{|x - x_0|}{r} \right)^{\alpha_0} dx \\ & \leq C \sum_{l=1}^{\infty} \int_{B_{2^l r}(x_0)} \left(\frac{|x - x_0|}{2^l r} \right)^{p\delta/3} (2^l)^{\alpha_0} (|g|^p + |f|^p) dx \\ & \quad + C \sum_{l=1}^{\infty} \int_{\mathbb{R}^d \setminus B_{2^l r}(x_0)} \left(\frac{2^l r}{|x - x_0|} \right)^{p\delta/3} (2^l)^{\alpha_0} (|g|^p + |f|^p) dx. \end{aligned}$$

For $\alpha_1 > \alpha_0$, we may estimate the last sum as

$$\begin{aligned} & \sum_{l=1}^{\infty} \int_{\mathbb{R}^d \setminus B_{2^l r}(x_0)} \left(\frac{2^l r}{|x - x_0|} \right)^{p\delta/3} (2^l)^{\alpha_0} (|g|^p + |f|^p) dx \\ & \leq C(\delta) \sum_{l=1}^{\infty} \sum_{n=l}^{\infty} \int_{B_{2^{n+1}r}(x_0) \setminus B_{2^n r}(x_0)} (2^{l-n})^{p\delta/3} (2^l)^{\alpha_0} (|g|^p + |f|^p) dx \\ & \leq C(\delta, \alpha_0, \alpha_1) \sum_{n=1}^{\infty} \int_{B_{2^{n+1}r}(x_0) \setminus B_{2^n r}(x_0)} (2^l)^{\alpha_1} (|g|^p + |f|^p) dx. \end{aligned}$$

If $\alpha_0 > 0$ is chosen small enough, the previous two estimates imply (53). \square

REFERENCES

- [1] S. Armstrong and P. Cardaliaguet. Stochastic homogenization of quasilinear Hamilton-Jacobi equations and geometric motions. *J. Eur. Math. Soc. (JEMS)*, 20(4):797–864, 2018.
- [2] S. Armstrong, S. Ferguson, and T. Kuusi. Homogenization, linearization and large-scale regularity for nonlinear elliptic equations. *Preprint*, 2018. arXiv:1805.00467.
- [3] S. Armstrong, S. Ferguson, and T. Kuusi. tba. *in preparation*, 2019.
- [4] S. Armstrong, T. Kuusi, and J.-C. Mourrat. The additive structure of elliptic homogenization. *Invent. Math.*, 208(3):999–1154, 2017.
- [5] S. Armstrong, T. Kuusi, and J.-C. Mourrat. *Quantitative stochastic homogenization and large-scale regularity*. Preprint, 2017. arXiv:1705.05300.
- [6] S. N. Armstrong, P. Cardaliaguet, and P. E. Souganidis. Error estimates and convergence rates for the stochastic homogenization of Hamilton-Jacobi equations. *J. Amer. Math. Soc.*, 27(2):479–540, 2014.
- [7] S. N. Armstrong and J.-C. Mourrat. Lipschitz regularity for elliptic equations with random coefficients. *Arch. Ration. Mech. Anal.*, 219(1):255–348, 2016.
- [8] S. N. Armstrong and C. K. Smart. Quantitative stochastic homogenization of elliptic equations in nondivergence form. *Arch. Ration. Mech. Anal.*, 214(3):867–911, 2014.
- [9] S. N. Armstrong and C. K. Smart. Quantitative stochastic homogenization of convex integral functionals. *Ann. Sci. Éc. Norm. Supér. (4)*, 49(2):423–481, 2016.
- [10] M. Avellaneda and F.-H. Lin. Compactness methods in the theory of homogenization. *Comm. Pure Appl. Math.*, 40(6):803–847, 1987.
- [11] J. M. Ball. Convexity conditions and existence theorems in nonlinear elasticity. *Arch. Rational Mech. Anal.*, 63(4):337–403, 1976/77.
- [12] M. Barchiesi and A. Gloria. New counterexamples to the cell formula in nonconvex homogenization. *Arch. Ration. Mech. Anal.*, 195(3):991–1024, 2010.
- [13] P. Bella, B. Fehrman, J. Fischer, and F. Otto. Stochastic homogenization of linear elliptic equations: Higher-order error estimates in weak norms via second-order correctors. *SIAM J. Math. Anal.*, 49(6):4658–4703, 2017.
- [14] M. Briane and G. A. Francfort. Loss of ellipticity through homogenization in linear elasticity. *Math. Models Methods Appl. Sci.*, 25(5):905–928, 2015.

- [15] L. A. Caffarelli and P. E. Souganidis. Rates of convergence for the homogenization of fully nonlinear uniformly elliptic pde in random media. *Invent. Math.*, 180(2):301–360, 2010.
- [16] S. Conti, G. Dolzmann, B. Kirchheim, and S. Müller. Sufficient conditions for the validity of the Cauchy-Born rule close to $SO(n)$. *J. Eur. Math. Soc.*, 8(3):515–530, 2006.
- [17] G. Dal Maso and L. Modica. Nonlinear stochastic homogenization. *Ann. Mat. Pura Appl. (4)*, 144:347–389, 1986.
- [18] G. Dal Maso and L. Modica. Nonlinear stochastic homogenization and ergodic theory. *J. Reine Angew. Math.*, 368:28–42, 1986.
- [19] M. Duerinckx and A. Gloria. Weighted functional inequalities: Concentration properties. *Preprint*, 2017. arXiv:1711.03148.
- [20] M. Duerinckx and A. Gloria. Weighted functional inequalities: Constructive approach. *Preprint*, 2017. arXiv:1711.03152.
- [21] M. Duerinckx, S. Neukamm, M. Ruf, and M. Schäffner. Quantitative stochastic homogenization in nonlinear elasticity at small loads. *in preparation*, 2019.
- [22] J. Fischer. The choice of representative volumes in the approximation of effective properties of random materials. *Preprint*, 2018. arXiv:1807.00834.
- [23] J. Fischer and C. Raithel. Liouville principles and a large-scale regularity theory for random elliptic operators on the half-space. *SIAM J. Math. Anal.*, 49(1):82–114, 2017.
- [24] G. A. Francfort and A. Gloria. Isotropy prohibits the loss of strong ellipticity through homogenization in linear elasticity. *C. R. Math. Acad. Sci. Paris*, 354(11):1139–1144, 2016.
- [25] G. Friesecke and F. Theil. Validity and failure of the Cauchy-Born hypothesis in a two-dimensional mass-spring lattice. *J. Nonlinear Sci.*, 12(5):445–478, 2002.
- [26] G. Geymonat, S. Müller, and N. Triantafyllidis. Homogenization of non-linearly elastic materials, microscopic bifurcation and macroscopic loss of rank-one convexity. *Arch. Rational Mech. Anal.*, 122(3):231–290, 1993.
- [27] M. Giaquinta and L. Martinazzi. *An introduction to the regularity theory for elliptic systems, harmonic maps and minimal graphs*, volume 11. Edizioni della Normale, Pisa, second edition, 2012.
- [28] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Springer, 2001.
- [29] A. Gloria, S. Neukamm, and F. Otto. A regularity theory for random elliptic operators. *Preprint*, 2014. arXiv:1409.2678.
- [30] A. Gloria, S. Neukamm, and F. Otto. Quantification of ergodicity in stochastic homogenization: optimal bounds via spectral gap on Glauber dynamics. *Invent. Math.*, 199(2):455–515, 2015.
- [31] A. Gloria, S. Neukamm, and F. Otto. Quantitative estimates in stochastic homogenization for correlated coefficient fields. *in preparation*, 2019.
- [32] A. Gloria and F. Otto. An optimal variance estimate in stochastic homogenization of discrete elliptic equations. *Ann. Probab.*, 39(3):779–856, 2011.
- [33] A. Gloria and F. Otto. An optimal error estimate in stochastic homogenization of discrete elliptic equations. *Ann. Appl. Probab.*, 22(1):1–28, 2012.
- [34] A. Gloria and F. Otto. The corrector in stochastic homogenization: Near-optimal rates with optimal stochastic integrability. *Preprint*, 2015. arXiv:1510.08290.
- [35] A. Gloria and M. Ruf. Loss of strong ellipticity through homogenization in 2D linear elasticity: a phase diagram. *Arch. Ration. Mech. Anal.*, 231(2):845–886, 2019.
- [36] S. Gutiérrez. Laminations in linearized elasticity: the isotropic non-very strongly elliptic case. *J. Elasticity*, 53(3):215–256, 1998/99.
- [37] H. Mitake, H. V. Tran, and Y. Yu. Rate of convergence in periodic homogenization of Hamilton-Jacobi equations: the convex setting. *Arch. Ration. Mech. Anal.*, 233(2):901–934, 2019.
- [38] A. Naddaf and T. Spencer. Estimates on the variance of some homogenization problems. *Unpublished preprint*, 1998.
- [39] S. Neukamm and M. Schäffner. Quantitative homogenization in nonlinear elasticity for small loads. *Arch. Ration. Mech. Anal.*, 230(1):343–396, 2018.
- [40] S. Neukamm and M. Schäffner. Lipschitz estimates and existence of correctors for nonlinearly elastic, periodic composites subject to small strains. *Calc. Var. Partial Differential Equations*, 58(2):Art. 46, 51, 2019.

- [41] K. Uhlenbeck. Regularity for a class of non-linear elliptic systems. *Acta Math.*, 138(3-4):219–240, 1977.
- [42] L. Wang, Q. Xu, and P. Zhao. Convergence rates on periodic homogenization of p-Laplace type equations. *Preprint*, 2018. arXiv:1812.04837.
- [43] L. Wang, Q. Xu, and P. Zhao. Quantitative estimates on periodic homogenization of nonlinear elliptic operators. *Preprint*, 2018. arXiv:1807.10865.
- [44] K. Zhang. Energy minimizers in nonlinear elastostatics and the implicit function theorem. *Arch. Rational Mech. Anal.*, 114(2):95–117, 1991.

IST AUSTRIA, AM CAMPUS 1, 3400 KLOSTERNEUBURG, AUSTRIA

TU DRESDEN, FACULTY OF MATHEMATICS, 01062 DRESDEN